

CRUDE DRUGS FROM AQUATIC PLANTS. VII.¹ FOUR NEW THIASPIRANE SULFOXIDE TYPE NUPHAR ALKALOIDS, NUPHARPUMILAMINES A, B, C, AND D, FROM CHINESE NUPHARIS RHIZOMA, THE RHIZOMA OF *NUPHAR PUMILUM* (TIMM.) DC. (NYMPHACEAE)[†]

Masayuki Yoshikawa,* Toshiyuki Murakami, Atsushi Ishikado, Shuji Wakao, Nobutoshi Murakami, Johji Yamahara, and Hisashi Matsuda

Kyoto Pharmaceutical University, 5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto 607, Japan

Abstract — Following the characterization of thiohemiaminal type nuphar alkaloids with immunosuppressive activity, four new thiaspirane sulfoxide type dimeric sesquiterpene alkaloids, nupharpumilamines A, B, C, and D, were isolated from Chinese Nupharis Rhizoma, the dried rhizoma of *Nuphar pumilum* (TIMM.) DC. The structures of nupharpumilamines A, B, C, and D were determined on the basis of chemical and physicochemical evidence.

In the course of our studies for the bioactive constituents of natural medicines originating from aquatic plants,² we found that the alkaloid fraction from Chinese Nupharis Rhizoma, the dried rhizoma of *Nuphar pumilum* (TIMM.) DC. (Nymphaeaceae), showed potent inhibitory activity on the *anti*-sheep erythrocyte plaque forming cell formation in mice spleen cells. From the alkaloid fraction, four thiohemiaminal type nuphar alkaloids, 6-hydroxythiobinupharidine (5), 6,6'-dihydroxythiobinupharidine (6), 6-hydroxythionupharlutine B (7), and 6'-hydroxythionupharlutine B (8), were isolated as the immunosuppressive principles together with inactive nine known nuphar alkaloids. Furthermore, by examination of the structure requirement for the immunosuppressive activity, the thiohemiaminal structure of those alkaloids was found to be essential to the activity.^{1,3} As a continuing study, we have isolated four new thiaspirane sulfoxide type dimeric sesquiterpene alkaloids named nupharpumilamines A (1), B (2), C (3), and D (4) from Chinese Nupharis Rhizoma. In this paper, we describe the isolation and structure elucidation of nupharpumilamines A (1)—D (4).⁴

The alkaloid fraction from Chinese Nupharis Rhizoma^{1,3} was subjected to normal silica gel and Chromatorex NH column chromatography and finally hplc (Develosil ODS HG-5, MeOH-H₂O-Et₂NH, MeCN-H₂O-Et₂NH) to afford 1 (0.0001%), 2 (0.0002%), 3 (0.0002%), and 4 (0.0005%).

Nupharpumilamine A (1)

Nupharpumilamine A (1) was obtained as colorless oil with negative optical rotation ($[\alpha]_D^{25}$ -33.4°). The IR spectrum of 1 showed absorption bands at 3360, 1024, and 874 cm⁻¹ ascribable to hydroxyl, sulfinyl, and furan functions and Bohlmann absorptions⁵ at 2750—2900 cm⁻¹ due to the *trans*-quinolizidine rings. The molecular formula C₃₀H₄₂N₂O₄S of 1 was

[†] Dedicated to the memory of the late Professor Shun-ichi Yamada





Figure 1

confirmed from the EI-MS and FAB-MS and by high-resolution MS measurement. Namely, the EI-MS and positive-ion FAB-MS of **1** showed dehydration peaks at m/z 508 ($M^+ - H_2O$) and 509 ($M + H - H_2O$)⁺, respectively, while a quasimolecular ion peak was observed at m/z 525 ($M - H$)⁻ in the negative-ion FAB-MS of **1**. Furthermore, fragment ion peaks characteristic of thiohemiaminal type nuphar alkaloid⁶ were observed at m/z 491, 447, 230, 178, and 94 in the EI-MS of **1**. The ¹H-NMR (CD₃OD) and ¹³C-NMR (Table 1) spectra⁷ of **1** showed two tertiary methyls [δ 0.92 (d, $J=6.4$ Hz, 11'-H₃) and 0.98 (d, $J=6.4$ Hz, 11-H₃)], two furans [δ 6.40 (dd, $J=0.6, 1.5$ Hz, 13, 13'-H), 7.37 (dd, $J=0.6, 1.5$ Hz, 16-H), 7.40 (dd, $J=0.6, 1.5$ Hz, 16'-H), and 7.43 (dd, $J=1.5, 1.5$ Hz, 14, 14'-H)], a thiolane sulfoxide [δ 2.02 (d, $J=15.0$ Hz, 17 β -H), 2.28 (d, $J=15.0$ Hz, 17 α -H), 2.32 (d, $J=14.3$ Hz, 17' α -H), and 2.92 (d, $J=14.3$ Hz, 17' β -H)], a hemiaminal-bearing quinolizidine ring [δ 2.35 (m, 10'-H), 3.68 (dd, $J=3.4, 10.7$ Hz, 4'-H), and 4.14 (s, 6'-H)], and a quinolizidine ring [δ 1.53 (m, 8 β -H), 1.64 (m, 10-H), 1.66 (m, 8 α -H), 1.89 (dd, $J=1.2, 11.3$ Hz, 6 β -H), 2.54 (dd, $J=2.7, 11.3$ Hz, 6 α -H), and 3.04 (dd, $J=3.1, 11.6$ Hz, 4-H)]. Comparison of the spectral data for **1** with those for 6-hydroxythiobinupharidine (**5**) and 6,6'-dihydroxythiobinupharidine (**6**)^{1,6} led us to presume the thiohemiaminal type dimeric sesquiterpene structure of **1**. Treatment of **1** with sodium borohydride (NaBH₄) in methanol furnished *anti*-thiobinupharidine sulfoxide (**9**), whose absolute stereostructure was previously reported.⁸


The position of the hydroxyl group in **1** was clarified by a heteronuclear multiple bond correlation (HMBC) experiment. Thus, long-range correlations were observed between the 6'-proton and the 4', 7', 8', 10'-carbons, between the 4'-proton and the 2', 3', 6', 12'-carbons, and between the 10'-proton and the 1', 4', 6'-carbons. The stereostructure of **1** including the configuration of the 6'-hydroxyl group was confirmed by a ¹H-NMR nuclear Overhauser and exchange spectroscopy (NOESY) experiment, which showed NOE correlations between the following protons; 6'-H and 17 β , 17' β , 8 α -H shown in Figure 3. On the basis of above-mentioned evidence, the structure of nupharpumilamine A (**1**) was determined as shown.

Nupharpumilamine B (2)


Nupharpumilamine B (2) was also isolated as colorless oil with negative optical rotation ($[\alpha]_D^{25} -60.1^\circ$). The IR spectrum of 2 showed absorption bands due to hydroxyl (3380 cm^{-1}), *trans*-fused quinolizidine ring (Bohlmann absorption, $2950\text{--}2750\text{ cm}^{-1}$), sulfinyl (1021 cm^{-1}), and furan (874 cm^{-1}) functions. The EI-MS and positive-ion FAB-MS of 2 showed dehydration peaks [m/z 508 ($M^+ - H_2O$), 509 ($M + H - H_2O$)] in addition of fragment ion peaks at m/z 491, 230, and 178. In the negative-ion FAB-MS of 2, a quasimolecular ion peak was observed at m/z 525 ($M - H$)⁻ and high-resolution MS measurement revealed the molecular formula of 2 to be $C_{30}H_{42}N_2O_4S$, which was identical with that of 1. The 1H -NMR (CD_3OD) and ^{13}C -NMR (Table 1) spectra⁷ of 2 indicated the presence of two tertiary methyls [δ 0.78 (d, $J=6.4$ Hz, $11'$ -H₃) and 0.81 (d, $J=6.4$ Hz, 11 -H₃)], two furans [δ 6.01 (dd, $J=0.6, 1.5$ Hz, $13'$ -H), 6.30 (dd, $J=0.6, 1.8$ Hz, 13 -H), 6.89 (dd, $J=1.5, 1.5$ Hz, $14'$ -H), 7.17 (dd, $J=0.6, 1.5$ Hz, $16'$ -H), 7.31 (dd, $J=1.5, 1.8$ Hz, 14 -H), and 7.35 (dd, $J=0.6, 1.5$ Hz, 16 -H)], a thiolane sulfoxide [δ




6-hydroxythiobinupharidine (5)



6,6'-dihydroxythiobinupharidine (6)



6-hydroxythionuphlutine B (7)



6'-hydroxythionuphlutine B (8)





*anti*-thiobinupharidine sulfoxide (9)*syn*-thiobinupharidine sulfoxide (10)thionuphlutine B β -sulfoxide (11)neothiobinupharidine β -sulfoxide (12)

Figure 2




Figure 3. NOE Correlations in the NOESY Spectra of **1**, **2**, **3**, and **4**

1.56 (d, $J=14.3$ Hz, $17\alpha\text{-H}$), 1.89 (d, $J=14.3$ Hz, $17\beta\text{-H}$) and 2.35 (d, $J=15.6$ Hz, $17'\alpha\text{-H}$), 2.97 (d, $J=15.6$ Hz, $12'\beta\text{-H}$), a hemiaminal-bearing quinolizidine ring [δ 3.53 (dd, $J=3.1$, 11.6 Hz, $4'\text{-H}$) and 4.08 (s, $6'\text{-H}$)], and a quinolizidine ring [δ 1.67 (dd, $J=0.9$, 12.8 Hz, $6\alpha\text{-H}$), 3.20 (dd, $J=2.4$, 12.8 Hz, $6\beta\text{-H}$), and 1.56 (m, 10-H)] From the comparison of the spectral data for **2** with those for **1** and 6'-hydroxythionuphlutine B (**8**), **2** was presumed to be a sulfoxide of **8**. In the HMBC experiment of **2**, long-range correlations were observed between the 6'-proton and the 4', 7', 8', 10'-carbons and between the 4'-proton and 6', 12'-carbons. The NOESY experiment of **2** showed NOE correlations as shown in Figure 3. The configuration of the sulfinyl group was clarified by comparison of the ^1H -NMR data for **2** with those for 6'-hydroxythionuphlutine B (**8**),^{1,6a} which showed the characteristic anisotropy and proximity effects of the *anti*-sulfinyl group.⁹ Finally, the NaBH_4 treatment of **2** yielded thionuphlutine B β -sulfoxide (**11**) Consequently, the structure of nupharpumilamine B (**2**) was characterized as shown.

Nupharpumilamine C (**3**)

Nupharpumilamine C (**3**), obtained as colorless oil with negative optical rotation ($[\alpha]_D^{25} -75.1^\circ$), showed Bohlmann absorption bands and absorption bands ascribable to sulfinyl and furan functions in the IR spectrum. The molecular formula $\text{C}_{30}\text{H}_{42}\text{N}_2\text{O}_3\text{S}$ of **3** was confirmed from the quasimolecular ion peak at m/z 511 ($\text{M}+\text{H}$)⁺ observed in the positive-ion FAB-MS and by high-resolution MS measurement. The ^1H -NMR (CD_3OD) and ^{13}C -NMR (Table 1) spectra⁷ of **3** showed signals

Table 1. ^{13}C -NMR Data for 1, 2, 3, 4, 9, 10, 11, and 12

	1 ¹⁾	2 ¹⁾	3 ²⁾	4 ²⁾	9 ²⁾	10 ²⁾	11 ²⁾	12 ²⁾
C-1	37.3	37.4	32.3	32.4	36.0	36.0	36.0	36.3
C-2	34.5	34.9	31.9 ^{a)}	32.0 ^{a)}	33.4	33.4	33.6	33.8
C-3	35.7	36.9	29.7	30.2	34.7	35.1 ^{a)}	35.8	35.0 ^{a)}
C-4	60.9	62.6	60.4	60.7	59.7	59.7 ^{b)}	61.1	61.1
C-6	59.7	56.6	59.8	57.2	59.2	56.3	55.9	55.4
C-7	68.2	71.6	66.0	66.3	66.2	66.0	70.2	68.4
C-8	30.8	34.8	28.2	31.5	28.3	31.7	34.3	32.5
C-9	28.5 ^{a)}	29.1	27.3	27.7	27.0	27.4	28.0	28.1
C-10	70.0	69.9	64.6	64.8	68.6	68.8	68.4	68.6
C-11	19.6	19.5	13.3	13.4	18.9	19.1	19.1	19.3
C-12	129.8	129.8	128.9	129.1	128.7	129.0 ^{c)}	128.5	128.8
C-13	110.3	111.2	109.1	109.1	109.2	109.4 ^{d)}	110.2	110.1
C-14	144.6	143.7	142.9	142.9	142.9	142.9	142.3	142.9
C-16	141.5	140.7	139.4	139.4	139.5	139.6 ^{e)}	139.1	139.3
C-17	42.8	47.1	40.9	40.2	41.8	41.1	47.0	49.3
C-1'	38.5	38.6	32.3	32.4	36.1	36.0	36.3	36.6
C-2'	34.3	34.2	31.8 ^{a)}	31.9 ^{a)}	33.4	33.5	33.4	33.6
C-3'	37.2	37.2	29.8	29.8	34.9	34.9 ^{a)}	35.0	36.0 ^{a)}
C-4'	56.0	56.0	60.3	60.4	59.5	60.0 ^{b)}	59.2	59.6
C-6'	96.4	96.6	64.7	64.7	64.2	64.2	65.9	65.4
C-7'	51.7	52.1	43.4	44.4	43.6	44.1	45.9	45.1
C-8'	32.9	33.4	38.2	40.3	38.4	40.2	38.8	39.8
C-9'	29.5 ^{a)}	30.3	28.0	27.8	27.8	27.5	29.0	28.5
C-10'	59.5	58.8	64.6	64.7	68.6	68.8	68.1	68.5
C-11'	19.4	19.5	13.3	13.3	19.0	19.1	19.0	19.2
C-12'	130.1	129.9	129.1	129.3	128.9	128.8 ^{c)}	128.7	129.6
C-13'	110.4	109.7	109.0	109.2	109.1	109.2 ^{d)}	108.4	109.2
C-14'	144.5	144.4	142.8	142.9	142.8	142.9	143.1	143.5
C-16'	141.4	141.0	139.3	139.4	139.4	139.5 ^{e)}	139.2	139.8
C-17'	58.2	56.9	60.3	59.7	60.0	59.4	59.6	57.5

The spectra were taken with CD_3OD ¹⁾ or CDCl_3 ²⁾.

a), b), c), d), e) Assignments may be interchangeable within the same column.

assignable to two tertiary methyls [δ 1.11 (d, $J=7.0$ Hz, 11'-H₃) and 1.13 (d, $J=7.0$ Hz, 11-H₃)], two furans [δ 6.34 (d, $J=1.5$ Hz, 13-H), 6.35 (d, $J=1.5$ Hz, 13'-H), 7.24 (br s, 16, 16'-H), 7.32 (dd, $J=1.5, 1.5$ Hz, 14-H), and 7.33 (dd, $J=1.5, 1.5$ Hz, 14'-H)], a thiolane sulfoxide [δ 1.92 (d, $J=14.3$ Hz, 17 α -H), 2.15 (d, $J=14.3$ Hz, 17 β -H), 2.43 (d, $J=13.4$ Hz, 17' β -H), and 2.56 (d, $J=13.4$ Hz, 17' α -H)], and two quinolizidine rings [δ 1.48 (d, $J=11.6$ Hz, 6 β -H), 1.69 (m, 1'-H), 1.75 (m, 1-H), 1.85 (d, $J=11.3$ Hz, 6 β -H), 1.90 (m, 10'-H), 2.01 (m, 10-H), 2.53 (dd, $J=2.4, 11.3$ Hz, 6 α -H), 2.80 (dd, $J=3.4, 11.6$ Hz, 4'-H), 2.87 (dd, $J=3.4, 11.6$ Hz, 4-H), 2.90 (dd, $J=2.4, 11.6$ Hz, 6' α -H)]. The proton and carbon signals in the ^1H -NMR and ^{13}C -NMR spectra of **3** were superimposable on those of *anti*-thiobinupharidine sulfoxide (**9**),⁸ except for the C-1—3 and C-1'—3' signals, which showed high-field shift by the axial 1 and 1'-methyl groups. The NOESY experiment of **3** showed NOE

correlations between the following protons : 4-H and 6 β , 10-H; 11-H₃ and 3 α , 9 α -H; 17 β -H and 9 α , 6' α -H; 17 α -H and 6 α , 9' α -H; 11'-H₃ and 3' α , 9' α -H; 4' β -H and 6' β , 10'-H, as shown in Figure 3, so that the stereostructure including the C-1 and C-1' configurations of **3** was characterized. Comparison of the ¹H-NMR and ¹³C-NMR data for **3** with those (Table 1) for *anti*- and *syn*-thiobinupharidine sulfoxides (**9**, **10**)¹⁰ and neothiobinupharidine β -sulfoxide (**12**)¹¹ led us to confirm the structure of nupharpumilamine C (**3**) as shown.

Nupharpumilamine D (**4**)

Nupharpumilamine D (**4**) was also obtained as colorless oil with positive optical rotation ($[\alpha]_D^{25} +20.8^\circ$) and its IR spectrum showed Bohlmann absorption and absorption bands due to sulfinyl and furan functions. Here again, the molecular formula C₃₀H₄₂N₂O₃S of **4**, which was identical with that of nupharpumilamine C (**3**), was determined from the quasimolecular ion peak at m/z 511 (M+H)⁺ in the positive-ion FAB-MS and by high-resolution MS measurement. The ¹H-NMR and ¹³C-NMR (Table 1) spectra⁷ of **4** indicated the presence of two methyls, two furans, a thiolane sulfoxide, and two quinolizidine rings. The proton and carbon signals in the ¹H-NMR and ¹³C-NMR (Table 1) spectra of **4** were very similar to those of *syn*-thiobinupharidine sulfoxide (**10**), except for the C-1—3 and C-1'—3' signals, which were significantly resembled with those of **3**. Furthermore, as is apparent from Figure 3, the NOE correlations in the NOESY experiment of **4** were almost the same as those of **3**. On the basis of this evidence, the structure of nupharpumilamine D (**4**) was characterized as shown.

EXPERIMENTAL

The instruments used to obtain physical data and the experimental conditions for chromatography were the same as described in our previous paper.²

Isolation of Nupharpumilamine A (**1**), B (**2**), C (**3**), and D (**4**) from the Alkaloid Fraction of Chinese Nupharis Rhizoma

The AcOEt soluble fraction (16.3g) from Chinese Nupharis Rhizoma (Osaka, 7.9 kg) was subjected to ordinary silica gel column chromatography [CHCl₃-AcOEt-Et₂NH (20 : 1 : 1)→MeOH-Et₂NH (10 : 1)] to give three fractions [Fr. 1 (9.5 g), Fr. 2 (3.0 g), and Fr. 3 (3.8 g)] as described previously.¹ Fraction 1 (8.7 g) was separated by Chromatorex NH DM1020 (Fuji Silysia Chemical Ltd.) column chromatography [260 g; *n*-hexane-CH₂Cl₂-AcOEt (100 : 100 : 1→20 : 20 : 1)→CHCl₃-MeOH (10 : 1)] followed by repeated hplc [Develosil ODS-HG-5, 250x20 mm i.d., i) MeOH-H₂O-Et₂NH (890 : 10 : 6), ii) MeOH-H₂O-Et₂NH (750 : 150 : 6), iii) MeOH-H₂O-Et₂NH (800 : 100 : 6), iv) MeCN-H₂O-Et₂NH (89 : 1 : 3), flow rate : 8.0 ml/min] to give **3** (10.4 mg) and **4** (17.8 mg). Chromatorex NH DM1020 column chromatography [90 g, *n*-hexane-AcOEt (3 : 1→1 : 5)→AcOEt-acetone (5 : 1→1 : 5)→MeOH-Et₂NH (9 : 1)] of fraction 2 furnished six fractions. Fraction 2-3 (234 mg) was subjected to HPLC (Develosil ODS-HG-5, 250x20 mm i.d., MeOH-H₂O-Et₂NH (850 : 50 : 6), flow rate : 8.0 ml/min) to give **1** (17.4 mg). The HPLC (same conditions as described above) of fraction 2-4 (235 mg) yielded **2** (40.6 mg).

Nupharpumilamine A (**1**) : colorless oil, positive for Dragendorff reagent, $[\alpha]_D^{25} - 33.4^\circ$ ($c=0.6$, CHCl₃). High-resolution EI-MS (m/z) : Calcd for C₃₀H₄₀N₂O₃S (M⁺-H₂O) : 508.2759; Found : 508.2752. High-resolution negative-ion FAB-MS (m/z) : Calcd for C₃₀H₄₁N₂O₄S (M-H)⁻ : 525.2787; Found : 525.2769. IR (film, cm⁻¹) : 3360, 2950—2750, 1501, 1456, 1439, 1159, 1024, 874. ¹H-NMR (CD₃OD, 500 MHz, δ) : 0.92 (3H, d, $J=6.4$ Hz, 11'-H₃), 0.98 (3H, d, $J=6.4$ Hz, 11-H₃), 1.33 (1H, m, 9 α -H), 1.53 (1H, m, 8 β -H), 1.64 (1H, m, 10-H), 1.66 (1H, m, 8 α -H), 1.89 (1H, dd, $J=1.2, 11.3$ Hz, 6 β -H), 2.02 (1H, d, $J=15.0$ Hz, 17 β -H), 2.28 (1H, d, $J=15.0$ Hz, 17 α -H), 2.32 (1H, dd, $J=14.3$ Hz, 17' α -H), 2.35 (1H, m, 10'-H),

2.54 (1H, dd, $J=2.7$, 11.3 Hz, 6 α -H), 2.92 (1H, d, $J=14.3$ Hz, 17 β -H), 3.04 (1H, dd, $J=3.1$, 11.6 Hz, 4-H), 3.68 (1H, dd, $J=3.4$, 10.7 Hz, 4'-H), 4.14 (1H, s, 6'-H), 6.40 (2H, dd, $J=0.6$, 1.5 Hz, 13, 13'-H), 7.37 (1H, dd, $J=0.6$, 1.5 Hz, 16-H), 7.40 (1H, dd, $J=0.6$, 1.5 Hz, 16'-H), 7.43 (2H, dd, $J=1.5$, 1.5 Hz, 14, 14'-H). $^{13}\text{C-NMR}$ (CD_3OD , 125 MHz, δ_{C}): see Table 1. EI-MS (m/z): 508 ($\text{M}^+-\text{H}_2\text{O}$), 491, 447, 230, 178, 94. Negative-ion FAB-MS (m/z): 525 (M-H) $^-$, 459 ($\text{M-C}_4\text{H}_3\text{O}$) $^-$, 305 ($\text{M-C}_{14}\text{H}_{22}\text{NO}$) $^-$. Positive-ion FAB-MS (m/z): 509 ($\text{M+H-H}_2\text{O}$) $^+$.

Nupharpumilamine B (2): colorless oil, positive for Dragendorff reagent, $[\alpha]_{\text{D}}^{25}$ -60.1 $^\circ$ ($c=1.6$, CHCl_3). High-resolution EI-MS (m/z): Calcd for $\text{C}_{30}\text{H}_{40}\text{N}_2\text{O}_3\text{S}$ ($\text{M}^+-\text{H}_2\text{O}$): 508.2760; Found: 508.2755. High-resolution negative-ion FAB-MS (m/z): Calcd for $\text{C}_{30}\text{H}_{41}\text{N}_2\text{O}_4\text{S}$ (M-H) $^-$: 525.2787; Found: 525.2806. IR (film, cm^{-1}): 3380, 2950—2750, 1503, 1439, 1157, 1021, 874. $^1\text{H-NMR}$ (CD_3OD , 500 MHz, δ): 0.78 (3H, d, $J=6.4$ Hz, 11'-H₃), 0.81 (3H, d, $J=6.4$ Hz, 11-H₃), 1.56 (1H, m, 10-H), 1.56 (1H, d, $J=14.3$ Hz, 17 α -H), 1.67 (1H, dd, $J=0.9$, 12.8 Hz, 6 α -H), 1.89 (1H, d, $J=14.3$ Hz, 17 β -H), 2.24 (1H, ddd, $J=2.4$, 8.5, 11.0 Hz, 10'-H), 2.35 (1H, d, $J=15.6$ Hz, 17 α -H), 2.91 (1H, dd, $J=6.4$, 7.3 Hz, 4-H), 2.97 (1H, d, $J=15.6$ Hz, 17 β -H), 3.20 (1H, dd, $J=2.4$, 12.8 Hz, 6 β -H), 3.53 (1H, dd, $J=3.1$, 11.6 Hz, 4'-H), 4.08 (1H, s, 6'-H), 6.01 (1H, dd, $J=0.6$, 1.5 Hz, 13'-H), 6.30 (1H, dd, $J=0.6$, 1.8 Hz, 13-H), 6.89 (1H, t, $J=1.5$ Hz, 14'-H), 7.17 (1H, dd, $J=0.6$, 1.5 Hz, 16'-H), 7.31 (1H, dd, $J=1.5$, 1.8 Hz, 14-H), 7.35 (1H, dd, $J=0.6$, 1.5 Hz, 16-H). $^{13}\text{C-NMR}$ (CD_3OD , 125 MHz, δ_{C}): see Table 1. EI-MS (m/z): 508 ($\text{M}^+-\text{H}_2\text{O}$), 491, 230, 178. Negative-ion FAB-MS (m/z): 525 (M-H) $^-$, 459 ($\text{M-C}_4\text{H}_3\text{O}$) $^-$. Positive-ion FAB-MS (m/z): 509 ($\text{M+H-H}_2\text{O}$) $^+$.

Nupharpumilamine C (3): colorless oil, positive for Dragendorff reagent, $[\alpha]_{\text{D}}^{25}$ -75.1 $^\circ$ ($c=0.5$, CHCl_3). High-resolution FAB-MS (m/z): Calcd for $\text{C}_{30}\text{H}_{43}\text{N}_2\text{O}_3\text{S}$ (M+H) $^+$: 511.2994; Found: 511.2976. IR (film, cm^{-1}): 2950—2750, 1501, 1449, 1381, 1159, 1026, 874. $^1\text{H-NMR}$ (CDCl_3 , 500 MHz, δ): 1.11 (3H, d, $J=7.0$ Hz, 11'-H₃), 1.13 (3H, d, $J=7.0$ Hz, 11-H₃), 1.48 (1H, d, $J=11.6$ Hz, 6 β -H), 1.69 (1H, m, 1'-H), 1.75 (1H, m, 1-H), 1.85 (1H, d, $J=11.3$ Hz, 6 β -H), 1.90 (1H, m, 10'-H), 1.92 (1H, d, $J=14.3$ Hz, 17 α -H), 2.01 (1H, m, 10-H), 2.15 (1H, d, $J=14.3$ Hz, 17 β -H), 2.43 (1H, d, $J=13.4$ Hz, 17 β -H), 2.53 (1H, dd, $J=2.4$, 11.3 Hz, 6 α -H), 2.56 (1H, d, $J=13.4$ Hz, 17 α -H), 2.80 (1H, dd, $J=3.4$, 11.6 Hz, 4'-H), 2.87 (1H, dd, $J=3.4$, 11.6 Hz, 4-H), 2.90 (1H, dd, $J=2.4$, 11.6 Hz, 6 α -H), 6.34 (1H, d, $J=1.5$ Hz, 13-H), 6.35 (1H, d, $J=1.5$ Hz, 13'-H), 7.24 (2H, br s, 16, 16'-H), 7.32 (1H, t, $J=1.5$ Hz, 14-H), 7.33 (1H, t, $J=1.5$ Hz, 14'-H). $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz, δ_{C}): see Table 1. FAB-MS (m/z): 511 (M+H) $^+$.

Nupharpumilamine D (4): colorless oil, positive for Dragendorff reagent, $[\alpha]_{\text{D}}^{25}$ +20.8 $^\circ$ ($c=0.9$, CHCl_3). High-resolution FAB-MS (m/z): Calcd for $\text{C}_{30}\text{H}_{43}\text{N}_2\text{O}_3\text{S}$ (M+H) $^+$: 511.2994; Found: 511.2980. IR (film, cm^{-1}): 2950—2750, 1501, 1451, 1381, 1159, 1021, 874. $^1\text{H-NMR}$ (CDCl_3 , 500 MHz, δ): 1.14 (3H, d, $J=7.0$ Hz, 11'-H₃), 1.16 (3H, d, $J=6.7$ Hz, 11-H₃), 1.47 (1H, d, $J=11.3$ Hz, 6 β -H), 1.94 (1H, m, 10'-H), 1.95 (1H, d, $J=11.3$ Hz, 6 α -H), 2.03 (1H, m, 10-H), 2.03 (1H, d, $J=14.3$ Hz, 17 α -H), 2.33 (1H, d, $J=14.3$ Hz, 17 β -H), 2.47 (1H, dd, $J=2.4$, 11.3 Hz, 6 α -H), 2.51 (2H, s, 17'-H₂), 2.83 (1H, dd, $J=3.4$, 11.9 Hz, 4'-H), 2.94 (1H, dd, $J=3.4$, 11.0 Hz, 4-H), 2.97 (1H, dd, $J=2.4$, 11.3 Hz, 6 β -H), 6.38 (1H, d-like, 13-H), 6.38 (1H, d-like, 13'-H), 7.25 (1H, br s, 16'-H), 7.28 (1H, br s, 16-H), 7.34 (1H, t-like, 14-H), 7.34 (1H, t-like, 14'-H). $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz, δ_{C}): Table 1. FAB-MS (m/z): 511 (M+H) $^+$.

Treatment of Nupharpumilamine A (1) with NaBH_4 Giving *anti*-Thiobinupharidine Sulfoxide (9)

A solution of nupharpumilamine A (1, 2.3 mg) in MeOH (0.5 mL) was treated with NaBH_4 (4.6 mg) and the whole mixture was stirred at rt for 30 min. After treatment of the reaction mixture with acetone (0.5 mL), the whole was poured into brine and then it was extracted with AcOEt. The AcOEt extract was washed with brine and dried over MgSO_4 . Removal of the solvent under reduced pressure gave *anti*-thiobinupharidine sulfoxide (9, 2.2 mg), which was identified by HPTLC, HPLC, IR, and $^1\text{H-NMR}$ (CD_3OD) spectra comparisons with an authentic sample¹

Treatment of Nupharpumilamine B (2) with NaBH₄ Giving Thionuphlutine B β -Sulfoxide (11)

A solution of nupharpumilamine B (2, 3.9 mg) in MeOH (0.5 mL) was treated with NaBH₄ (10.0 mg) and the whole mixture was stirred at rt for 30 min. The reaction mixture was worked up as described above to yield thionuphlutine B β -sulfoxide (11, 3.8 mg), which was identified by HPTLC, HPLC, IR, and ¹H-NMR (CD₃OD) spectra comparisons with an authentic sample.¹

ACKNOWLEDGMENT

The authors are grateful to the Ministry of Education, Science, Sports and Culture of Japan for Grant-in-Aid for Scientific Research (C) (No. 08672461).

REFERENCES AND NOTES

1. Part VI : M. Yoshikawa, T. Murakami, S. Wakao, A. Ishikado, H. Shimoda, N. Murakami, J. Yamahara, and H. Matsuda, *Chem. Pharm. Bull.* to be published.
2. a) M. Yoshikawa, S. Hatakeyama, N. Tanaka, Y. Fukuda, J. Yamahara, and N. Murakami, *Chem. Pharm. Bull.*, 1993, **41**, 1948; b) M. Yoshikawa, S. Hatakeyama, N. Tanaka, T. Matsuoka, J. Yamahara, and N. Murakami, *ibid.*, 1993, **41**, 2109; c) M. Yoshikawa, S. Yamaguchi, N. Chatani, Y. Nishino, T. Matsuoka, J. Yamahara, N. Murakami, H. Matsuda, and M. Kubo, *Yakugaku Zasshi*, 1994, **114**, 241; d) M. Yoshikawa, S. Yamaguchi, H. Matsuda, Y. Kohda, H. Ishiyama, N. Tanaka, J. Yamahara, and N. Murakami, *Chem. Pharm. Bull.*, 1994, **42**, 1813; e) M. Yoshikawa, S. Yamaguchi, H. Matsuda, N. Tanaka, J. Yamahara, and N. Murakami, *ibid.*, 1994, **42**, 2430.
3. J. Yamahara, H. Shimoda, H. Matsuda, and M. Yoshikawa, *Biol. Pharm. Bull.*, 1996, **19**, 1241.
4. M. Yoshikawa, T. Murakami, A. Ishikado, H. Shimoda, N. Murakami, H. Matsuda, and J. Yamahara, presented at the 43rd Annual Meeting of the Japanese Society of Pharmacognosy, Tokyo, September 1996, Abstract of Papers; p 206.
5. F. Bohlmann, *Angew. Chem.*, 1957, **69**, 641.
6. a) R. T. LaLonde, C. F. Wong, and W. P. Cullen, *Tetrahedron Lett.*, 1970, **51**, 4477; b) R. T. LaLonde, C. F. Wong, and K. C. Das, *J. Org. Chem.*, 1974, **39**, 2892; c) T. I. Martin, D. B. MacLean, J. T. Wrobel, A. Iwanow, and W. Starzec, *Can. J. Chem.*, 1974, **52**, 2705; d) R. T. LaLonde, C. F. Wong, and K. C. Das, *ibid.*, 1974, **52**, 2714; d) R. T. LaLonde, C. F. Wong, and A. I. M. Tsai, *Org. Mass. Spectr.*, 1976, **11**, 183.
7. The ¹H-NMR and ¹³C-NMR spectra of **1**, **2**, **3**, and **4** were assigned with the aid of homo and hetero correlation spectroscopy (¹H-¹H, ¹H-¹³C COSY), distortionless enhancement by polarization transfer (DEPT), HMBC, and NOESY experiment.
8. A. Iwanow, K. Wojtasiewicz, and J. T. Wrobel, *Phytochemistry*, 1986, **25**, 2227.
9. a) J. T. Wrobel, J. Ruskowska, and K. Kabzinska, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1977, **XXV**, 927; b) J. T. Wrobel, J. Ruskowska, and K. Wojtasiewicz, *J. Mol. Str.*, 1977, **41**, 287; c) J. T. Wrobel, J. Ruskowska, and H. Bielawska, *Polish J. Chem.*, 1979, **53**, 39.
10. a) J. T. Wrobel, A. Iwanow, and K. Wojtasiewicz, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.*, 1976, **XXIV**, 99; b) J. T. Wrobel, J. Ruskowska, and K. Wojtasiewicz, *J. Mol. Str.*, 1978, **50**, 299.
11. R. T. LaLonde, C. F. Wong, A. I. M. Tsai, J. T. Wrobel, J. Ruskowska, K. Kabzinska, T. I. Martin, and D. B. MacLean, *Can. J. Chem.*, 1976, **54**, 3861.