


SYNTHESIS OF HETEROARENECARBONITRILES BY ELECTROPHILIC CYANATION; REACTION OF METALATED HETEROARENES WITH *p*-TOLUENESULFONYL CYANIDE ¹

Izuru Nagasaki, Yumiko Suzuki, Ken-ichi Iwamoto, Takeo Higashino,
and Akira Miyashita*

*School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Shizuoka
422, Japan*

Abstract—Several heteroarenecarbonitriles (**5**) were synthesized in moderate yields from heteroarenes (**3**) through metalation, followed by electrophilic cyanation using *p*-toluenesulfonyl cyanide. Similarly, trimethylsilylheteroarenes (**8**) were converted to heteroarenecarbonitriles (**5**) in good yields by treatment with *p*-toluenesulfonyl cyanide.

In the previous paper,¹ we showed that *p*-toluenesulfonyl cyanide (TsCN) is a useful reagent for electrophilic cyanation. Namely, in the presence of base, mercaptoheteroarenes (**1**) reacted with TsCN to give the corresponding thiocyanatoheteroarenes (**2**) in good yields. In this reaction, TsCN provides the cyanogen cation (CN⁺). To extend the scope of this electrophilic cyanation using TsCN, we applied this method to the



Scheme 1

preparation of heteroarenecarbonitriles (**5**). We anticipated that heteroarenyl carbanions (Het[−]) would react with TsCN to give the corresponding heteroarenecarbonitriles (**5**). This is a carbon–carbon bond-forming reaction and represents a method for electrophilic introduction of a cyano group into heteroarenes. Many methods for preparing heteroarenecarbonitriles (**5**) have been reported,² but there are only a few papers describing electrophilic cyanation.³ Here we wish to report the results of the reaction of metalated heteroarenes (**4**) with TsCN.


Metalation is a useful method to produce carbanions from organic compounds. Thus, metalloheteroarenes (**4**) should react with TsCN to give the heteroarene carbonitriles (**5**) by electrophilic cyanation. We examined several kinds of metalloheteroarenes.

It is known that imidazo[1,5-*a*]pyridine (**3a**) is lithiated at the 3-position by treatment with *n*-BuLi.⁴ Treatment of imidazo[1,5-*a*]pyridine (**3a**) with *n*-BuLi, followed by dropwise addition of TsCN (Method A) gave imidazo[1,5-*a*]pyridine-3-carbonitrile (**5a**) in 59% yield. However, when the solution of lithiated imidazo[1,5-*a*]pyridine was added dropwise to TsCN (Method B), **5a** was formed in 65% yield. In the preparation of 2-benzofurancarbonitrile (**5f**), only 26% yield was obtained by Method A, and 71% yield by Method B (Scheme 2). The difference may arise because the produced heteroarene carbonitriles (**5**) can further react with lithioheteroarenes. In the cyanation of benzothiazole (**3e**) by method A, di-2-benzothiazolyl ketone (**7**) was the main product in 61% yield, together with a trace of 2-benzothiazole carbonitrile (**5e**), as shown in Scheme 3.

$\text{Het-H} \xrightarrow{n\text{-BuLi}} \left[\text{Het-Li} \right] \xrightarrow{\text{TsCN}} \text{Het-CN}$ (or Het-X)					
Het—		5, Yield (%)		Het— or Ar—	5 or 6, Yield (%)
		Method A	Method B		Method A Method B
	5a	59	65		5f 26 71
	5b	25	50		5g 52
	5c	40	77		5h 66 73
	5d	39	57		6j 57
	5e	trace	33		6k 24

Scheme 2


Cyanogen bromide (BrCN) is also used in electrophilic cyanation. In the synthesis of thiocyanatoheteroarenes (**2**), BrCN and TsCN, both gave thiocyanatoheteroarenes in good yields.¹ However, Bolze and Dell reported



Scheme 3


that cyanation by treatment of lithioarenes with BrCN does not proceed, namely, treatment of 2-lithio-3-methoxyanisole with BrCN gave 2-bromo-3-methoxyanisole.⁵ The expected 3,6-dimethoxybenzonitrile (**6j**) could not be obtained. This result indicates that BrCN affords the bromo cation (Br^+) under these conditions. For direct introduction of a cyano group into heteroarenes, we found that TsCN is an effective reagent, presumably because the cyano cation (CN^+) is readily formed. TsCN provides a so-called soft cyano cation in comparison with BrCN because of the nature of the binding groups, bromine and sulfone.

To find a more suitable method for this electrophilic cyanation, we tried other methods to generate the heteroarenyl carbanion. It is known that the carbon-silicon bond of trimethylsilylheteroarenes cleaves easily, resulting in the formation of the heteroarenyl carbanion.⁶ This led us to examine the use of trimethylsilylheteroarenes (**8**) as the heteroarenyl carbanion donors. As shown in Scheme 4, we obtained good results in electrophilic cyanation using trimethylsilylheteroarenes compared with that using lithioheteroarenes. However, it seems that the donor effect of the arenyl carbanion of trimethylsilylarenes (**8**) is less than that of lithioarenes (**4**).




Scheme 4

Grignard reagents also provide carbanions. Treatment of the pyrrole (**9**) with methylmagnesium iodide followed by reaction with TsCN furnished 2-pyrrolecarbonitrile (**10**) in 53% yield. We have applied this



Scheme 5

electrophilic cyanation to regioselective synthesis of indolecarbonitriles,⁷ and have already shown that 1-benzenesulfonylindole (**3h**) affords 1-benzenesulfonyl-2-indolecarbonitrile (**5h**) *via* lithiation. Treatment of indole with sodium hydride followed by cyanation with TsCN gave 1-indolecarbonitrile (**12**). 3-Indolecarbonitrile (**13**) was obtained by forming indolylmagnesium iodide, followed by cyanation with TsCN. Appropriate metalations can thus be used to obtain regioselective cyanation of indoles (**11**).



Scheme 6

In conclusion, electrophilic cyanation of heteroarenes (**3**) using TsCN can be achieved through metalation. By this procedure, several heteroarenecarbonitriles (**5**) were synthesized from metalated heteroarenes, such as lithioheteroarenes (**4**), heteroarenylmagnesium halide, and trimethylsilylheteroarenes (**8**).

EXPERIMENTAL

All melting points are uncorrected. IR spectra were recorded on a JASCO A-102 diffraction grating IR spectrophotometer. ¹H-NMR spectra were measured at 60 MHz on a JEOL PMX60SI NMR spectrometer, and at 270 MHz on a JEOL JNM-GSX270 FT-NMR spectrometer. Chemical shifts are quoted in parts per million (ppm) with tetramethylsilane as an internal standard, and coupling constants (*J*) are given in Hz.

Synthesis of Imidazo[1,5-*a*]pyridine-2-carbonitriles; A Typical Procedure.

Method A. A solution of *n*-BuLi in *n*-hexane (1.63 mol/L, 1.8 mL, 3.0 mmol) was slowly added to a solution of imidazo[1,5-*a*]pyridine (3.0 mmol) in 6 mL of THF with stirring at -78°C under an argon atmosphere and the solution was further stirred for 30 min at -78°C . A solution of TsCN (815 mg, 4.5 mmol) in 4 mL of THF was added dropwise to the resulting solution at -78°C and the solution was further stirred for 30 min. The cooling bath was removed and the reaction mixture was further stirred for 2 h, then poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on SiO_2 with *n*-hexane, benzene, and CH_2Cl_2 .

Method B. A solution of *n*-BuLi in *n*-hexane (1.63 mol/L, 1.8 mL, 3.0 mmol) was slowly added to a solution of imidazo[1,5-*a*]pyridine (3.0 mmol) in 6 mL of THF with stirring at -78°C under an argon atmosphere and the solution was further stirred for 30 min at -78°C . The resulting solution was added dropwise to a solution of TsCN (815 mg, 4.5 mmol) in 4 mL of THF at -78°C , and the whole was further stirred for 30 min. The cooling bath was removed and the reaction mixture was further stirred for 2 h. The reaction mixture was poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography on SiO_2 with *n*-hexane, benzene, and/or CH_2Cl_2 .

2-Lithio-1-methylimidazole (**4c**), 2-lithio-1-methylbenzimidazole (**4d**), 2-lithio-1-benzothiazole (**4e**), and 2-lithio-1-benzenesulfonylindole (**4h**) were prepared similarly.

The lithiation of other heteroarenes was carried out according to the reported procedures; 3-lithiopyridine (**4b**),⁸ 2-lithiobenzofuran (**4f**),⁹ 2-lithio-1-benzothiophene (**4g**),¹⁰ 2-lithio-3-methoxyanisole,⁸ and 2-lithio-*N,N*-dimethylaniline.¹¹

Imidazo[1,5-*a*]pyridine-3-carbonitrile (**5a**): The fraction eluted with *n*-hexane–benzene (1:1) gave **5a**. Colorless needles (*n*-hexane–benzene), mp $123\text{--}124^{\circ}\text{C}$. *Anal.* Calcd for $\text{C}_8\text{H}_5\text{N}_3$: C, 67.13; H, 3.52; N, 29.35. Found: C, 67.12; H, 3.23; N, 29.16. IR (KBr) cm^{-1} : 2218 (CN). $^1\text{H-NMR}$ (CDCl_3) δ : 8.25 (1H, d, $J = 5.8$, $\text{C}^5\text{-H}$), 7.78–6.78 (3H, m, $\text{C}^{6-8}\text{-H}$), 7.57 (1H, s, $\text{C}^1\text{-H}$).

3-Pyridinecarbonitrile (**5b**): The fraction eluted with CH_2Cl_2 gave **5b**.

1-Methyl-2-imidazolecarbonitrile (**5c**): Colorless oil (lit.,¹² $65\text{--}70^{\circ}\text{C}/0.4\text{ mm}$). IR (neat) cm^{-1} : 2232 (CN). MS (m/z): 107 (M^+).

1-Methyl-2-benzimidazolecarbonitrile (**5d**): The fraction eluted with *n*-hexane–benzene (1:1) gave **5d**. Colorless needles (benzene), mp $177\text{--}178^{\circ}\text{C}$. *Anal.* Calcd for $\text{C}_9\text{H}_7\text{N}_3$: C, 68.78; H, 4.49; N, 26.74. Found: C, 68.71; H, 4.34; N, 27.05. IR (KBr) cm^{-1} : 2234 (CN). $^1\text{H-NMR}$ (CDCl_3) δ : 7.85 (1H, d, $J = 8.1$, $\text{C}^4\text{-H}$), 7.53–7.39 (3H, m, $\text{C}^{5-7}\text{-H}$), 4.01 (3H, s, Me). MS (m/z): 157 (M^+).

2-Benzothiazolecarbonitrile (**5e**): The fraction eluted with *n*-hexane– CH_2Cl_2 (2:1) gave **5e**. Colorless needles (*n*-hexane–benzene), mp 82°C . *Anal.* Calcd for $\text{C}_8\text{H}_4\text{N}_2\text{S}$: C, 59.98; H, 2.52; N, 17.49. Found: C, 59.79; H, 2.21; N, 17.67. IR (KBr) cm^{-1} : 2224 (CN). $^1\text{H-NMR}$ (CDCl_3) δ : 8.40–7.85 (2H, m, C^4 and $\text{C}^7\text{-H}$), 7.80–7.40 (2H, m, C^5 and $\text{C}^6\text{-H}$). MS (m/z): 160 (M^+).

2-Benzofurancarbonitrile (**5f**): The fraction eluted with *n*-hexane– CH_2Cl_2 (2:1) gave **5f**. Colorless needles (*n*-hexane–AcOEt), mp 36°C (lit.,¹³ 36°C). IR (neat) cm^{-1} : 2228 (CN). $^1\text{H-NMR}$ (CDCl_3) δ : 7.77–7.15 (5H, m, $\text{C}^{3-7}\text{-H}$). MS (m/z): 143 (M^+).

2-Benzothiophenecarbonitrile (**5g**): Colorless oil. IR (KBr) cm^{-1} : 2214 (CN). $^1\text{H-NMR}$ (CDCl_3) δ : 7.90–7.84 (3H, m, C^2 , C^4 and $\text{C}^7\text{-H}$), 7.56–7.45 (2H, m, C^5 and $\text{C}^6\text{-H}$). MS (m/z): 159 (M^+).

1-Benzenesulfonyl-2-indolecarbonitrile (**5h**): The fraction eluted with *n*-hexane– CH_2Cl_2 (1:2) gave **5h**.

Colorless needles (*n*-hexane), mp 126–128 °C. *Anal.* Calcd for $C_{15}H_{10}N_2O_2S$: C, 63.82; H, 3.57; N, 9.92. Found: C, 63.78; H, 3.28; N, 9.72. IR (KBr) cm^{-1} : 2220 (CN). 1H -NMR ($CDCl_3$) δ : 8.31–7.80 (3H, m, C^3 , C^4 and C^7 -H), 7.70–7.20 (7H, m, C^5 , C^6 -H and Ph).

2,6-Dimethoxybenzonitrile (**6j**): Colorless needles (*n*-hexane–acetone), mp 118 °C (lit.,¹⁴ 118 °C). IR (KBr) cm^{-1} : 2214 (CN). 1H -NMR ($CDCl_3$) δ : 7.44 (1H, t, J = 8.4, C^4 -H), 6.56 (2H, d, J = 8.4, C^3 and C^5 -H).

2-Dimethylaminobenzonitrile (**6k**): Colorless oil (lit.,¹⁵ bp 58–65 °C/0.6 mm). IR (neat) cm^{-1} : 2210 (CN).

Reaction of 2-Lithiobenzothiazole with TsCN. A solution of *n*-BuLi in *n*-hexane (1.63 mol/L, 1.8 mL, 3.0 mmol) was slowly added to a solution of benzothiazole (405 mg, 3.0 mmol) in 6 mL of ether with stirring at –78 °C under an argon atmosphere and the solution was further stirred for 5 min. A solution of TsCN (815 mg, 4.5 mmol) in 4 mL of ether was added dropwise to the resulting solution at –78 °C and stirring was continued for 30 min. The cooling bath was removed and the reaction mixture was further stirred for 2 h, then poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography on SiO_2 with *n*-hexane and CH_2Cl_2 . The fraction eluted with *n*-hexane– CH_2Cl_2 gave di-2-benzothiazolyl ketone (**7**). Orange needles (*n*-hexane–acetone), mp 182–183 °C (lit.,¹⁶ 184 °C). IR (KBr) cm^{-1} : 1657 (CO). 1H -NMR ($CDCl_3$) δ : 8.38 (2H, d, J = 6.8, C^4 and C^4' -H or C^5 and C^5' -H), 8.06 (2H, d, J = 6.8, C^4 and C^4' -H or C^5 and C^5' -H), 7.67–7.57 (4H, m, C^{6-7} and $C^{6'-7'}$ -H). ^{13}C -NMR ($CDCl_3$) δ : 176.5 (CO), 162.2, 153.3, 137.9, 128.4, 127.4, 126.3, 122.2. MS (m/z): 296 (M^+).

Synthesis of Heteroarene carbonitriles (**5**) by Reaction of Trimethylsilyl heteroarenes (**8**) with TsCN.

1-Methyl-2-benzimidazole carbonitrile (5d**).** A solution of 1-methyl-2-trimethylsilylbenzimidazole (**8d**, 2.10 g, 10.3 mmol) and TsCN (1.86 g, 10.3 mmol) in THF (20 mL) was refluxed for 5 h under an argon atmosphere. The reaction mixture was poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on SiO_2 with *n*-hexane and CH_2Cl_2 . The fraction eluted with CH_2Cl_2 gave **5d** in 82% yield (1.33 g).

2-Benzothiazole carbonitrile (5e**).** A solution of 2-trimethylsilylbenzothiazole (**8e**, 400 mg, 1.9 mmol) and TsCN (350 mg, 1.9 mmol) in THF (10 mL) was refluxed for 5 h under an argon atmosphere. The reaction mixture was poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 , and concentrated. The residue was purified by column chromatography on SiO_2 with *n*-hexane and CH_2Cl_2 . The fraction eluted with *n*-hexane– CH_2Cl_2 (1:1) gave **5e** in 82% yield (250 mg).

2-Thiazole carbonitrile (5i**).** A mixture of 2-trimethylsilylthiazole (**8i**, 296 mg, 1.9 mmol) and TsCN (342 mg, 1.9 mmol) was heated at 70 °C for 5 h under an argon atmosphere. The reaction mixture was poured into ice- H_2O and extracted with CH_2Cl_2 . The organic layer was washed with H_2O , dried over Na_2SO_4 , and

concentrated. The residue was purified by column chromatography on SiO₂ with *n*-hexane and CH₂Cl₂. The fraction eluted with CH₂Cl₂ gave **5i** in 73% yield (153 mg). Colorless oil (lit.,¹⁷ bp 98 °C/24 mm). IR (neat) cm⁻¹: 2228 (CN). ¹H-NMR (CDCl₃) δ: 8.10 (1H, d, *J* = 2.7, C⁴-H), 7.80 (1H, d, *J* = 2.7, C⁵-H). MS (*m/z*): 110 (M⁺).

When above reaction was carried out in THF (10 mL), 2-thiazolecarbonitrile (**5i**) was obtained in 30% yield (62 mg).

2-Pyrrolicarbonitrile (10). A solution of methylmagnesium iodide (0.31 mol/L in ether, 9 mL, 2.8 mmol) was added to a solution of pyrrole (**9**, 134 mg, 2.0 mmol) in 6 mL of ether, and the whole was stirred for 30 min at rt. A solution of TsCN (434 mg, 2.4 mmol) in ether (4 mL) was added to the resulting solution and the mixture was further stirred for 2 h at -78 °C, then poured into ice-H₂O and extracted with CH₂Cl₂. The organic layer was washed with H₂O, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on SiO₂ with *n*-hexane and CH₂Cl₂. The fraction eluted with CH₂Cl₂ gave 2-pyrrolicarbonitrile (**10**) in 53% yield (97 mg). Colorless oil.¹⁸ IR (neat) cm⁻¹: 2218 (CN). ¹H-NMR (CDCl₃) δ: 9.07 (1H, br s, NH), 6.97 (1H, m, C³-H), 6.87 (1H, m, C⁵-H), 6.29 (1H, m, C⁴-H). MS (*m/z*): 92 (M⁺).

1-Indolecarbonitrile (12). Sodium hydride (NaH, 60% in oil, 144 mg, 3.6 mmol) was added to a solution of indole (**11**, 351 mg, 3.0 mmol) in 10 mL of THF at rt under an argon atmosphere, and the mixture was stirred for 10 min. TsCN (652 mg, 3.6 mmol) was further added to the resulting mixture, and the whole was stirred for 3 h at rt, then poured into ice-H₂O, and extracted with AcOEt. The organic layer was washed with H₂O, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on SiO₂ with *n*-hexane. The first fraction gave **12**. Colorless oil. IR (neat) cm⁻¹: 2242 (CN). ¹H-NMR (CDCl₃) δ: 7.80–7.15 (5H, m, C² and C^{4–7}-H), 6.65 (1H, d, *J* = 3.8, C³-H). MS (*m/z*): 142 (M⁺).

3-Indolecarbonitrile (13). A solution of methylmagnesium iodide (0.31 mol/L in ether, 9 mL, 2.8 mmol) was added to a solution of indole (**11**, 234 mg, 2.0 mmol) in 6 mL of ether, and the whole was stirred at rt for 30 min. A solution of TsCN (434 mg, 2.4 mmol) in ether (4 mL) was added to the resulting solution and the mixture was further stirred for 1 h at -50 °C, then poured into ice-H₂O and extracted with CH₂Cl₂. The organic layer was washed with H₂O, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography on SiO₂ with *n*-hexane and CH₂Cl₂. The fraction eluted with CH₂Cl₂ gave 3-indolecarbonitrile (**13**) in 93% yield (264 mg). Colorless needles (*n*-hexane), mp 178 °C (lit.,¹⁹ 180 °C). Anal. Calcd for C₉H₆N₂: C, 76.04; H, 4.25; N, 19.71. Found: C, 75.85; H, 4.06; N, 19.46. IR (neat) cm⁻¹: 2222 (CN). ¹H-NMR (DMSO-*d*₆) δ: 12.22 (1H, bs, NH), 8.26 (1H, d, *J* = 2.7, C²-H), 7.65 (1H, d, *J* = 7.3, C⁷-H), 7.57 (1H, d, *J* = 8.4, C⁴-H), 7.32–7.22 (2H, m, C⁵ and C⁶-H).

REFERENCES AND NOTES

1. This paper forms Part II of "Electrophilic Cyanations": Part I; A. Miyashita, I. Nagasaki, Y. Suzuki, K. Iwamoto, and T. Higashino, *Heterocycles*, 1997, **45**, 745.
2. (a) K. Friedrich and K. Wallenfels, " *The Chemistry of the Cyano Group*", 1970, 67, ed. by Z. Rappoport, The series of " *The Chemistry of Functional Groups*", John Wiley & Sons, London; (b) E. Hayashi, N. Shimada, and A. Miyashita, *Yakugaku Zasshi*, 1976, **96**, 1370; (c) A. Miyashita, T. Kawashima, C. Iijima, and T. Higashino, *Heterocycles*, 1992, **33**, 211; (d) C. Iijima, and A. Miyashita, *Chem. Pharm. Bull.*, 1990, **38**, 661; (e) A. Miyashita, Y. Suzuki, K. Ohta, and T. Higashino, *Heterocycles*, 1994, **39**, 345.
3. (a) C. Grundmann, *Methoden Org. Chem. (Houben-Weyl)*, 1985, **E5**, 1313, and 1441; (b) M. E. Kuehne, *J. Am. Chem. Soc.*, 1959, **81**, 5400; (c) P. J. Dudfield, C. T. Ekwuru, K. Hamilton, C. E. Osbourn, and D. J. Simpson, *Synlett*, 1990, 277.
4. (a) O. Fuents and W. W. Paudler, *J. Heterocycl. Chem.*, 1975, **12**, 375; (b) W. W. Paudler, C. I. P. Chao, and L. S. Helmick, *J. Heterocycl. Chem.*, 1972, **9**, 1157.
5. K. H. Bolze and H. D. Dell, *Liebigs Ann. Chem.*, 1967, **709**, 63.
6. W. P. Weber, " *Silicon Reagents for Organic Synthesis*". The series of " *Reactivity and Structure Concepts in Organic Chemistry*", Vol. 14, ed. by K. Hafner, J-M. Lehn, C. W. Rees, P. R. Schleyer, B. M. Trost, and R. Zahradnik, Springer-Verlag, New York, 1983.
7. R. A. Jones, " *Comprehensive Heterocyclic Chemistry*", ed. by A. R. Katritzky and C. W. Rees, Pergamon Press, Oxford, 1985, Vol. 4, 235.
8. P. Jacob and A. T. Shulgin, *Synthetic Comm.*, 1981, **11**, 957.
9. H. Gilman and J. W. Morton Jr., *Org. Reac.*, 1954, **8**, 287.
10. D. A. Shirley and M. D. Cameron, *J. Am. Chem. Soc.*, 1950, **72**, 2788.
11. D. W. Slocum, G. Book, and C. A. Jennings, *Tetrahedron Lett.*, 1970, 3443.
12. K. Yoshida and H. Kitabayashi, *Bull. Chem. Soc. Jpn.*, **60**, 1987, 3693.
13. L. Rene and R. Royer, *Bull. Soc. Chim. Fr.*, 1971, **12**, 4329.
14. (a) A. Russell and W. G. Tebbens, *Org. Synth.*, Coll. Vol. III, 293; (b) Mauthner, *J. Prakt. Chem.* (2), 1929, **121**, 259.
15. J. P. Idoux, J. T. Gupton, and C. Colon, *Synthetic Comm.*, 1982, **12**, 907.
16. E. Regal, *Liebigs Ann. Chem.*, 1977, 159.
17. T. Honma and Y. Tada, *Heterocycles*, 1977, **6**, 1985.
18. Y. Tamura, T. Kawasaki, M. Adachi, M. Tanio, and Y. Kita, *Tetrahedron Lett.*, 1977, 4417.
19. G. Mehta, *Synthesis*, 1978, 375.