

SYNTHESIS OF (*S*)-(+)- α -METHOXYMETHYLDIHYDRO-COUMARIN: ABSOLUTE CONFIGURATION OF THE REDUCED PRODUCT IN ENANTIOSELECTIVE RADICAL-MEDIATED REDUCTION†


Masatoshi Murakata, Hideyuki Tsutsui and Osamu Hoshino*

*Faculty of Pharmaceutical Sciences, Science University of Tokyo, Shinjuku-ku,
Tokyo 162, Japan*

Abstract — (*S*)-(+)- α -Methoxymethyldihydrocoumarin was synthesized starting from (*R*)-2-benzyl-3-hydroxypropyl acetate. Consequently, the absolute configuration of the reduced product obtained by radical-mediated reduction of α -iodo- α -methoxymethyldihydrocoumarin was determined to be *R*.


Asymmetric induction in radical-mediated reactions using organotin reagents has been currently focused in synthetic organic chemistry.¹ Although diastereoselective radical reactions have been reported, little has been known about the enantioselective radical reaction.² Recently,^{2a} we have reported the first example of an efficient enantioselective radical-mediated reduction of alkyl halide using organotin compound in the presense of a chiral ligand and a Lewis acid; the synthesis of optically active α -methoxymethyldihydrocoumarin (**2**) could be achieved by treatment of α -iodolactone (**1**) with tributyltin hydride coupled with a chiral amine and magnesium iodide as shown in Scheme 1. However, the absolute configuration of the reduced product (**2**) obtained by the asymmetric reaction was not determined, because **2** had not been known previously in optically active form. We describe here the synthesis of the title compound and the absolute configuration of the reduced product (**2**).

† This paper is dedicated to the memory of Professor Emeritus Shun-ichi Yamada.



Scheme 1

In our synthetic strategy, (*R*)-2-benzyl-3-hydroxypropyl acetate (**3**)³, which is prepared from 2-benzylpropanediol by lipase-catalyzed transesterification reported by Achiwa *et al.*, was chosen as a starting chiral synthon. The synthesis of (*S*)-(+)-**2** is shown in Scheme 2.



Scheme 2

The hydroxyl group of (*R*)-**3** [97% optical yield; $[\alpha]_D +28^\circ$ (*c*, 1.32, CHCl_3)] was protected with *t*-butyldimethylsilyl (TBDMS) group to form **4** in 94% yield, hydrolysis of which with LiOH in THF–H₂O gave alcohol (**5**) in 80% yield. For methylation of the hydroxyl group, at first, **5** was treated with NaH and MeI in DMF to give methyl ether (**6**). However, the specific rotation did not show significant value. This result suggested that racemization occurred by the treatment of **5** with NaH. Therefore, methylation under neutral conditions was considered. The reaction of **5** with MeI in MeCN in the presence of Ag₂O was carried out to afford desired optically active methyl ether (**6**) [$[\alpha]_D -11^\circ$ (*c*, 1.22, MeOH)] in 99%

yield. Deprotection of TBDMS group in **6** using Bu₄NF gave alcohol (**7**) {[α]_D +13° (c, 1.02, MeOH)} in 99% yield, Jones oxidation of which provided carboxylic acid (**8**) in 33% yield. Indanone (**9**) was obtained in 87% yield by the treatment of **8** with (COCl)₂ in the presence of a catalytic amount of DMF followed by Friedel-Crafts acylation using AlCl₃ in CH₂Cl₂. Finally, Baeyer-Villiger reaction of **9** with *m*-CPBA afforded (*S*)- α -methoxymethyl-dihydrocoumarin (**2**) in 57% yield. The enantiomeric excess (ee) was determined to be 40% by HPLC analysis using a chiral column, although partial racemization stage has not been clear. The specific rotation of (*S*)-**2** synthesized from (*R*)-**3** showed [α]_D +8° (c, 1.19, benzene). Thus, we concluded that *S*-enantiomer of α -methoxymethyldihydrocoumarin (**2**) has plus sign of rotation. Consequently, the absolute configuration of **2** {62% ee; [α]_D -12° (c, 1.01, benzene)} in the enantioselective radical-mediated reduction previously reported^{2a} was determined to be *R* (Scheme 1).

EXPERIMENTAL

All melting points were measured on a Yanagimoto (hot plate) melting point apparatus and are uncorrected. IR spectra were performed with a Hitachi 260-10 spectrophotometer and ¹H-NMR (270 MHz) and ¹³C-NMR (67.5 MHz) spectra were recorded with a JEOL EX-270 (270 MHz) spectrometer in CDCl₃ solution using tetramethylsilane as an internal standard. Mass spectra were measured on a Hitachi M-80, a JEOL JMS D-300 or a JEOL JMS-SX102A spectrometer. Column chromatography was performed on silica gel.

(*S*)-2-Benzyl-3-*tert*-butyldimethylsilyloxypropyl acetate (**4**)

To a solution of (*R*)-2-benzyl-3-hydroxypropyl acetate (**3**)³ {optical yield: 97%; [α]_D +28° (c, 1.32, CHCl₃)} (5 g, 24.0 mmol) in DMF (70 mL) were added imidazole (2.7 g, 39.7 mmol) and TBDMSCl (4 g, 26.5 mmol). The whole was stirred at rt for 2 h. The mixture was diluted with AcOEt (100 mL)-benzene (50 mL). The resulting mixture was washed successively with water (100 mL), 0.5N HCl (50 mL), water (50 mL), saturated NaHCO₃, and saturated NaCl, and dried over Na₂SO₄. The solvent was removed under reduced pressure, and the crude product was purified by distillation to afford **4** (7.3 g, 94%) as an oil: bp 140-145 °C / 4 mmHg; [α]_D +1° (c, 1.39, benzene); IR (neat) 2940, 2850, 1740, 1600 cm⁻¹; ¹H-NMR δ 0.03 (6H, s), 0.88 (9H, s), 2.02 (3H, s), 2.02-2.14 (1H, m), 2.58 (1H, dd, *J* = 7.1, 13.7 Hz), 2.69 (1H, dd, *J* = 7.8, 13.7 Hz), 3.49 (1H, dd, *J* = 5.3, 10.2 Hz), 3.55 (1H, dd, *J* = 4.8, 10.2 Hz), 4.03 (2H, d, *J* = 5.9 Hz), 7.13-7.26 (5H, m); ¹³C-NMR δ -5.6, -5.5, 20.9, 25.9, 34.2,

42.2, 61.8, 64.3, 126.0, 128.3, 129.1, 139.8, 171.1; MS m/z 307 ($M^+ - 15$). HRMS calcd for $C_{17}H_{27}O_3Si$ ($M^+ - 15$) 307.1727, found 307.1706.

(S)-2-Benzyl-3-tert-butyldimethylsilyloxypropanol (5)

A solution of acetate (4) (7.2 g, 22.2 mmol) and $LiOH \cdot H_2O$ (1.9 g, 44.3 mmol) in THF (45 mL)-water (20 mL) was refluxed for 25 h. The solvent was removed under reduced pressure, and the crude product was diluted with Et_2O (300 mL). The resulting mixture was washed successively with water (100 mL) and saturated NaCl, and dried over Na_2SO_4 . The solvent was removed under reduced pressure, and the crude product was purified by distillation to afford 5 (5 g, 80%) as an oil: bp 145 °C / 1.5 mmHg; $[\alpha]_D -18^\circ$ (c, 1.08, MeOH); IR (neat) 3600–3200, 2950, 2860, 1600 cm^{-1} ; 1H -NMR δ 0.03 (6H, s), 0.85 (9H, s), 1.90–2.03 (1H, m), 2.57 (2H, d, $J = 7.6$ Hz), 2.67 (1H, br), 3.54–3.63 (2H, m), 3.68–3.75 (2H, m), 7.12–7.24 (5H, m); ^{13}C -NMR δ -5.6, 18.2, 25.9, 34.2, 43.9, 65.8, 66.2, 126.0, 128.4, 129.0, 140.1; MS m/z 280 (M^+). HRMS calcd for $C_{16}H_{28}O_2Si$ (M^+) 280.1856, found 280.1835.

(S)-2-Benzyl-3-tert-butyldimethylsilyloxypropyl methyl ether (6)

To a solution of alcohol (5) (2.1 g, 7.5 mmol) in MeCN (5 mL) were added MeI (4 mL, 64 mmol) and Ag_2O (3 g, 12.9 mmol). The whole was refluxed for 18 h. After filtration, the solvent was removed under reduced pressure. The crude product was purified by column chromatography (hexane : benzene = 2 : 1, then benzene) followed by bulb-to-bulb distillation to give 6 (2.2 g, 99%) as an oil: bp 140 °C / 2 mmHg; $[\alpha]_D -11^\circ$ (c, 1.22, MeOH); IR (neat) 2930, 2850, 1600 cm^{-1} ; 1H -NMR δ 0.04 (6H, s), 0.88 (9H, s), 1.90–2.06 (1H, m), 2.58 (1H, dd, $J = 7.1, 13.5$ Hz), 2.66 (1H, dd, $J = 7.6, 13.5$ Hz), 3.28 (3H, s), 3.28 (2H, d, $J = 5.6$ Hz), 3.50 (1H, dd, $J = 5.4, 9.9$ Hz), 3.54 (1H, dd, $J = 5.3, 9.9$ Hz), 7.14–7.25 (5H, m); ^{13}C -NMR δ -5.5 (SiC and SiCH₃), 25.9, 34.2, 43.4, 58.8, 62.2, 72.3, 125.8, 128.2, 129.2, 140.6; MS m/z 294 (M^+). HRMS calcd for $C_{17}H_{30}O_2Si$ (M^+) 294.2013, found 294.1988.

(R)-2-Benzyl-3-methoxypropanol (7)

A solution of ether (6) (2.2 g, 7.3 mmol) and $Bu_4NF \cdot xH_2O$ (4 g) in THF (50 mL) was stirred for 2 h. The solvent was removed under reduced pressure, and the crude product was purified by column chromatography (hexane : AcOEt = 2 : 1, then 1 : 1) followed by bulb-to-bulb distillation to afford 7 (1.3 g, 99%) as an oil: bp 160 °C / 2 mmHg; $[\alpha]_D +13^\circ$ (c, 1.02, MeOH); IR (neat) 3600–3200, 2900, 1600 cm^{-1} ; 1H -NMR δ 2.05–2.18 (1H, m), 2.52 (1H, br), 2.61 (1H, dd, $J = 7.3, 13.5$ Hz), 2.67 (1H, dd, $J =$

7.6, 13.5 Hz), 3.34 (3H, s), 3.39 (1H, dd, $J = 6.9, 9.2$ Hz), 3.50 (1H, dd, $J = 4.3, 9.2$ Hz), 3.63 (1H, dd, $J = 6.6, 10.9$ Hz), 3.73 (1H, dd, $J = 3.6, 10.9$ Hz), 7.17–7.32 (5H, m); ^{13}C -NMR δ 34.9, 42.9, 59.6, 66.1, 76.2, 126.5, 128.9, 129.5, 140.4; MS m/z 180 (M^+). HRMS calcd for $\text{C}_{11}\text{H}_{16}\text{O}_2$ (M^+) 180.1148, found 180.1146.

(S)-2-Benzyl-3-methoxypropionic acid (8)

To a solution of alcohol (7) (680.6 mg, 3.78 mmol) in acetone (10 mL) was added dropwise Jones reagent at 0 °C until the solution turned brown. The reaction mixture was stirred until TLC analysis indicated complete oxidation of the alcohol (7). The mixture was quenched with *i*-PrOH and filtered. After addition of water, the filtrate was extracted with Et_2O (50 mL x 3). The organic extracts were washed with saturated NaCl, and dried over MgSO_4 . The solvent was removed under reduced pressure, and the crude product was diluted with Et_2O . The resulting mixture was extracted with saturated NaHCO_3 (50 mL x 3). The extracts were washed with Et_2O , and then acidified with 30% H_2SO_4 . The aqueous solution was extracted with Et_2O (50 mL x 3). The extracts were washed with saturated NaCl, and dried over MgSO_4 . The solvent was removed under reduced pressure to afford 8 (240.4 mg, 33%) as an oil: $[\alpha]_{\text{D}} -4^\circ$ (c , 1.05, c -hexane); IR (neat) 3700–2300, 1710, 1600 cm^{-1} ; ^1H -NMR δ 2.84 (1H, dd, $J = 6.9, 12.2$ Hz), 2.89–2.99 (1H, m), 3.04 (1H, dd, $J = 6.3, 12.2$ Hz), 3.35 (3H, s), 3.47–3.56 (2H, m), 7.18–7.33 (5H, m); ^{13}C -NMR δ 34.1, 47.2, 59.0, 71.8, 76.5, 126.6, 128.5, 128.9, 138.4, 178.4; MS m/z 194 (M^+). HRMS calcd for $\text{C}_{11}\text{H}_{14}\text{O}_3$ (M^+) 194.0943, found 194.0944.

(S)-2-Methoxymethylindanone (9)

To a solution of acid (8) (75.7 mg, 0.39 mmol) in CH_2Cl_2 (2 mL) were added $(\text{COCl})_2$ (0.05 mL, 0.57 mmol) and a catalytic amount of DMF (2 drops) at 0 °C. The whole was stirred at 0 °C for 30 min. The solvent was removed under reduced pressure, and the crude product was dissolved in CH_2Cl_2 (4 mL). To this solution was added AlCl_3 (98.4 mg, 0.74 mmol) at 0 °C. The reaction mixture was stirred at rt for 30 min. After addition of saturated NaHCO_3 , the mixture was extracted with Et_2O (50 mL x 3). The extracts were washed with saturated NaCl, and dried over Na_2SO_4 . The solvent was removed under reduced pressure, and the crude product was purified by column chromatography (hexane : $\text{Et}_2\text{O} = 4 : 1$) followed by bulb-to-bulb distillation to afford 9 (59.4 mg, 87%) as an oil: bp 123 °C / 2 mmHg; $[\alpha]_{\text{D}} -23^\circ$ (c , 1.41, benzene); IR (neat) 2900, 1705, 1600 cm^{-1} ; ^1H -NMR δ 2.88–2.96 (1H, m), 3.10–3.36 (2H, m), 3.34 (3H, s), 3.69–3.80 (2H, m), 7.33–7.39 (1H, m), 7.46–7.49 (1H, m), 7.56–7.62 (1H, m), 7.74–

7.77 (1H, m); ^{13}C -NMR δ 30.4, 47.9, 59.1, 72.3, 123.9, 126.6, 127.3, 134.9, 136.8, 154.3, 206.5; MS m/z 176 (M^+). HRMS calcd for $\text{C}_{11}\text{H}_{12}\text{O}_2$ (M^+) 176.0837, found 176.0845.

(S)-2-Methoxymethyldihydrocoumarin (2)

A solution of indanone (9) (50.1 mg, 0.29 mmol) and 70% *m*-CPBA (500 mg, 2.0 mmol) in CH_2Cl_2 (3 mL) was refluxed for 12 h. The solvent was removed under reduced pressure, and the crude product was taken up in Et_2O (30 mL). After filtration, the filtrate was diluted with hexane, and then washed with saturated NaHCO_3 and saturated NaCl , and dried over MgSO_4 . The solvent was removed under reduced pressure, and the crude product was purified by column chromatography (benzene) followed by bulb-to-bulb distillation to give 2 (31 mg, 57%) as an oil: bp $120^\circ\text{C} / 2\text{ mmHg}$; 40% ee; HPLC [chiralcel OB; hexane : 2-propanol = 50 : 1; flow rate 0.5 mL / min]; $[\alpha]_{\text{D}}^{+8}$ (c , 1.19, benzene); IR (neat) 2925, 1760, 1600 cm^{-1} ; ^1H -NMR δ 2.89–3.17 (3H, m), 3.41 (3H, s), 3.68 (1H, dd, $J = 6.9, 9.6\text{ Hz}$), 3.85 (1H, dd, $J = 4.1, 9.6\text{ Hz}$), 7.03–7.26 (4H, m); ^{13}C -NMR δ 27.0, 39.7, 59.2, 70.7, 116.6, 122.5, 124.4, 128.2 ($\text{CH} \times 2$), 151.5, 168.9; MS m/z 192 (M^+). HRMS calcd for $\text{C}_{11}\text{H}_{12}\text{O}_3$ (M^+) 192.0785, found 192.0788. These spectral data were identical with those for the reduced product obtained by enantioselective radical reaction.^{2a}

REFERENCES AND NOTES

1. Reviews: N. A. Porter, B. Giese, and D. P. Curran, *Acc. Chem. Res.*, 1991, **24**, 296; W. Smadja, *Synlett*, 1994, 1 and references cited therein.
2. Reports on enantioselective radical reactions: (a) M. Murakata, H. Tsutsui, and O. Hoshino, *J. Chem. Soc., Chem. Commun.*, 1995, 481. (b) H. Urabe, K. Yamashita, K. Suzuki, K. Kobayashi, and F. Sato, *J. Org. Chem.*, 1995, **60**, 3576. (c) J. H. Wu, R. Radinov, and N. A. Porter, *J. Am. Chem. Soc.*, 1995, **117**, 11029. (d) M. Nishida, H. Hayashi, A. Nishida, and N. Kawahara, *J. Chem. Soc., Chem. Commun.*, 1996, 579. (e) M. P. Sibi and J. Ji, *J. Am. Chem. Soc.*, 1996, **118**, 9200.
3. The synthesis of (*R*)-(+)-2-benzyl-3-hydroxypropyl acetate using lipase has been reported: lipase-catalyzed transesterification: K. Tsuji, Y. Terao, and K. Achiwa, *Tetrahedron Lett.*, 1989, **30**, 6189; lipase-catalyzed hydrolysis: K. Mori and N. Chiba, *Liebigs Ann. Chem.*, 1989, 957.

Received, 25th March, 1997