


SYNTHESIS AND REACTIONS OF CHIRAL CYCLIC NITRONES DERIVED FROM D-RIBOSE

Cedric W. Holzapfel* and Renier Crous

Department of Chemistry and Biochemistry, Rand Afrikaans University, P.O. Box 524, Auckland Park, 2006, South Africa

Abstract- A facile route to chiral cyclic nitrones derived from D-ribose is described. Their versatility as substrates for 1,3-dipolar cycloadditions and nucleophilic additions is demonstrated.


In the search for effective, non-toxic, antiviral agents a variety of modified nucleoside analogues are currently being developed.¹ Carbonucleosides,² thionucleosides³ and diheterosubstituted nucleosides⁴ have received much attention. As part of an ongoing investigation in our laboratories into the synthesis of analogues of biologically active compounds,⁵ we were interested in synthesising the aza analogues of β -C-nucleosides and β -C-glycosides.⁶ Our strategy involves the synthesis of a carbohydrate derived chiral cyclic nitrone (**1**) as the key synthon (Scheme 1). The rich chemistry displayed by these 1,3-dipoles⁷ makes them attractive targets, but the preparative methods for chiral cyclic nitrones of type (**1**) remain limited.



Scheme 1

Van den Broek⁸ has described the synthesis of cyclic nitrones *via* the oxidation of 1-deoxynojirimycin using 2,2-dimethyldioxirane. The oxidation was not regiospecific and separation of the isomers proved

to be tedious. Recently, the synthesis of pyrrolidine⁹ and piperidine based nitrones¹⁰ derived from carbohydrates was reported. The nitrone functionality was obtained by intramolecular conjugate addition of an oxime.¹¹ In both cases the nitrones were isolated as a mixture of diastereoisomers. Clearly, a stereospecific synthesis of these highly versatile intermediates is lacking. Herein we wish to report a facile route towards cyclic nitrones starting from suitably protected hemiacetals of D-ribofuranose. Readily available tri-*O*-benzyl-D-ribofuranose¹² was allowed to react with hydroxylamine hydrochloride to afford acyclic (**3**). Selective silylation followed by iodination at C-5 (with inversion of configuration) furnished the cyclisation precursor (**4**). Anhydrous TBAF-mediated desilylation and subsequent intramolecular nucleophilic attack¹³ afforded the cyclic nitrone (**5**)¹⁴ in an isolated yield of 86% (Scheme 2).

















Reagents: i) $\text{NH}_2\text{OH}\cdot\text{HCl}$ / pyridine / rt / 3 h ii) $\text{tBuPh}_2\text{SiCl}$ / pyridine / rt / 12 h
 iii) PPh_3 / imidazole / I_2 / toluene / reflux / 0.5 h iv) TBAF anhydrous
 / benzene / reflux / 0.5 h

Scheme 2


The 1,3-dipolar cycloaddition reaction of a nitrone to an alkene is an extremely powerful synthetic method for the creation of complex heterocyclic structures.¹⁵ For example, the labile nature of the N-O bond under mild reducing conditions in the formed isoxazolidines, makes them suitable precursors of 1,3-amino alcohols. Reaction of nitrone (**5**) with α,β -unsaturated carbonyl compounds furnished the corresponding cycloadducts (**6**, **7** and **8**)¹⁶ in good diastereomeric excess (Table 1). The *exo*-product was isolated as the major isomer in each case. The structures of all products were confirmed by NMR and MS analysis, and the stereochemistries by means of ROESY-spectra. The high diastereofacial selectivities observed were comparable to results reported by Ishikawa *et al.*⁹ for a different cyclic nitrone. The reaction of a variety of carbon nucleophiles, including a Grignard reagent, with **5** (Table 1) led to the formation of interesting β -C-glycoside analogues (**9**, **10**, **11** and **12**).¹⁷ The nitrone (**5**) was converted into the thymine C- nucleoside

Table 1: Reactions of **5** with dipolarophiles and carbon nucleophiles.

Nitrone	Reagent	Major Product	Yield ^a
 5		 6	92%^b (78% de)
		 7	95%^b (65% de)
		 8	86%^b (60% de)
	MeMgI	 9	83%^c
		 10	59%^d
		 11	75%^d
		 12	73%^d


- a) Total cycloadduct yield. b) Alkene (2 mmol) and nitrone (1 mmol) in dry benzene (2 mL) were reacted at 60°C for 3 h. c) Grignard reagent (1.1 mmol) and nitrone (1 mmol) in dry THF (2 mL) were reacted at 0°C for 10 min. d) Lithiated species (1.1 mmol) and nitrone (1 mmol) in dry THF (2 mL) were reacted at -30°C for 1 h.

analogue (13) using a method reported by Tronchet and co-workers¹⁸ (Scheme 3).



Scheme 3

Following the same protocol as for the synthesis of **5**, 2,3-isopropylidene-5-*O*-trityl-D-ribofuranose (**14**)¹⁹ was converted to the corresponding nitron (**15**).²⁰ The reaction of **15** with *N*-benzylmaleimide afforded cycloadduct (**16**)²¹ in a yield of 95% (diastereomeric excess 83%).



This work clearly shows that the construction of chiral cyclic nitrones from D-ribose derivatives is an extremely efficient and simple procedure. Furthermore, 1,3-dipolar cycloadditions of the nitrones to appropriate olefins proceed smoothly in high yielding diastereoselective reactions.

ACKNOWLEDGEMENTS

We thank the FRD (South Africa), AECI and SASOL for funding.

REFERENCES

- For reviews, see: E. De Clercq, *AIDS Res. Humn. Retroviruses*, 1992, **8**, 119; V. Nair and T.S. Jahnke, *Antimicrob. Agents Chemother.*, 1995, **39**, 1017; E. De Clercq, *Nucleosides Nucleotides*, 1994, **12**, 1271.

2. L. Agrofoglio, E. Suhas, A. Farese, R. Condom, S.R. Challand, R.A. Earl, and R. Guedji, *Tetrahedron*, 1994, **50**, 10611.
3. M.R. Dyson, O.L. Coe, and R.T. Walker, *J. Med. Chem.*, 1991, **34**, 2782.
4. D.W. Norbeck, S. Spanton, S. Border, and H. Mitsuya, *Tetrahedron Lett.*, 1989, **30**, 6263; B.R. Belleau, C.A. Evans, H.L.A. Tse, H. Jin, D.M. Dixit, and T.S. Mansour, *Tetrahedron Lett.*, 1992, **33**, 6949.
5. C.W. Holzapfel and M. Portwig, *Heterocycles*, 1997, **45**, 1433.
6. M. Yokoyama, T. Akiba, Y. Ochiai, A. Momotake, and H. Togo, *J. Org. Chem.*, 1996, **61**, 6079; G. Just and P. Donnini, *Can. J. Chem.*, 1977, **55**, 2998; B.A. Horenstein, R.F. Zabinski, and V.L. Schramm, *Tetrahedron Lett.*, 1993, **34**, 7213.
7. K.B.G. Torrsell, 'Nitrile Oxides, Nitrones, and Nitronates in Organic Synthesis', Chapter 3, VCH Publishers, New York, 1988, pp. 75-89.
8. L.A.G.M. Van den Broek, *Tetrahedron*, 1996, **52**, 4467.
9. T. Ishikawa, Y. Tajima, M. Fukui, and S. Saito, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1863.
10. P. Herczegh, I. Kovács, L. Szilágyi, T. Varga, Z. Dinya, and F. Sztaricskai, *Tetrahedron Lett.*, 1993, **34**, 1211.
11. P. Armstrong, R. Grigg, S. Surendrakumar, and W.J. Warnock, *J. Chem. Soc., Chem. Commun.*, 1987, 1327.
12. R. Barker and H.G. Fletcher Jr., *J. Org. Chem.*, 1961, **26**, 4605.
13. N.A. LeBel and N. Balasubramanian, *Tetrahedron Lett.*, 1985, **26**, 4331.
14. **5**: colourless oil; $[\alpha]_D +8.21^\circ$ ($c=2.1$, CHCl_3), ^1H NMR (300 MHz, CDCl_3): δ 7.31-7.27 (15H, m), 6.91 (1H, s), 4.69-4.36 (7H, m), 4.43 (1H, t, $J=5.4$ Hz), 4.12 (1H, dd, $J=10.7$ and 2.5 Hz), 4.08 (1H, m), 3.61 (1H, dd, $J=10.7$ and 2.1 Hz), ^{13}C NMR (300 MHz, CDCl_3): δ 137.61, 137.41, 137.23, 133.43, 128.63, 128.16, 128.11, 127.86, 127.68, 76.34, 75.35, 74.46, 73.41, 72.41, 72.06, 64.72, MS: m/z 418 (M^+).
15. For review, see: M. Frederickson, *Tetrahedron*, 1997, **53**, 403.
16. **6**: mp $63-64^\circ\text{C}$ (ethyl acetate/hexane); $[\alpha]_D +0.58^\circ$ ($c=2.4$, CHCl_3), ^1H NMR (300 MHz, CDCl_3): δ 7.32-7.27 (15H, m), 4.87 (1H, dt, $J=5.7$ and 2.0 Hz), 4.68-4.46 (6H, m), 4.34 (2H, d, $J=2.1$ Hz), 4.23 (1H, dd, $J=6.3$ and 2.4 Hz), 4.11 (1H, dd, $J=6.3$ and 4.5 Hz), 3.94 (1H, dd, $J=6.0$ and 2.4 Hz), 3.85 (1H, t, $J=4.2$ Hz), 3.56-3.41 (3H, m), ^{13}C NMR (300 MHz, CDCl_3): δ 177.47, 137.91, 137.72, 137.56, 128.51, 128.47, 128.42, 127.96, 127.89, 127.81, 78.66, 78.32, 76.68, 73.44, 72.74, 72.56, 71.14, 70.75, 69.92, 69.77, 50.57, MS: m/z 501 (M^+).

17. **11**: mp 84-85°C (ethyl acetate/hexane); $[\alpha]_D +5.93^\circ$ ($c=1.8$, CHCl_3), ^1H NMR (300 MHz, CDCl_3): δ 7.41 (1H, dd, $J=7.5$ and 1.8 Hz), 7.31-7.26 (16H, m), 6.88 (2H, m), 5.29 (1H, br s), 4.68 (1H, d, $J=4.8$ Hz), 4.65-4.43 (6H, m), 3.91 (1H, dd, $J=7.5$ and 5.7 Hz), 3.81 (1H, t, $J=5.5$ Hz), 3.77 (3H, s), 3.76 (1H, dd, $J=10.2$ and 3.9 Hz), 3.67 (1H, dd, $J=10.2$ and 4.2 Hz), 3.61 (1H, m), ^{13}C NMR (300 MHz, CDCl_3): δ 157.49, 138.48, 138.36, 138.23, 128.32, 128.25, 128.17, 127.99, 127.83, 127.59, 120.66, 110.63, 80.05, 75.29, 73.21, 71.62, 71.13, 71.04, 70.35, 69.11, 55.29, MS: m/z 525 (M^+).
18. J.M.J. Tronchet, G. Zosimo-Landolfo, M. Balkadjian, A. Ricca, M. Zsély, F. Barabalt-Rey, D. Cabrini, P. Lichtle, and M. Geoffroy, *Tetrahedron Lett.*, 1991, **32**, 4129.
19. N. Katagiri, K. Takashima, and T. Kato, *J. Chem. Soc., Perkin Trans. I*, 1983, 201.
20. **15**: colourless oil; $[\alpha]_D +2.22^\circ$ ($c=2.6$, CHCl_3), ^1H NMR (300 MHz, CDCl_3): δ 7.33-7.24 (15H, m), 7.09 (1H, s), 5.34 (1H, d, $J=6.3$ Hz), 4.52 (1H, d, $J=6.3$ Hz), 4.07 (1H, dd, $J=10.2$ and 2.7 Hz), 4.02 (1H, m), 3.13 (1H, dd, $J=10.2$ and 2.1 Hz), 1.42 (3H, s), 1.31 (3H, s), ^{13}C NMR (300 MHz, CDCl_3): δ 143.26, 133.31, 128.46, 128.11, 127.43, 111.79, 86.99, 79.19, 79.14, 77.32, 59.48, 27.26, 25.81, MS: m/z 430 (M^+).
21. **16**: colourless oil; $[\alpha]_D +0.45^\circ$ ($c=1.2$, CHCl_3), ^1H NMR (300 MHz, CDCl_3): δ 7.31-7.21 (20H, m), 4.81, (1H, t, $J=5.1$ Hz), 4.73 (2H, s), 4.70 (1H, d, $J=7.5$ Hz), 4.69 (1H, d, $J=5.1$ Hz), 4.19 (1H, d, $J=5.1$ Hz), 3.94 (2H, m), 3.24 (1H, dd, $J=9.9$ and 4.2 Hz), 3.14 (1H, dd, $J=9.9$ and 4.2 Hz), 1.42 (3H, s), 1.26 (3H, s), ^{13}C NMR (300 MHz, CDCl_3): δ 175.82, 175.11, 143.35, 135.27, 128.78, 128.59, 128.53, 127.99, 127.26, 113.08, 87.47, 85.94, 82.86, 75.68, 72.84, 71.59, 63.53, 50.24, 42.55, 26.31, 24.06, MS: m/z 616 (M^+).

Received, 9th March, 1998