

SYNTHESIS OF *N*-ARYLROLIPRAM DERIVATIVES - POTENT AND SELECTIVE PHOSPHODIESTERASE-IV INHIBITORS - BY COPPER CATALYZED LACTAM-ARYL HALIDE COUPLING

Esther Aebischer, Edmond Bacher, F. W. Joachim Demnitz,^{*1} Thomas H. Keller, Miriam Kurzmeyer, Marta L. Ortiz (in part),² Esteban Pombo-Villar, and Hans-Peter Weber


NOVARTIS Pharma AG, Preclinical Research, CH-4002 Basel, Switzerland

Abstract – The copper catalysed coupling of rolipram (**1**) with a wide variety of aryl halides (**2**) affords *N*-arylrolipram derivatives (**3**), potent and selective phosphodiesterase type-IV inhibitors.

Much attention has been paid to the therapeutic potential of phosphodiesterase (PDE) type-IV inhibitors for the treatment of asthma. This stems primarily from the wide-ranging anti-inflammatory properties exhibited by PDE-IV inhibitors *in vitro* and *in vivo*.³ The archetypal PDE-IV inhibitor *rolipram* (**1**) suppresses the proinflammatory function of a range of immune cells including mast cells, macrophages, lymphocytes and eosinophils.⁴

Structure-activity studies involving *rolipram* have implied that its receptor tolerates bulky substituents at the pyrrolidinone nitrogen.⁵ In order to exploit this finding we embarked upon the synthesis of a wide range of *N*-arylrolipram derivatives. The requirement of an expedient and general entry into this compound class was apparent. We report herein our results in preparing a large series of mono-, di- and tri-substituted *N*-arylroliprams (**3**). Due to the ready availability of *rolipram* on a large scale,⁶ we employed this compound as a starting material and found it an admirably suitable substrate in copper catalysed pyrrolidinone-aryl halide coupling.⁷ This proved to be a very versatile procedure for the synthesis of a wide range of *N*-arylrolipram derivatives in moderate to high yields (Table 1). *Rolipram* itself was very robust under the reaction conditions, often being partially recovered even after prolonged reaction at high temperature. Furthermore, we were satisfied to observe that subjection of optically pure *rolipram* enantiomers⁸ to the coupling conditions in the presence of 1-iodo-3-(3'-methoxybenzyloxy)benzene (**2i**) provided the corresponding *N*-arylrolipram derivative (**3i**) in good yield with no racemisation at the benzylic carbon (>99% optical purity by chiral HPLC).


Reactions were typically worked up upon completion of *rolipram* consumption, except in cases of prolonged reaction times (>12 h) where unchanged starting material was often recovered. In order to ensure good yields at acceptable reaction rates the appropriate aryl iodides were employed.

Table 1: Copper catalysed coupling of *rolipram* with aryl halides

Aryl Halide (2)					Conditions	Time (h) / Temp (°C)	Product ^o Yield (%)	MP (°C)
X	R ²	R ³	R ⁴	R ⁵				
Br	H	H	H	H	a)	3.5 / 150	3a (88)	124-126
I	H	OMe	H	OMe ¹⁰	b)	4.5 / reflux	3b (61)	52-54
I	H	OBn	H	OBn ¹¹	b)	3.5 / 150	3c (30)	116-119
Br	NO ₂	H	H	H	a)	5min / 150	3d (90)	117-119
Cl	NO ₂	OMe	H	OMe ¹²	a)	6 / 150	3e (32)	Foam
I	H	H	OMe	H	b)	6 / reflux	3f (16)	75-76
I	H	OBn	OMe	H	b)	4 / 150	3g (33)	88-89
I	H	OBn	H	H ¹³	b)	22 / 150	3h (50)	103-105
I	H	O(<i>m</i> -OMe)Bn	H	H ¹⁴	c)	4 / 140	(+)-3i (81) (-)-3i (81)	61-62 57-59
I	H	OBn	H	MeO ₂ C	a)	4 / 140	3j (59)	Foam
3-Bromo-5-benzoyloxy pyridine					a)	48 / 140	3k (74)	117-118
2-Bromo-4-benzoyloxy pyridine					a)	48 / 140	3l (54)	101-103
I	H	CO ₂ Me	H	H	b)	16 / reflux	3m (70)	129-131
Br	H	NO ₂	H	H	a)	20 / 150	3n (73)	99-101
Br	H	CHO	H	H	a)	22 / 135	3o (68)	Resin
I	H	O(<i>m</i> -OMe)Bn	H	O(<i>m</i> -OMe)Bn	b)	43 / 140	3p (75)	70-75
I	H	Br	H	H	c)	18 / 150	3q (75)	88-89

Conditions: a) Cu(1.85 eq)/K₂CO₃(1.46 eq)/KI(1.56 eq)/aryl halide(2-6 eq)/neat; b) Cu(0.16 eq)/K₂CO₃(1.3-1.4 eq)/aryl halide(1.3-1.5 eq)/DMF; c) Cu(17-22 eq)/K₂CO₃(1.5 eq)/aryl halide(0.9-1 eq)/DMF

Bromides, and to a lesser extent chlorides (2e), were only suitable in the case of bromobenzene (2a, no deactivating substituents on the benzene ring) or when electron withdrawing substituents were present in the aromatic ring. Even then coupling was sluggish, albeit in good yield (3n,o) unless the activating sub-


**Table 2:** Further derivatisation of *N*-arylrolipram derivatives

Arylrolipram			Conditions	Product			Yield (%)	MP (°C)
	R ²	R ³			R ²	R ³		
3d	NO ₂	H	H ₂ /10% Pd-C/MeOH	3r	NH ₂	H	100	159-161
3h	H	OBn	H ₂ /10% Pd-C/MeOH	3s	H	OH	100	158-160
3n	H	NO ₂	1) H ₂ /10% Pd-C/MeOH 2) ClCO(<i>p</i> -OMe)Ph	3t	H	NHCO(<i>p</i> -OMe)Ph	99	Foam
3o	H	CHO	Ph ₃ PCH ₂ (<i>p</i> -OMe)Ph/BuLi/THF	3u	H	<i>trans</i> -CH=CH(<i>p</i> -OMe)Ph	33*	163-165
3u			H ₂ /10% Pd-C/MeOH	3v	H	C ₂ H ₄ (<i>p</i> -OMe)Ph	27 (from 3o)	91-93
3s	H	OH	NaH/DMF/ClCO(<i>p</i> -OMe)Ph	3w	H	OCO(<i>p</i> -OMe)Ph	73	90-92
3s	H	OH	Cs ₂ CO ₃ /IBMK ^a /80° C/ ClCH ₂ (<i>p</i> -OMe)Ph	3x	H	OCH ₂ (<i>p</i> -OMe)Ph	57	59-61
3s	H	OH	Cs ₂ CO ₃ /IBMK/80° C/ BrCH ₂ CO(<i>m</i> , <i>p</i> -di-OMe)Ph	3y	H	OCH ₂ CO(<i>m</i> , <i>p</i> -di-OMe)Ph	80	Resin

* Isolated pure after chromatography. Reaction yield: 83%, 1:1 *cis:trans* mixture; ^a *iso*-butylmethyl ketone

tituent was *ortho* to the bromine. Thus 2-bromonitrobenzene (**2d**) reacted in 90% within 5 min with *rolipram*! Most of the other *rolipram* derivatives prepared had electron rich *N*-aryl substituents. The aryl iodide precursors provided the respective products (**3**) in good yields in these cases except when the deactivating aromatic methoxy group was in the *para* position (**2f,g**). Although not studied in detail, the amount of copper powder used as catalyst had a profound impact on both reaction time and yield, as shown by a comparison of cases **h** and **i**. The dihalide 3-bromoiodobenzene reacted, as expected, preferentially at the iodine-bearing carbon providing *N*-3'-bromophenylrolipram (**3q**) in good yield. Finally, many of the *N*-arylroliprams (**3**) shown in **Table 1** not only proved to be potent and selective PDE-IV inhibitors, but served as starting points for further derivatisation; (**Table 2**). *In vitro* PDE-IV inhibition results indicated, that bulky, lipophilic 3'-mono-, or 3',5'-disubstitution at the *rolipram* *N*-phenyl group significantly improved human neutrophil PDE-IV potency.¹⁵ Such derivatives were either prepared by direct halide coupling; (**Table 1**) or by manipulation of appropriate *N*-arylroliprams; (**Table**

2). However, unsymmetrical *N*-arylroliprams carrying a bulky, lipophilic group in one *meta* position and a polar group to improve aqueous solubility in the other, were particularly interesting from a pharmacological point of view. The synthesis of such derivatives, exemplified by **3za** and **3zb** having unsymmetrical *N*-phenyl 3',5'-disubstitution posed a selectivity problem. This could be nicely solved by the controlled partial hydrogenolysis of the symmetrical *di*-methoxybenzyl ether (**3p**) in ethyl acetate in acceptable yields,¹⁶ thus obviating the need for more circuitous routes which could not make use of the readily available 3,5-dihydroxyiodobenzene as a precursor.



d): H₂/10% Pd-C/EtOAc e): 1) Cs₂CO₃/IBMK/CIC₂H₄Cl/100°C 2) Me₂NH/EtOH 3) HCl; f): 1) K₂CO₃/BrCH₂CO₂Me 2) NaOH/H₂O/MeOH

Where not commercially available, aryl halide coupling substrates were prepared by standard procedures. 3,5-dimethoxy- (**2b**)¹⁰ and 3,5-dibenzyloxyphenyl iodide (**2c**)¹¹ are known and the 3,5-di-(3'-methoxybenzyloxy) derivative (**2p**) was prepared in an analogous fashion from 3,5-dihydroxyphenyl iodide by appropriate 3'-methoxybenzylation. The chloro-dimethoxynitrobenzene (**2e**)¹² was prepared by nitration of 3,5-dimethoxychlorobenzene. 3-Benzyloxy-4-methoxyiodobenzene (**2g**) was made from 3-hydroxy-4-methoxyaniline by diazotisation/potassium iodide followed by benzylation, whilst the benzyl ethers (**2h**)¹³ and (**2i**)¹⁴ and the methyl ester (**2m**) were prepared from the respective free hydroxy precursor. Finally, the bromopyridines (**2k**) and (**2l**) were synthesised without event using standard methodology from 5-amino-3-bromopyridine¹⁷ and 4-amino-2-bromopyridine respectively.¹⁸

In summary we have established the copper catalysed pyrrolidinone-aryl halide coupling to be a very useful and general preparation of *N*-aryl substituted derivatives. The reaction works well for phenyl- and pyridyl halide substrates bearing a variety of other aromatic substitution groups and has served as a quick entry by means of which we have been able to synthesise a considerable number of members of this compound class for pharmacological testing.

REFERENCES AND NOTES

1. New address: Departamento de Química Fundamental, Universidade Federal de Pernambuco, Cidade Universitária, CEP 50.670-901, Recife – PE, Brazil; FAX: +55 81 271 8442, e-mail: jdemnitz@npd.ufpe.br
2. *Praktikantin* from the University of Santiago de Compostela, Spain, 1993.
3. C. Schudt, G. Dent, and K. F. Rabe, eds., 'Phosphodiesterase Inhibitors', Academic Press, London, 1996; L. Sekut, D. Yarnall, S. A. Stimpson, L. S. Noel, R. Bateman-Fite, R. L. Clark, M. F. Brackeen, J. A. Menius, and K. M. Conolly, *Clin. Exp. Immunol.*, 1995, **100**, 126.
4. N. Sommer, P. A. Loeschmann, G. H. Northoff, M. Weller, A. Steinbrecher, J. P. Steinbach, R. Richtenfels, R. Meyermann, A. Reithmueller, A. Fontana, J. Dichgans, and R. Martin, *Nature Med.*, 1995, **1**, 244.
5. M. C. Marivet, J. J. Bourguignon, C. Lugnier, A. Mann, J. C. Stoclet, and C. G. Wermuth, *J. Med. Chem.*, 1989, **32**, 1450; P. W. Baures, D. S. Eggleston, S. Drake, K. F. Erhard, L. B. Cieslinski, Th. J. Torphy, and S. B. Christensen, *J. Med. Chem.*, 1993, **36**, 3274.
6. R. Backstrom, E. Honkanen, A. Raasmaja, and I. B. Linden, *PCT International Application* WO 9116303 A1 (*Chem. Abstr.*, 1992, **116**, 83532); K. Petzoldt, R. Schmiechen, and K. Hamp, *Ger. Offen.*, 3921593 A1 (*Chem. Abstr.*, 1991, **114**, 143134). We thank Dr. L. LaVecchia from the (former SANDOZ) kilo laboratory for providing large quantities of *rolipram*.
7. Y. Yamamoto and I. Kurate, *Can. J. Chem.*, 1983, **61**, 86; I. Goldberg, *Ber.*, 1907, **40**, 4541.
8. F. W. J. Demnitz, L. LaVecchia, E. Bacher, Th. Keller, Th. Mueller, F. Schürch, H.-P. Weber, and E. Pombo-Villar, *Molecules*, 1998, **3**, 107.
9. All new compounds provided satisfactory NMR, MS and microanalytical data.
10. R. A. Benkeser, R. A. Hickner, D. I. Hoke, and O. H. Thomas, *J. Am. Chem. Soc.*, 1958, **80**, 5289.
11. P. N. Edwards, G. C. Crawley, and J. M. M. M. Girondeau, *EP*, 385662.
12. H. H. Hodgson and J. S. Wignall, *J. Chem. Soc.*, 1928, 329; H. H. Hodgson and W. E. Batty, *J. Chem. Soc.*, 1934, 1433.
13. F. B. Kipping and J. J. Wren, *J. Chem. Soc.*, 1957, 3246.
14. We thank Mr. Thomas Wegmann for carrying out this reaction.
15. E. Bacher, Ch. Boer, K. Bray-French, F. W. J. Demnitz, Th. Keller, L. Mazzoni, Th. Mueller, and Ch. Walker, *Bioorg. Med. Chem. Let.*, 1998, **41**, submitted.
16. The corresponding over-hydrogenolysed diphenol product could be recycled by 3-methoxybenzylation and renewed hydrogenolysis.
17. A. Marcinkow and E. Plazek, *Roczniki Chem.*, 1936, **16**, 136; H. J. den Hertog, F. R. Schepman, J. de Bruyn, and G. J. E. Thyse, *Rec. Trav. Chim. Pays-Bas*, 1950, **69**, 1281.
18. H. J. den Hertog, *Rec. Trav. Chim. Pays-Bas*, 1945, **64**, 85; H. J. den Hertog, C. R. Kolder, and W. P. Combé, *Rec. Trav. Chim. Pays-Bas*, 1951, **70**, 591.

Received, 5th August, 1998