

A ONE-STEP TRANSFORMATION OF (*S*)-1-BENZOYL-3-[(*E*)-DIMETHYLAMINOMETHYLIDENE]-5-METHOXYCARBONYL-PYRROLIDIN-2-ONE INTO QUINOLIZINYL- AND 2*H*-2-PYRANONYL-SUBSTITUTED ALANINE DERIVATIVES

Marko Škof, Jurij Svete,* and Branko Stanovnik*

Faculty of Chemistry and Chemical Technology, University of Ljubljana,
Aškerčeva 5, 1000 Ljubljana, Slovenia


Abstract - (*S*)-1-Benzoyl-3-[(*E*)-dimethylaminomethylidene]-5-methoxycarbonylpyrrolidin-2-one (**1**) was transformed in one step with various 1,3-dinucleophiles (**2-14**) into the corresponding *N*-benzoyl-3-quinolizinyll- (**15-18**) and *N*-benzoyl-3-(2*H*-2-pyranonyl)alanine methyl esters (**19-27**).

Due to their occurrence in nature, biological activity, and synthetic utility, there has been in the last few decades a significant interest for the synthesis of 3-heteroarylalanines.¹ Among various synthetic approaches, transformations of commercially available α -amino acids, such as serine, aspartic acid, and glutamic acid, are often used for the preparation of 3-heteroarylalanines.² Among them, only few preparations of pyranonylalanines have been reported.³ Recently, Young and coworkers reported on the synthesis of 3-pyrazolyl-, 3-isoxazolyl-, and 3-pyrimidinylalanines from (*S*)-3-formylpyroglutamic acid derivatives, using a 'ring switching' strategy.⁴ On the other hand, we have previously shown that 3-dimethylaminopropenoates can serve as versatile reagents for the preparation of a variety of heterocyclic systems.⁵⁻⁷ In continuation of our work in this field, we report on efficient, one step synthesis of novel *N*-benzoyl-3-heteroarylalanine esters (**15-27**) from easily available (*S*)-1-benzoyl-3-[(*E*)-dimethylaminomethylidene]-5-methoxycarbonylpyrrolidin-2-one (**1**) and by using an improved 'ring switching' strategy.

The starting compound, (*S*)-1-benzoyl-3-[(*E*)-dimethylaminomethylidene]-5-methoxycarbonylpyrrolidin-2-one (**1**), was prepared from L-pyroglutamic acid in three steps according to the procedure described previously.⁷ This was then treated with two types of 1,3-dinucleophiles: 2-(pyridinyl-2)acetic acid derivatives (**2-5**) and carbocyclic and heterocyclic 1,3-dicarbonyl compounds and their analogs (**6-14**). Treatment of **1** with *C,N*-dinucleophiles (**2-5**) afforded quinolizinyll substituted alanine esters (**15-18**),


while with *C,O*-dinucleophiles (**6-14**), 3-heteroarylalanines (**19-27**) with fused 2*H*-pyran-2-one heterocyclic residue were formed (Scheme 1).

Scheme 1



Thus, methyl 2-(pyridinyl-2)acetate (**2**), ethyl 2-(pyridinyl-2)acetate (**3**), 2-(pyridinyl-2)acetonitrile (**4**), methyl 2-(quinolinyl-2)acetate (**5**), cyclohexane-1,3-dione (**6**), indane-1,3-dione (**7**), 4-hydroxy-6-methyl-2*H*-pyran-2-one (**8**), 4-hydroxy-2*H*-benzo[*b*]pyran-2-one (**9**), 2,4-dihydropyridine (**10**), barbituric acid (**11**), 1,3-dimethylbarbituric acid (**12**), 1,3-diphenyl-1*H*,4*H*-pyrazol-5-one (**13**), and 3-methyl-1-phenyl-1*H*,4*H*-pyrazol-5-one (**14**) furnished, after treatment with **1** in refluxing acetic acid, the corresponding (*S*)-*N*-benzoyl-3-heteroarylalanine methyl esters with the following heteroaryl residues, 1-methoxycarbonyl-4-oxo-4*H*-quinoliziny-3 (**15**), 1-ethoxycarbonyl-4-oxo-4*H*-quinoliziny-3 (**16**), 1-cyano-4-oxo-4*H*-quinoliziny-3 (**17**), 1-methoxycarbonyl-4-oxo-4*H*-benzo[*c*]quinoliziny-3 (**18**), 2,5-dioxo-5,6,7,8-tetrahydro-2*H*-benzo[*b*]pyrany-3 (**19**), 2,5-dioxo-2*H*,5*H*-indeno[1,2-*b*]pyrany-3 (**20**), 2,5-dioxo-7-methyl-2*H*,5*H*-pyrano[4,3-*b*]pyrany-3 (**21**), 2,5-dioxo-2*H*,5*H*-benzo[*b*]pyrano[4,3-*b*]pyrany-3 (**22**), 2,5-dioxo-5,6-dihydro-2*H*-pyrano[3,2-*c*]pyridiny-3 (**23**), 2,4,7-trioxo-1,2,3,4-tetrahydro-7*H*-pyrano[2,3-*d*]pyrimidiny-6 (**24**), 1,3-dimethyl-2,4,7-trioxo-1,2,3,4-tetrahydro-7*H*-pyrano[2,3-*d*]pyrimidiny-6 (**25**), 1,3-diphenyl-6-oxo-1*H*,6*H*-pyrano[2,3-*c*]pyrazoly-5 (**26**), and 3-methyl-1-phenyl-6-oxo-1*H*,6*H*-pyrano[2,3-*c*]pyrazoly-5 (**27**), respectively (Scheme 2).

Scheme 2



EXPERIMENTAL

Melting points were taken on a Kofler micro hot stage. The ^1H NMR spectra and ^{13}C NMR spectra were obtained on a Bruker Avance DPX 300 (300 MHz) spectrometer with $\text{DMSO}-d_6$ and CDCl_3 as solvents and Me_4Si as internal standard. The microanalyses for C, H, and N were obtained on a Perkin-Elmer CHN Analyser 2400. The optical rotations were measured on a Perkin-Elmer 241 MC Polarimeter.

(*S*)-1-Benzoyl-3-[(*E*)-dimethylaminomethylidene]-5-methoxycarbonylpyrrolidin-2-one (**1**) was prepared according to the procedure described in the literature.⁷

General Procedure for the Preparation of (*S*)-*N*-Benzoyl-3-heteroarylalanine methyl esters (15-27): A mixture of (*S*)-1-benzoyl-3-[(*E*)-dimethylaminomethylidene]-5-methoxycarbonylpyrrolidin-2-one (**1**) (1 mmol), 1,3-dinucleophile (**2-14**) (1 mmol), and glacial acetic acid (4 mL) was heated at reflux temperature for 2-5 h. Volatile components were evaporated *in vacuo*, the solid residue crystallised from methanol, and the precipitate collected by filtration to give *N*-benzoyl-3-heteroarylalanine methyl ester (**15-27**).

(*S*)-*N*-Benzoyl-3-(1-methoxycarbonyl-4-oxo-4*H*-quinolizinyl-3)alanine methyl ester (15). This compound was prepared from methyl 2-(pyridinyl-2)acetate (**2**), reflux for 2 h; yield 68%; mp 223-225°C (from methanol); $[\alpha]_{\text{D}}^{23} = +35.8^\circ$ ($c = 0.84$, CHCl_3). ¹H NMR (300 MHz, DMSO- d_6): δ 3.05 (1H, dd, $J = 9.8, 13.4$ Hz, 3-Ha), 3.35 (1H, dd, $J = 4.8, 13.5$ Hz, 3-Hb), 3.66 (3H, s, OMe), 3.80 (3H, s, OMe), 4.81 (1H, ddd, $J = 4.8, 7.9, 9.7$ Hz, 2-H), 7.43-7.53 (4H, m, 3H-Ph 7'-H), 7.76-7.79 (2H, m, 2H-Ph), 7.88-7.89 (1H, m, 8'-H), 8.39 (1H, s, 2'-H), 8.85 (1H, d, $J = 7.8$ Hz, NH), 9.05 (1H, d, $J = 9.1$ Hz, 9'-H), 9.22 (1H, d, $J = 7.1$ Hz, 6'-H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 33.3, 52.6, 52.6, 52.9, 101.2, 115.4, 118.0, 123.9, 128.1, 128.9, 129.1, 132.3, 134.6, 135.1, 141.1, 144.2, 158.7, 165.8, 167.4, 172.9. *Anal.* Calcd for $\text{C}_{22}\text{H}_{20}\text{N}_2\text{O}_6$: C, 64.70; H, 4.94; N, 6.86. Found: C, 64.60; H, 4.80; N, 6.78.

(*S*)-*N*-Benzoyl-3-(1-ethoxycarbonyl-4-oxo-4*H*-quinolizinyl-3)alanine methyl ester (16). This compound was prepared from ethyl 2-(pyridinyl-2)acetate (**3**), reflux for 3 h; yield 90%; mp 207-209°C (from methanol); $[\alpha]_{\text{D}}^{23} = +27.2^\circ$ ($c = 1.12$, CHCl_3). ¹H NMR (300 MHz, DMSO- d_6): δ 1.28 (3H, t, $J = 7.2$ Hz, CH_3CH_2), 3.05 (1H, dd, $J = 10.0, 13.4$ Hz, 3-Ha), 3.36 (1H, dd, $J = 4.5, 13.6$ Hz, 3-Hb), 3.67 (3H, s, OMe), 4.26 (2H, q, $J = 7.2$, CH_2CH_3), 4.79 (1H, ddd, $J = 4.9, 7.9, 9.8$ Hz, 2-H), 7.40-7.47 (3H, m, 3H-Ph), 7.50-7.56 (1H, m, 7'-H), 7.77-7.80 (2H, m, 2H-Ph), 7.87 (1H, ddd, $J = 1.5, 6.8, 9.4$ Hz, 8'-H), 8.40 (1H, s, 2'-H), 8.88 (1H, d, $J = 7.9$ Hz, NH), 9.06 (1H, dt, $J = 1.1, 9.4$ Hz, 9'-H), 9.22 (1H, dt, $J = 1.1, 7.5$ Hz, 6'-H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 15.0, 33.2, 52.6, 52.9, 61.2, 101.5, 115.4, 117.9, 123.9, 128.1, 128.9, 129.1, 132.3, 134.6, 135.0, 141.0, 144.2, 158.7, 165.4, 167.3, 172.8. *Anal.* Calcd for $\text{C}_{23}\text{H}_{22}\text{N}_2\text{O}_6$: C, 65.39; H, 5.25; N, 6.63. Found: C, 65.20; H, 5.42; N, 6.55.

(*S*)-*N*-Benzoyl-3-(1-cyano-4-oxo-4*H*-quinolizinyl-3)alanine methyl ester (17). This compound was prepared from 2-(pyridinyl-2)acetonitrile (**4**), reflux for 3 h; yield 71%; mp 231-233°C (from methanol); $[\alpha]_{\text{D}}^{23} = -2.5^\circ$ ($c = 1.01$, CHCl_3). ¹H NMR (300 MHz, DMSO- d_6): δ 3.04 (1H, dd, $J = 9.8, 13.6$ Hz, 3-Ha),

3.30 (1H, dd, $J = 5.1, 13.4$ Hz, 3-Hb), 3.66 (3H, s, OMe), 4.84 (1H, ddd, $J = 5.1, 7.7, 9.8$ Hz, 2-H), 7.43-7.56 (4H, m, 3H-Ph and 7'-H), 7.77-7.80 (2H, m, 2H-Ph), 7.90-7.93 (2H, m, 8'-H and NH), 8.07 (1H, s, 2'-H), 8.80 (1H, d, $J = 7.9$ Hz, 9'-H), 9.16 (1H, dt, $J = 1.0, 7.2$ Hz, 6'-H). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 32.3, 51.4, 52.0, 82.8, 116.4, 117.2, 117.6, 122.6, 127.2, 128.2, 128.3, 131.5, 133.6, 135.2, 139.8, 144.5, 157.2, 166.5, 171.9. *Anal.* Calcd for $\text{C}_{21}\text{H}_{17}\text{N}_3\text{O}_4$: C, 67.19; H, 4.56; N, 11.19. Found: C, 67.16; H, 4.78; N, 11.24.

(*S*)-*N*-Benzoyl-3-(1-methoxycarbonyl-4-oxo-4*H*-benzo[*c*]quinoliziny-3)alanine methyl ester (18). This compound was prepared from methyl 2-(quinolinyl-2)acetate (5), reflux for 3 h; yield 71%; mp 245-246°C (from methanol); $[\alpha]_D^{23} = +44.1^\circ$ ($c = 0.99$, CHCl_3). ^1H NMR (300 MHz, DMSO- d_6): δ 3.04 (1H, dd, $J = 10.2, 13.6$ Hz, 3-Ha), 3.36 (1H, dd, $J = 4.9, 13.2$ Hz, 3-Hb), 3.68 (3H, s, OMe), 3.79 (3H, s, OMe), 4.91 (1H, ddd, $J = 4.5, 8.3, 9.4$ Hz, 2-H), 7.42-7.58 (3H, m, 3H-Ph), 7.60-7.67 (2H, m, 7'-H, 8'-H), 7.76-7.79 (2H, m, 2H-Ph), 7.86 (1H, dd, $J = 2.1, 7.4$ Hz, 9'-H), 7.90 (1H, d, $J = 9.4$ Hz, 10'-H), 8.15 (1H, s, 2'-H), 8.63 (1H, d, $J = 9.8$ Hz, 11'-H), 8.83 (1H, d, $J = 7.9$ Hz, NH), 9.36 (1H, dd, $J = 1.1, 7.9$ Hz, 6'-H). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 33.8, 52.1, 52.9, 52.9, 104.4, 120.9, 122.7, 123.2, 126.3, 127.2, 128.0, 128.1, 128.8, 129.1, 129.5, 132.3, 133.9, 134.7, 138.9, 144.6, 164.2, 166.1, 167.4, 172.8. *Anal.* Calcd for $\text{C}_{26}\text{H}_{22}\text{N}_2\text{O}_6$: C, 68.12; H, 4.84; N, 6.11. Found: C, 68.05; H, 4.88; N, 6.29.

(*S*)-*N*-Benzoyl-3-(2,5-dioxo-5,6,7,8-tetrahydro-2*H*-benzo[*b*]pyranyl-3)alanine methyl ester (19). This compound was prepared from cyclohexane-1,3-dione (6), reflux for 3 h; yield 59%; mp 185-188°C (from methanol); $[\alpha]_D^{23} = +2.6^\circ$ ($c = 0.78$, CHCl_3). ^1H NMR (300 MHz, DMSO- d_6): δ 2.02 (2H, deg tt, $J = 6.3, 6.3$ Hz, 7'-CH₂), 2.44 (2H, t, $J = 6.6$ Hz, 6'-CH₂), 2.83 (2H, t, $J = 6.6$ Hz, 8'-CH₂), 2.87 (1H, dd, $J = 10.2, 14.3$ Hz, 3-Ha), 3.06 (1H, dd, $J = 4.9, 14.3$ Hz, 3-Hb), 3.66 (3H, s, OMe), 4.76 (1H, ddd, $J = 4.7, 8.1, 10.2$ Hz, 2-H), 7.43-7.57 (3H, m, 3H-Ph), 7.73 (1H, s, 4'-H), 7.77-7.81 (2H, m, 2H-Ph), 8.77 (1H, d, $J = 8.3$ Hz, NH). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 19.7, 27.1, 31.7, 35.9, 50.7, 52.1, 113.7, 121.8, 127.2, 128.3, 131.6, 133.5, 137.2, 160.7, 166.8, 171.6, 173.8, 194.2. *Anal.* Calcd for $\text{C}_{20}\text{H}_{19}\text{NO}_6$: C, 65.03; H, 5.18; N, 3.79. Found: C, 64.92; H, 5.35; N, 3.87.

(*S*)-*N*-Benzoyl-3-(2,5-dioxo-2*H*,5*H*-indeno[1,2-*b*]pyranyl-3)alanine methyl ester (20). This compound was prepared from indane-1,3-dione (7), reflux for 3 h; yield 84%; mp 151-153°C (from methanol); $[\alpha]_D^{23} = -21.0^\circ$ ($c = 1.01$, CHCl_3). ^1H NMR (300 MHz, DMSO- d_6): δ 2.92 (1H, dd, $J = 10.2, 13.9$ Hz, 3-Ha), 3.12 (1H, dd, $J = 4.8, 13.7$ Hz, 3-Hb), 3.69 (3H, s, OMe), 4.82 (1H, ddd, $J = 4.9, 8.1, 10.0$ Hz, 2-H), 7.43-7.64 (8H, m, 3H-Ph and 5H-Het), 7.78-7.81 (2H, m, 2H-Ph), 8.79 (1H, d, $J = 8.3$ Hz, NH). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 33.5, 51.6, 53.0, 111.5, 121.1, 122.3, 124.2, 128.1, 129.2, 132.5, 133.0, 133.4, 134.4, 135.2,

135.9, 136.2, 161.6, 167.6, 172.4, 175.3, 188.3. *Anal.* Calcd for $C_{23}H_{17}NO_6$: C, 68.48; H, 4.25; N, 3.47. Found: C, 68.50; H, 4.28; N, 3.63.

(S)-N-Benzoyl-3-(2,5-dioxo-7-methyl-2H,5H-pyrano[4,3-b]pyranyl-3)alanine methyl ester (21). This compound was prepared from 4-hydroxy-6-methyl-2H-pyran-2-one (**8**), reflux for 2 h; yield 82%; mp 220–222°C; $[\alpha]_D^{23} = -21.3^\circ$ ($c = 0.77$, $CHCl_3$). 1H NMR (300 MHz, $DMSO-d_6$): δ 2.30 (3H, s, 7-Me), 2.91 (1H, dd, $J = 10.2, 13.9$ Hz, 3-Ha), 3.11 (1H, dd, $J = 4.5, 13.9$ Hz, 3-Hb), 3.68 (3H, s, OMe), 4.79 (1H, ddd, $J = 4.9, 7.9, 10.2$ Hz, 2-H), 6.63 (1H, s, 8'-H), 7.43–7.56 (3H, m, 3H-Ph), 7.76–7.80 (3H, m, 2H-Ph and 4'-H), 8.78 (1H, d, $J = 7.9$ Hz, NH). ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 19.8, 32.0, 50.7, 52.1, 98.7, 100.8, 122.5, 127.2, 128.3, 131.5, 133.6, 137.8, 159.5, 159.8, 164.8, 165.7, 166.6, 171.5. *Anal.* Calcd for $C_{20}H_{17}NO_7$: C, 62.66; H, 4.47; N, 3.65. Found: C, 62.29; H, 4.51; N, 3.45.

(S)-N-Benzoyl-3-(2,5-dioxo-2H,5H-benzo[b]pyrano[4,3-b]pyranyl-3)alanine methyl ester (22). This compound was prepared from 4-hydroxy-2H-benzo[b]pyran-2-one (**9**), reflux for 2 h; yield 93%; mp 220–222°C (from methanol); $[\alpha]_D^{23} = -52.6^\circ$ ($c = 0.88$, $CHCl_3$). 1H NMR (300 MHz, $DMSO-d_6$): δ 3.00 (1H, dd, $J = 10.1, 13.9$ Hz, 3-Ha), 3.18 (1H, dd, $J = 4.6, 13.9$ Hz, 3-Hb), 3.70 (3H, s, OMe), 4.85 (1H, ddd, $J = 4.9, 7.9, 10.1$ Hz, 2-H), 7.45–7.54 (5H, m, 3H-Ph, 8'-H, 9'-H), 7.77–7.81 (3H, m, 2H-Ph, 7'-H), 7.93 (1H, s, 4'-H), 8.00 (1H, dd, $J = 1.0, 6.7$ Hz, 10'-H), 8.82 (1H, d, $J = 8.0$ Hz, NH). ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 33.0, 51.6, 53.1, 104.3, 113.7, 118.0, 123.8, 125.5, 126.2, 128.2, 129.2, 132.4, 134.5, 135.2, 139.0, 153.6, 159.3, 160.0, 160.8, 167.6, 172.4. *Anal.* Calcd for $C_{23}H_{17}NO_7$: C, 65.87; H, 4.09; N, 3.34. Found: C, 65.77; H, 4.05; N, 3.34.

(S)-N-Benzoyl-3-(2,5-dioxo-5,6-dihydro-2H-pyrano[3,2-c]pyridinyl-3)alanine methyl ester (23). This compound was prepared from 2,4-dihydroxypyridine (**10**), reflux for 2 h; yield 80%; mp 276–279°C (from DMF/methanol); $[\alpha]_D^{23} = -62.9^\circ$ ($c = 1.12$, DMF). 1H NMR (300 MHz, $DMSO-d_6$): δ -ppm: 2.91 (1H, dd, $J = 10.2, 14.0$ Hz, 3-Ha), 3.12 (1H, dd, $J = 4.7, 13.8$ Hz, 3-Hb), 3.67 (3H, s, OMe), 4.80 (1H, ddd, $J = 4.5, 7.9, 9.8$ Hz, 2-H), 6.33 (1H, d, $J = 7.2, 8'$ -H), 7.43–7.58 (4H, m, 3H-Ph and 7'-H), 7.77–7.80 (2H, m, 2H-Ph), 7.88 (1H, s, 4'-H), 8.79 (1H, d, $J = 7.9$ Hz, NH), 11.91 (1H, s, 6'-H). ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 32.1, 50.8, 52.0, 97.3, 108.0, 121.4, 127.2, 128.2, 131.4, 133.6, 138.0, 138.1, 159.8, 160.2, 162.5, 166.5, 171.6. *Anal.* Calcd for $C_{19}H_{16}N_2O_6$: C, 61.96; H, 4.38; N, 7.61. Found: C, 61.66; H, 4.43; N, 7.52.

(S)-N-Benzoyl-3-(2,4,7-trioxo-1,2,3,4-tetrahydro-7H-pyrano[2,3-d]pyrimidinyl-6)alanine methyl ester (24). This compound was prepared from barbituric acid (**11**), reflux for 5 h; yield 69%; mp 273–275°C (from

methanol); $[\alpha]_D^{23} = -107.6^\circ$ ($c = 0.68$, DMF). ^1H NMR (300 MHz, DMSO- d_6): δ 2.84 (1H, dd, $J = 10.2$, 14.0 Hz, 3-Ha), 3.02 (1H, dd, $J = 4.7$, 13.7 Hz, 3-Hb), 3.67 (3H, s, OMe), 4.74 (1H, ddd, $J = 4.7$, 8.1, 9.8 Hz, 2-H), 7.43-7.57 (3H, m, 3H-Ph), 7.75 (1H, s, 4'-H), 7.78-7.81 (2H, m, 2H-Ph), 8.25 (1H, br s, NH), 8.74 (1H, d, $J = 8.3$ Hz, NH), 11.26 (1H, s, NH). ^{13}C NMR (75 MHz, DMSO- d_6): δ 31.5, 50.9, 52.1, 91.7, 114.6, 127.2, 128.3, 131.5, 133.6, 139.4, 148.9, 158.8, 159.3, 160.4, 166.7, 171.6. *Anal.* Calcd for $\text{C}_{18}\text{H}_{15}\text{N}_3\text{O}_7$: C, 56.11; H, 3.92; N, 10.90. Found: C, 55.90; H, 3.90; N, 10.89.

(S)-N-Benzoyl-3-(1,3-dimethyl-2,4,7-trioxo-1,2,3,4-tetrahydro-7H-pyrano[2,3-d]-pyrimidinyl-6)alanine methyl ester (25). This compound was prepared from 1,3-dimethylbarbituric acid (12), reflux for 3 h; yield 76%; mp 218-220°C (from methanol); $[\alpha]_D^{23} = -40.0^\circ$ ($c = 0.78$, CH_2Cl_2). ^1H NMR (300 MHz, DMSO- d_6): δ 2.90 (1H, dd, $J = 10.2$, 14.0 Hz, 3-Ha), 3.07 (1H, dd, $J = 4.5$, 13.9 Hz, 3-Hb), 3.20 (3H, s, N-Me), 3.37 (3H, s, N-Me), 3.78 (3H, s, OMe), 4.77 (1H, ddd, $J = 4.7$, 8.1, 10.2 Hz, 2-H), 7.43-7.57 (3H, m, 3H-Ph), 7.79-7.82 (2H, m, 2H-Ph), 7.93 (1H, s, 4'-H); 8.77 (1H, d, $J = 7.9$ Hz, NH). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 28.8, 29.7, 32.2, 51.8, 53.0, 92.9, 115.1, 128.2, 129.2, 132.4, 134.5, 141.3, 150.2, 158.5, 159.3, 159.6, 167.5, 172.6. *Anal.* Calcd for $\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_7$: C, 58.11; H, 4.63; N, 10.16. Found: C, 58.10; H, 4.69; N, 10.20.

(S)-N-Benzoyl-3-(1,3-diphenyl-6-oxo-1H,6H-pyrano[2,3-c]pyrazolyl-5)alanine methyl ester (26). This compound was prepared from 1,3-diphenyl-1H,4H-pyrazol-5-one (13), reflux for 3 h; yield 51%; mp 208-210°C (from methanol); $[\alpha]_D^{23} = +20.6^\circ$ ($c = 0.95$, CHCl_3). ^1H NMR (300 MHz, DMSO- d_6): δ 2.97 (1H, dd, $J = 10.4$, 13.7 Hz, 3-Ha), 3.17 (1H, dd, $J = 4.9$, 13.4 Hz, 3-Hb), 3.70 (3H, s, OMe), 4.81 (1H, ddd, $J = 4.7$, 8.1, 10.0 Hz, 2-H), 7.38-7.54 (7H, m, 7H-Ph), 7.59-7.65 (2H, m, 2H-Ph), 7.80-7.84 (4H, m, 4H-Ph), 7.87-7.90 (2H, m, 2H-Ph), 8.32 (1H, s, 4'-H), 8.82 (1H, d, $J = 8.0$ Hz, NH). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 33.3, 52.0, 53.0, 100.5, 117.1, 121.8, 127.6, 128.2, 128.5, 129.1, 129.9, 130.1, 130.5, 132.2, 132.4, 134.5, 137.3, 139.1, 145.8, 151.0, 160.1, 167.5, 172.6. *Anal.* Calcd for $\text{C}_{29}\text{H}_{23}\text{N}_3\text{O}_5$: C, 70.58; H, 4.70; N, 8.51. Found: C, 70.21; H, 4.93; N, 8.42.

(S)-N-Benzoyl-3-(3-methyl-1-phenyl-6-oxo-1H,6H-pyrano[2,3-c]pyrazolyl-5)alanine methyl ester (27). This compound was prepared from 3-methyl-1-phenyl-1H,4H-pyrazol-5-one (14), reflux for 3 h; yield 49%; mp 130-132°C (from methanol); $[\alpha]_D^{23} = +54.3^\circ$ ($c = 0.75$, CHCl_3). ^1H NMR (300 MHz, DMSO- d_6): δ 2.31 (3H, s, 3'-Me), 2.91 (1H, dd, $J = 9.4$, 13.9 Hz, 3-Ha), 3.10 (1H, dd, $J = 4.9$, 13.9 Hz, 3-Hb), 3.67 (3H, s, OMe), 4.76 (1H, ddd, $J = 5.4$, 7.7, 9.6 Hz, 2-H), 7.37-7.60 (6H, m, 6H-Ph), 7.76-7.81 (4H, m, 4H-Ph), 7.95 (1H, s, 4'-H), 8.78 (1H, d, $J = 7.6$ Hz, NH). ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 12.2, 33.6, 52.1, 53.0, 82.3,

102.1, 115.7, 121.2, 128.1, 129.2, 130.5, 132.5, 134.5, 137.3, 138.7, 145.1, 150.3, 160.6, 167.7, 172.6. *Anal.* Calcd for $C_{24}H_{21}N_3O_5$: C, 66.81; H, 4.91; N, 9.74. Found: C, 66.58; H, 5.06; N, 9.57.

ACKNOWLEDGEMENT

The financial support from the Ministry of Science and Technology of Slovenia is gratefully acknowledged.

REFERENCES AND NOTES

1. For a review see: P. Kolar, A. Petrič, and M. Tišler, *J. Heterocycl. Chem.*, 1997, **34**, 1067.
2. J. J. Nestor, Jr., B. L. Horner, T. L. Ho, G. H. Jones, and G. I. McRae, *J. Med. Chem.*, 1984, **27**, 320; B. Ebert, S. Lenz, L. Brehm, P. Bregnedal, J. J. Hansen, K. Fredriksen, K. P. Bøgesø, and P. Krogsgaard-Larsen, *J. Med. Chem.*, 1994, **37**, 878; W. C. Patt, R. W. Skeeane, and B. A. Steinbaugh, *Synth. Commun.*, 1990, **20**, 3097; K. Burger, M. Gold, H. Neuhauser, M. Rudolph, and E. Höß, *Synthesis*, 1992, 1145; J. Svete, B. Stanovnik, and M. Tišler, *J. Heterocycl. Chem.*, 1994, **31**, 1259; M. Škof, J. Svete, and B. Stanovnik, *J. Heterocycl. Chem.*, 1997, **34**, 853.
3. J. W. Tilley, P. Levitan, and R. V. Kierstead, *J. Heterocycl. Chem.*, 1979, **16**, 333; H. Döpp, S. Maurer, A. N. Sasaki, and H. Musso, *Liebigs Ann. Chem.*, 1982, 254; J. E. Baldwin, M. R. Spyvee, and R. C. Whitehead, *Tetrahedron Lett.*, 1994, **35**, 6575.
4. A. N. Bowler, P. M. Doyle, and D. W. Young, *J. Chem. Soc., Chem. Commun.*, 1991, 314; A. Dinsmore, P. M. Doyle, and D. W. Young, *Tetrahedron Lett.*, 1995, **36**, 7503; A. N. Bowler, A. Dinsmore, P. M. Doyle, and D. W. Young, *J. Chem. Soc., Perkin Trans. 1*, 1997, 1297.
5. For short reviews see: B. Stanovnik, *Molecules*, 1996, **1**, 123; B. Stanovnik, 'Methyl 2-Benzoylamino-3-dimethylaminopropenoate in the Synthesis of Heterocyclic Systems' in 'Progress in Heterocyclic Chemistry,' Vol 5, ed. by H. Suschitzky and E. F. V. Scriven, Pergamon Press, Oxford, 1993, pp. 34-53.
6. For recent publications see: L. Selič, S. Golič-Grdadolnik, and B. Stanovnik, *Helv. Chim. Acta*, 1997, **80**, 2418; J. Smodiš and B. Stanovnik, *Tetrahedron*, 1998, **54**, 9799; L. Pizzioli, B. Ornik, J. Svete, and B. Stanovnik, *Helv. Chim. Acta*, 1998, **81**, 231; L. Selič and B. Stanovnik, *Helv. Chim. Acta*, 1998, **81**, 1634; L. Selič, S. Strah, R. Toplak, and B. Stanovnik, *Heterocycles*, 1998, **47**, 1017.
7. M. Škof, J. Svete, B. Stanovnik, L. Golič, S. Golič-Grdadolnik, and L. Selič, *Helv. Chim. Acta*, 1998, **81**, 2332.

Received, 11th January, 1999