

REACTION OF *rel*-(4*R*,5*R*)-4-BENZOYLAMINO-5-PHENYL-3-PYRAZOLIDINONE WITH ALIPHATIC 1,3-DICARBONYL COMPOUNDS. A 'RING SWITCH' SYNTHESIS OF *rel*-(2*R*,3*R*)-3-PHENYL-3-(PYRAZOLYL-1)ALANINE ESTERS

Jurij Svete and Branko Stanovnik*


Faculty of Chemistry and Chemical Technology, University of Ljubljana,
Aškerčeva 5, 1000 Ljubljana, Slovenia

Abstract - *rel*-(2*R*,3*R*)-*N*-Benzoyl-3-phenyl-3-(pyrazolyl-1)alanine esters (**5-8**) were prepared by acid-catalysed treatment of *rel*-(4*R*,5*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**) with various aliphatic 1,3-dicarbonyl compounds (**2-4**).

In last few decades, several synthetic approaches for the preparation of heteroarylalanines have been developed.¹ Considerable attention has been paid to the synthesis of 3-(pyrazolyl-1)alanine (**1**), which was isolated from the semen of *Citrullus vulgaris* and was also used as constituent of highly potent renine inhibitors.^{2,3} A 'ring switch' strategy, introduced by Young and coworkers, is a versatile synthetic method, since it enables the preparation of alanines with various heteroaryl residues.^{4,5} Previously, we reported a preparation of various 3-heteroarylalanine esters and their homologs starting from aspartic acid, pyroglutamic acid, and substituted pyrazolo[1,2-*a*]pyrazolones.⁵⁻¹⁰


In continuation of our work in this field, we now report a novel one-step 'ring switch' preparation of *rel*-(2*R*,3*R*)-*N*-benzoyl-3-phenyl-3-(substituted pyrazolyl-1)alanine esters (**5-8**) from *rel*-(4*R*,5*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**). Thus, treatment of **1**, accessible in two steps from *N*-benzoylglycine,¹¹ with the following 1,3-dicarbonyl compounds: 1,1,3,3-tetraethoxypropane (**2**), 3-oxobutylaldehyde dimethyl acetal (**3**), and pentane-2,4-dione (**4**) in methanol in the presence of trifluoroacetic acid gave the corresponding *rel*-(2*R*,3*R*)-*N*-benzoyl-3-phenyl-3-(substituted pyrazolyl-1)alanine esters (**5-7**). Similar treatment of **1** with pentane-2,4-dione (**4**) in ethanol afforded the corresponding pyrazolylalanine ethyl ester (**8**). Reaction mechanism can be explained by initial formation of the corresponding azomethine imine (**9**) which is in equilibrium with non-dipolar bicyclic isomer (**10**). Intermediate (**10**) then undergoes nucleophilic attack of alcohol followed by elimination of water giving pyrazolylalanine derivative (**11**) as product. However, this proposed mechanism is only hypothetical, since we were so far not able to isolate any of the intermediates (**9**, **10**) (Scheme 1).

Scheme 1



Structures of compounds (**5-8**) were confirmed by spectral characterisations and elemental analyses. Theoretically, the reaction of pyrazolidinone (**1**) with 3-oxobutylaldehyde dimethyl acetal (**3**) could give two isomeric products (**6**) or (**12**) with methyl group attached to the position 5 or 3 in the pyrazole residue, respectively. However, only *N*-benzoyl-3-(5-methyl-1-pyrazolyl)alanine methyl ester (**6**) was formed under these reaction conditions. The structure of the pyrazole residue in compound (**6**) was determined by ¹³C NMR spectroscopy. The ¹³C NMR chemical shifts for 3'-CH₃, 5'-CH₃, 3'-C, and 5'-C are in agreement with the literature data for parent pyrazole systems.^{12,13} (Scheme 2).

Scheme 2

¹³C NMR chemical shifts. δ (ppm)

Compound	3'-CH ₃	5'-CH ₃	3'-C	5'-C	Lit.
1-methylpyrazole	-	-	139.6	130.6	12a
1,3-dimethylpyrazole	13.9	-	148.8	131.0	12a
1,5-dimethylpyrazole	-	11.9	138.7	139.1	12a
1,3,5-trimethylpyrazole	14.0	11.7	147.5	139.8	12a
	-	-	140.3	130.4	this paper
5					
	-	11.3	139.2	139.7	this paper
6					
	14.2	11.2	148.0	140.4	this paper
7					

EXPERIMENTAL

Melting points were taken on a Kofler micro hot stage. The ¹H NMR spectra and ¹³C NMR spectra were obtained by Varian E-360 (60 MHz) and Bruker Avance DPX 300 (300 MHz) spectrometers with CDCl₃

as solvent and Me_4Si as internal standard. The microanalyses for C, H, and N were obtained on a Perkin-Elmer CHN Analyser 2400. *rel*-(2*R*,3*R*)-4-Benzoylamino-5-phenyl-3-pyrazolidinone (**1**) was prepared according to the procedure described in the literature.¹¹

***rel*-(2*R*,3*R*)-*N*-Benzoyl-3-phenyl-3-(pyrazolyl-1)alanine methyl ester (**5**).** A mixture of *rel*-(2*R*,3*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**) (1.405 g, 5 mmol), 1,1,3,3-tetraethoxypropane (**2**) (1.320 g, 6 mmol), methanol (20 mL), and trifluoroacetic acid (0.5 mL) was heated at reflux temperature for 2 h. The resulting solution was cooled, left to stand at 20° for 7 days, and the precipitate collected by filtration to give **5**; yield 1.081 g (62%); mp 159-160°C (from methanol). ¹H NMR (300 MHz, CDCl_3): δ 3.66 (3H, s, OMe), 5.61 (1H, dd, J = 6.0, 7.9 Hz, 2-H), 6.03 (1H, d, J = 6.0 Hz, 3-H), 6.27 (1H, t, J = 2.1 Hz, 4'-H), 6.90 (1H, br d, J = 7.5 Hz, NH), 7.28-7.70 (12H, m, 10H-Ph, 3'-H, 5'-H). ¹³C NMR (75.5 MHz, CDCl_3): δ 53.1, 56.8, 66.8, 106.34, 127.5, 127.9, 128.9, 129.0, 129.1, 130.4, 132.3, 133.9, 137.1, 140.3, 167.5, 170.8. *Anal.* Calcd for $\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_3$: C, 68.75; H, 5.48; N, 12.03. Found: C, 68.38; H, 5.47; N, 11.83.

***rel*-(2*R*,3*R*)-*N*-Benzoyl-3-(5-methylpyrazolyl-1)-3-phenylalanine methyl ester (**6**).** A mixture of *rel*-(2*R*,3*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**) (1.405 g, 5 mmol), 3-oxobutylaldehyde dimethyl acetal (**3**) (0.792 g, 6 mmol), methanol (20 mL), and trifluoroacetic acid (1 mL) was stirred at 20° for 5 days. Volatile components were evaporated *in vacuo*, the residue triturated with ether (20 mL), and the precipitate collected by filtration to give **6**; yield 1.131 g (62%); mp 141-142°C (from *n*-heptane/benzene, 5:1). ¹H NMR (300 MHz, CDCl_3): δ 2.14 (3H, s, 5'-Me), 3.64 (3H, s, OMe), 5.52 (1H, dd, J = 6.0, 7.2 Hz, 2-H), 6.01 (1H, d, J = 6.0 Hz, 3-H), 6.03 (1H, m, 4'-H), 6.97 (1H, br d, J = 7.2 Hz, NH), 7.22-7.68 (11H, m, 10H-Ph, 3'-H). ¹³C NMR (75.5 MHz, CDCl_3): δ 11.3, 52.9, 57.2, 62.6, 106.1, 127.4, 127.7, 128.4, 128.9, 129.0, 132.3, 134.0, 137.9, 139.2, 139.7, 167.6, 170.8. *Anal.* Calcd for $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_3$: C, 69.41; H, 5.82; N, 11.56. Found: C, 69.09; H, 5.62; N, 11.44.

***rel*-(2*R*,3*R*)-*N*-Benzoyl-3-(3,5-dimethylpyrazolyl-1)-3-phenylalanine methyl ester (**7**).** A mixture of *rel*-(2*R*,3*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**) (1.405 g, 5 mmol), pentane-2,4-dione (**4**) (0.600 g, 6 mmol), methanol (20 mL), and trifluoroacetic acid (1 mL) was stirred at 20° for 2 h. Volatile components were evaporated *in vacuo*, the residue triturated with ether (20 mL), and the precipitate collected by filtration to give **7**; yield 1.656 g (88%); mp 140-141°C (from methanol). ¹H NMR (300 MHz, CDCl_3): δ 2.08 (3H, s, 5'-Me), 2.25 (3H, s, 3'-Me), 3.65 (3H, s, OMe), 5.45 (1H, dd, J = 6.0, 7.0 Hz, 2-H), 5.80 (1H, br s, 4'-H), 5.90 (1H, d, J = 5.7 Hz, 3-H), 7.03 (1H, br d, J = 6.8 Hz, NH), 7.02-7.68 (10H, m, 10H-Ph). ¹³C NMR (75.5 MHz, CDCl_3): δ 11.2, 14.2, 52.8, 57.5, 62.4, 105.8, 127.4, 127.7,

128.2, 128.8, 129.0, 132.2, 134.2, 138.2, 140.4, 148.0, 167.6, 170.7. *Anal.* Calcd for $C_{22}H_{23}N_3O_3$: C, 70.01; H, 6.14; N, 11.13. Found: C, 70.00; H, 6.13; N, 11.49.

***rel*-(2*R*,3*R*)-*N*-Benzoyl-3-(3,5-dimethylpyrazolyl-1)-3-phenylalanine ethyl ester (8).** A mixture of *rel*-(2*R*,3*R*)-4-benzoylamino-5-phenyl-3-pyrazolidinone (**1**) (0.281 g, 1 mmol), pentane-2,4-dione (**4**) (0.600 g, 1.2 mmol), anhydrous ethanol (3 mL), and trifluoroacetic acid (0.2 mL) was heated at reflux temperature for 30 min. Volatile components were evaporated *in vacuo*, the residue triturated with ether (3 mL), and the precipitate collected by filtration to give **8**; yield 0.189g (48%); mp 142-143°C (from *n*-heptane). 1H NMR (60 MHz, $CDCl_3$): δ 1.08 (3H, t, $J = 7.1$ Hz, CH_3CH_2), 2.09 (3H, s, 5'-Me), 2.23 (3H, s, 3'-Me), 4.28 (2H, q, $J = 7.2$ Hz, CH_2CH_3), 5.52 (1H, t, $J = 6.6$ Hz, 2-H), 5.87 (1H, s, 4'-H), 5.99 (1H, d, $J = 6.2$ Hz, 3-H), 7.08-7.91 (11H, m, 10H-Ph, NH). *Anal.* Calcd for $C_{23}H_{25}N_3O_3$: C, 70.57; H, 6.44; N, 10.73. Found: C, 70.26; H, 6.55; N, 10.62.

ACKNOWLEDGEMENT

The financial support by the Ministry of Science and Technology of Slovenia is gratefully acknowledged.

REFERENCES AND NOTES

1. For a review see: P. Kolar, A. Petrič, and M. Tišler, *J. Heterocycl. Chem.*, **1997**, *34*, 1067.
2. N. Sugimoto, H. Watanabe, and A. Ide, *Tetrahedron*, **1960**, *11*, 231; I. L. Finar and K. Utting, *J. Chem. Soc.*, **1960**, 5272; H. Reimlinger and J. F. M. Oth, *Chem. Ber.*, **1964**, *97*, 331; L. D. Arnold, R. G. May, and J. C. Vederas, *J. Am. Chem. Soc.*, **1988**, *110*, 2237; J. E. Baldwin, A. C. Spivey, and C. J. Schofield, *Tetrahedron Asymmetry*, **1990**, *1*, 881; M. Perez and R. Pleixats, *Tetrahedron*, **1995**, *51*, 8355.
3. S. H. Rosenberg, K. P. Spina, K. W. Woods, J. Polakowski, D. L. Martin, Z. Yao, H. H. Stein, J. Cohen, J. L. Barlow, D. A. Egan, K. A. Tricarico, W. R. Baker, and H. D. Kleinert, *J. Med. Chem.*, **1993**, *36*, 449; S. H. Rosenberg, K. P. Spina, S. L. Condon, J. Polakowski, Z. Yao, P. Kovar, H. H. Stein, J. Cohen, J. L. Barlow, V. Klinghofer, D. A. Egan, K. A. Tricarico, T. J. Perun, W. R. Baker, and H. D. Kleinert, *J. Med. Chem.*, **1993**, *36*, 460.
4. A. N. Bowler, P. M. Doyle, and D. W. Young, *J. Chem. Soc., Chem. Commun.*, **1991**, 314; A. Dinsmore, P. M. Doyle, and D. W. Young, *Tetrahedron Lett.*, **1995**, *36*, 7503; A. N. Bowler, A. Dinsmore, P. M. Doyle, and D. W. Young, *J. Chem. Soc., Perkin Trans. 1*, **1997**, 1297.

5. M. Škof, J. Svete, and B. Stanovnik, *Heterocycles*, **1999**, *51*, 1051.
6. J. Svete, B. Stanovnik, and M. Tišler, *J. Heterocycl. Chem.*, **1994**, *31*, 1259.
7. L. Jukić, U. Bratušek, M. Škof, J. Svete, and B. Stanovnik, *Chemistry of Heterocyclic Compounds*, **1996**, 1510.
8. U. Bratušek, I. Kejžar, J. Svete, and B. Stanovnik, *Acta Chim. Slov.*, **1996**, *43*, 105.
9. M. Škof, J. Svete, and B. Stanovnik, *J. Heterocycl. Chem.*, **1997**, *34*, 853.
10. A. Prešeren, J. Svete, and B. Stanovnik, *J. Heterocycl. Chem.*, **1999**, *36*, in print.
11. J. Svete, A. Prešeren, B. Stanovnik, L. Golič, and S. Golič-Grdadolnik, *J. Heterocycl. Chem.*, **1997**, *34*, 1323.
12. a) J. Elguero, C. Marzin, and J. D. Roberts, *J. Org. Chem.*, **1974**, *39*, 357. b) E. Gonzales, R. Faure, E.-J. Vincent, M. Espada, and J. Elguero, *Org. Magn. Reson.*, **1979**, *12*, 587. c) P. Cabildo, R. M. Claramunt, and J. Elguero, *Org. Magn. Reson.*, **1984**, *22*, 603.
13. For a review on ^{13}C NMR of pyrazoles see: M. Begtrup, G. Boyer, P. Cabildo, C. Cativiela, R. M. Claramunt, J. Elguero, J. I. Garcia, C. Toiron, and P. Vedsø, *Magn. Reson. Chem.*, **1993**, *31*, 107, and references cited therein.

Received, 23rd March, 1999