


AN EFFICIENT TOTAL SYNTHESIS OF ELLIPTICINE

Minoru Ishikura,* Ayako Hino, and Nobuya Katagiri

Faculty of Pharmaceutical Sciences,
Health Sciences University of Hokkaido,
Ishikari-Tobetsu, Hokkaido 061-0293, Japan






Abstract - A total synthesis of ellipticine could be attained through the palladium catalyzed tandem cyclization-cross-coupling reaction of indolylborate (**2b**) with vinyl bromide (**3**) as a key reaction.

Ellipticine, first isolated from *Ochrosia elliptica* Labill in 1959,¹ is a member of pyrido[4,3-*b*]carbazole, which displays potent antitumor activity.² Because of its cardiovascular toxicity and hemolysis, interest was shifted to the investigation of the structure-activity relationships of structural analogues of ellipticine, which promoted the pronounced development of synthetic methodologies to pyrido[4,3-*b*]carbazoles.³ As a part of our ongoing studies of synthetic application of indolylborate,⁴ the palladium catalyzed tandem cyclization-cross-coupling reaction with 1-methylindolylborate (**2a**) was previously developed for a concise construction of 6-methylellipticine derivatives.⁵ At this point, use of pyrido[4,3-*b*]carbazole bearing a removable protecting group at the 6-position (indole-1-*N*) is indispensable to complete the synthesis of ellipticine. This paper describes a successful use of this protocol for the total synthesis of ellipticine.



Requisite vinyl bromide (**3**) as (*E*)/(*Z*) mixture was prepared from a mixture of *cis*- and *trans*-crotyl alcohol via sequences reported previously.⁵

The cross-coupling protocol was successfully extended to the reaction of 1-(*tert*-butoxycarbonyl)indolylborate (**2b**), generated from indole (**1b**) *in situ*,⁶ with **3** using a 1:4 ratio of Pd₂ (dba)₃•CHCl₃ and Ph₃P, producing hexatriene (**4**)⁷ in 64%



Scheme

yield (Scheme). Irradiation of **4** with high-pressure mercury lamp in benzene gave rise to pyridocarbazole (**5**) as an oxidized form⁸ in 41% yield. Then, treatment of **5** with TFA in CH₂Cl₂ afforded **6** in 88 % yield, and the deprotection of carbobenzyloxy group in **6** by catalytic hydrogenation using 20% Pd(OH)₂ on carbon produced **7** in 90% yield. To complete the sequence, dehydrogenation of **7** was examined next. However, the dehydrogenation of **7** with MnO₂ in AcOEt under reflux or 10% Pd on carbon in decalin at 200°C was frustrated by fairly low yield of ellipticine, which reveals that the presence of free indole-1-NH would be problematic.⁹ For successful completion of synthesis of ellipticine, **8** bearing a Boc group at the indole-1-N derived from **5** by catalytic hydrogenation was subjected to the dehydrogenation with MnO₂ in AcOEt under reflux to give **9** in 65% yield. Finally, deprotection of the Boc group in **9** with TFA provided ellipticine in 84% yield.

In summary, a novel and concise route to ellipticine was realized based on the palladium catalyzed tandem cyclization-cross-coupling reaction of indolylborate (**2b**) with vinyl bromide (**3**) as a key step.

ACKNOWLEDGEMENTS

This work was supported in part by a Grant-in Aid for Scientific Research (No. 09672151) and a Grant-in Aid for High Technology Research Program from the Ministry of Education, Sciences, Sports, and Culture.

REFERENCES AND NOTES

1. S. Goodwin, A. F. Smith, and E. C. Horning, *J. Am. Chem. Soc.* 1959, **81**, 1903.
2. L. K. Dalton, S. Demerac, B. C. Elmes, J. W. Loder, J. M. Swan, and T. Teitei, *Aust. J. Chem.*, 1967, **20**, 2715; M. G. Ferlin, G. Chiarello, C. Marzano, E. Severin, F. Baccichetti, F. Carlasse, M. Simonato, and F. Bordin, *Il Farmaco*, 1998, **53**, 431; L. M. Shi, Y. Fan, T. G. Meyer, P. M. O'Connor, K. D. Paull, S. H. Friend, and J. N. Weinstein, *J. Chem. Inf. Comput. Sci.*, 1998, **38**, 189.
3. G. W. Gribble, "Advances in Heterocyclic Natural Product Synthesis: Synthetic Approached to the Ellipticine Alkaloids via Metalation and Cycloaddition Chemistry," Vol 1, ed. by W. H. Pearson, JAI Press Inc., Greenwich, 1990, p. 43; M. Alvarez and J. A. Joule, "The Chemistry of Heterocyclic Compounds: The Uleine-Ellipticine-Vallesamine Ngouniensine Group," Supplement to Vol. 25, part 4, ed. by J. E. Saxton, John Wiley & Sons, Ltd., Chichester, 1994, p. 261; S. Blechert, R. Knier, H. Schroers, and T. Wirth, *Synthesis*, 1995, 593; A. Godard, F. Marsais, N. Ple, F. Trecourt, A. Turck, and G. Queguiner, *Heterocycles*, 1995, **40**, 1055; H. Haider, K. Mereiter, and R. Wanko,

Heterocycles, 1995, **41**, 1445; M. Amat, S. Hadida, S. Sathyanarayana, and J. Bosch, *Tetrahedron Lett.*, 1996, **37**, 3071; Y. Miki, Y. Tada, N. Yanase, H. Hachiken, and K. Matsushita, *Tetrahedron Lett.*, 1996, **37**, 7753; R. A. Jones, J. Pastor, J. Siro, and T. N. Voro, *Tetrahedron*, 1997, **53**, 479; N. Harada, T. Kawaguchi, I. Inoue, M. Ohashi, K. Oda, T. Hashiyama, and K. Tsujihara, *Chem. Pharm. Bull.*, 1997, **45**, 134; Y. Miki, Y. Tada, and K. Matsushita, *Heterocycles*, 1998, **48**, 1593; Y. Erbun, S. Patir, and G. Okay, *J. Heterocycl. Chem.* 1998, **35**, 1445; M. T. Diaz, A. Cobas, E. Guitian, and L. Castedo, *Synlett*, 1998, 157; A. Poumaroux, Z. Bouaziz, H. Fillion, M. Domard, J. Giraud, and A. Petavy, *Chem. Pharm. Bull.*, 1999, **47**, 643; Y. Murakami, *Yakugaku Zasshi*, 1999, **119**, 35.

4. M. Ishikura and I. Agata, *Recent Res. Devel. in Organic Chem.*, 1997, **1**, 145.
5. M. Ishikura, T. Yaginuma, I. Agata, Y. Miwa, R. Yanada, and T. Taga, *Synlett*, 1997, 214.
6. M. Ishikura, Y. Matsuzaki, I. Agata, and N. Katagiri, *Tetrahedron*, 1998, **54**, 13929.
7. Hexatriene (**4**) as (E)/(Z) mixture: $^1\text{H-NMR}$ (CDCl_3): δ 1.30-1.50 (m, 3H), 1.61 (s, 9H), 1.90 (s, 3H), 2.32 (br s, 1H), 2.64 (br s, 1H), 3.50-3.80 (m, 3H), 4.40-4.55 (m, 1H), 5.17 (s, 2H), 5.10-5.25 (m, 1H), 7.17 (t, 1H, $J=7.3$ Hz), 7.23 (t, 1H, $J=7.3$ Hz), 7.30-7.45 (m, 6H), 8.14 (d, 1H, $J=8.3$ Hz). $^{13}\text{C-NMR}$ (CDCl_3): δ 13.4, 20.3, 28.0, 30.3, 42.8, 45.6, 67.1, 83.3, 106.4, 115.3, 120.1, 122.6, 123.3, 123.9, 125.1, 127.8, 127.9, 128.4, 129.6, 133.7, 135.1, 136.4, 136.8, 142.7, 149.9, 155.4.
8. Compound (**5**): mp 145°C (recrystallized from AcOEt). $^1\text{H-NMR}$ (CDCl_3): δ 1.67 (s, 9H), 2.31 (s, 3H), 2.70 (br s, 3H), 2.92 (br s, 2H), 3.75-3.83 (m, 2H), 4.78 (s, 2H), 5.21 (s, 2H), 7.30-7.45 (m, 7H), 8.08 (d, 1H, $J=8.3$ Hz), 8.12 (d, 1H, $J=7.8$ Hz). $^{13}\text{C-NMR}$ (CDCl_3): δ 15.1, 17.2, 27.3, 28.1, 41.2, 44.5, 67.1, 83.7, 114.9, 117.0, 122.4, 122.9, 123.5, 124.2, 126.0, 127.3, 127.9, 128.0, 128.5, 132.7, 136.9, 138.0, 140.8, 151.3, 155.4. UV λ_{nm} (CHCl_3): 240, 270, 290.
9. The dehydrogenation of 6-methyl (indole-N-Me) derivative of **7** with MnO_2 in AcOEt under reflux produced 6-methylellipticine in 62% yield (unpublished result by M. Ishikura, *et al.*).