

DIRECT ARYLATION OF 2-PYRIDONES; PHOTOSTIMULATED $S_{RN}1$ REACTION BETWEEN CESIUM PHENOXIDES AND CHLORO-2-PYRIDONES

Hiroyuki Higuchi, Masayo Hattori, and Shigeru Ohmiya,*

*Institute of Medicinal Chemistry, Hoshi University, Ebara 2-4-41, Shinagawa-ku,
Tokyo 142-8501, Japan*

E-mail: ohmiya@hoshi.ac.jp

Abstract ----- 4-Hydroxyphenyl- (**3** and **4**) and 6-hydroxyphenyl-1-methyl-2-pyridones (**6** and **7**) were obtained by the photostimulated reaction of 4-chloro- (**1**) and 6-chloro-1-methyl-2-pyridone (**5**), respectively, with variously substituted cesium phenoxides (**2**). The reactions were proposed to proceed *via* an $S_{RN}1$ mechanism.


INTRODUCTION

The 2-pyridone ring is an important structural element of a number of biologically and structurally interesting molecules.¹ 2-Pyridones are versatile synthetic intermediates for many alkaloids.² We have recently reported that irradiation of 2-pyridones in the presence of electron donors such as aliphatic amines³ or π -excessive heteroaromatic compounds (pyrroles and indoles)⁴ causes a single electron transfer (SET) to yield addition products, 4-substituted 3,4-dihydropyridin-2(1*H*)-ones and 6-substituted 3,6-dihydropyridin-2(1*H*)-ones. As part of our investigations on the photochemistry of 2-pyridones *via* a SET process, we have studied the photostimulated substitution reaction of halo-2-pyridones with phenoxides under the intention of synthesizing aryl-2-pyridone derivatives, some of which are known to have biological activities such as elastase inhibitor,^{1b} cardiac,⁵ and anti-inflammatory.⁶ There is no literature precedent for a direct arylation of 2-pyridones to our knowledge though the syntheses of this class of compounds have been investigated extensively.^{2a} We now report the photosubstitution reaction of 4-chloro- (**1**) and 6-chloro-1-methyl-2-pyridone (**5**) with the cesium salts (**2**) of phenol and its derivatives having an electron-withdrawing (EWG) or an electron-donating group (EDG). Preliminary experiments revealed that 3-chloro-1-methyl-2-pyridone did not take place any reaction and 5-chloro-1-methyl-2-pyridone gave the substitution products only in low yields though the starting 2-pyridone was completely consumed.

RESULTS AND DISCUSSION

The photoreaction was carried out by irradiation of a deaerated solution of chloro-1-methyl-2-pyridone (**1**) or

(5) and cesium phenoxides (2) in dimethyl sulfoxide (DMSO) with a 400W high-pressure mercury lamp through a Pyrex filter at room temperature. The reaction mixture was separated by column chromatography on silica gel with CH_2Cl_2 -MeOH solvent systems. The structures of the products obtained were established by spectroscopic methods.



Scheme

Irradiation of 4-chloro-1-methyl-2-pyridone (1) with variously substituted phenoxides (2) underwent substitution reaction to yield 4-hydroxyphenyl-1-methyl-2-pyridones (3 and 4) in which the hydroxyl group on the benzene ring was located either *ortho* (*ortho*-substitution) or *para* (*para*-substitution) to the newly formed C-C bond. Similarly, 6-hydroxyphenyl-1-methyl-2-pyridones (6 and/or 7) were formed by the photoreaction of 6-chloro-1-methyl-2-pyridone (5) with the phenoxides (2) (*C*-substitution). The latter reactions were accompanied by the formation of 6-phenoxy-1-methyl-2-pyridones (8) (*O*-substitution) except for the case with *p*-cyanophenoxide (2g). The results are summarized in Table 1.

The common features of the photoreaction of chloro-2-pyridones (1 and 5) with cesium phenoxides in DMSO are as follows: 1) the *ortho*-substitution was preferable to the *para*-substitution irrespective of the species (EWG or EDG) of a substituent on the phenoxides; 2) the presence of an EWG at the *ortho* position of phenoxides significantly reduced the formation of the substitution product; 3) in the photoreaction with 6-chloro-2-pyridone (5), the presence of an EWG on the phenoxides increased the ratio of *C*-substitution/*O*-substitution.

When a solution of phenoxide (2d) and 4-chloro-2-pyridone (1) in DMSO, which gave *C*-substitution product (3d) on irradiation, was treated in the dark, no reaction occurred. On the other hand, only the *O*-substitution product (8d) was obtained in 80% yield in the dark reaction of 6-chloro-2-pyridone (5) with 2d. The above results demonstrate that the *C*-substitution proceeded *via* a photochemical process and the *O*-substitution *via* a thermal process. Irradiation of phenoxy-2-pyridone (8d) in DMSO did not cause any reaction, indicating that the phenoxy-2-pyridone derivatives might not be an intermediate for the *C*-substitution. It has recently been reported that hydroxybiaryls are synthesized both photochemically⁷ and electrochemically⁸ starting from haloarenes and phenoxides or naphthoxides. These substitution reactions are proved to proceed *via* SRN1 mechanism. The photostimulated reaction of 1 with 2f was significantly inhibited in the presence of *p*-dinitrobenzene (*p*-DNB, 0.1 eq. to 1), a well-known inhibitor of SRN1 .

reactions (Table 1), in analogy with those of haloarenes and phenoxides.^{7,8} The formation of *C*-substitution product (**6d**) also decreased, in contrast to a significant increase of the *O*-substitution product (**8d**) by addition of the inhibitor to the photoreaction of **5** with **2d**. All these results let us to propose an $S_{RN}1$ mechanism for the photostimulated substitution reaction leading to the formation of hydroxyphenyl-2-pyridone derivatives.

Table 1. Photostimulated Reaction of Chloro-2-pyridones (**1** and **5**) with Phenoxides (**2**) in DMSO

2-pyridones	phenoxides	products (yields %) ^a			
1	2a; R=H ^c				
	2b; R=2-OMe	3b; R=3'-OMe (40)	4b; R=3'-OMe (19)		
	2c; R=3-OMe	3c; R=4'-OMe (8)	4c; R=2'-OMe (8)		
		3c'; R=6'-OMe (12)			
	2d; R=4-OMe	3d; R=5'-OMe (27)			
	2e; R=2-CN	3e; R=3'-CN (5)	4e; R=3'-CN (5)		
	2f; R=3-CN	3f; R=4'-CN (20)	4f; R=2'-CN (12)		
		(10) ^b	(6) ^b		
	2g; R=4-CN	3g; R=5'-CN (18)			
5	2a	6a; R=H (32)	7a; R=H (10)	8a; R=H (10)	
	2b		7b; R=3'-OMe (5)	8b; 2'-OMe (50)	
	2c	6c; 4'-OMe (20)		8c; 3'-OMe (20)	
	2d	6d; 5'-OMe (22)		8d; 4'-OMe (28)	
		(8) ^b		(82) ^b	
	2e ^c				
	2f	6f; R=6'-CN (40)		8f; R=3'-CN (25)	
	2g	6g; R=5'-CN (54)			

a, Isolated yields. Products (**3** and **4**, and **6a** and **7a**) were obtained as an inseparable mixture of regioisomers, ratios of which were estimated by ¹H-NMR spectra.

b, In the presence of *p*-dinitrobenzene (10 mol %).

c, Complex mixture.

EXPERIMENTAL

General Procedures

Melting points were determined on a Yanagimoto micro melting apparatus and are uncorrected. ¹H NMR spectra were measured in DMSO-*d*₆ unless otherwise noted, with a JEOL JNM-GX 400 or a JEOL JNM-GSX 270 spectrometer, and chemical shifts (δ) are given in parts per million (ppm) down field from tetramethylsilane as an internal standard. Coupling constants (*J*) are given in Hertz (Hz). EI-MS were taken on

JEOL D-300 instrument. IR spectra were measured with Hitachi 215 spectrophotometer at KBr discs. Analytical GLC was carried out with Hitachi 263-30 gas chromatograph equipped with a column (0.3 x 200 cm) of silicon OV-17 on Gas Chrom Q. GLCEI-MS were measured with the same instrument as used for EI-MS, and GLC conditions were the same as in the analytical GLC except for the use of He instead of N₂ as a carrier gas. A Riko 400W high-pressure mercury lamp was used as the irradiation sources. 6-Chloro-2-pyridinol and phenols were obtained from commercial sources and purified by distillation or recrystallization before use. 4-Chloro-2-pyridinol was synthesized by the reported method.⁹ **1** and **5** were prepared in 92% and 82% yields, respectively, by stirring a solution of the chloro-2-pyridinols (0.65 g, 5.0 mmol), methyl iodide (2.13 g, 15.0 mmol) and tetrabutylammonium iodide (0.18 g, 0.5 mmol) in benzene (10 mL) in the presence of anhydrous K₂CO₃ (2.07 g, 15.0 mmol) for 24 h at rt; **1**; mp 74-75 °C (sublimation); ¹H-NMR (CDCl₃) δ: 6.61 (1H, d, *J* = 2.4, 3-H), 6.19 (1H, dd, *J* = 7.3 and 2.4, 5-H), 7.23 (1H, d, *J* = 7.3, 6-H). **5**: needles from *n*-C₅H₁₂-Et₂O, mp 63-65 °C, that was identical with the reported value.¹⁰ The cesium salts of phenols were prepared analogously to the reported procedure.¹¹ DMSO was purified by distillation after being dried with CaH₂ and stored on molecular sieves (4Å).

General procedure for the photochemical reactions. A solution of the chloro-2-pyridones (0.6 mmol) and the cesium salts (3.0 mmol) in dry DMSO (6 mL) was flushed with N₂ for 15 min and then sealed under reduced pressure. The solution was irradiated externally for 7 h through a Pyrex filter with a 400W high-pressure mercury lamp. After removal of the solvent under reduced pressure, the residue was treated with water, 5% HCl and then extracted with CH₂Cl₂ several times. The combined extracts were dried on Na₂SO₄ and evaporated to dryness *in vacuo*. The residue was separated by repeated column chromatography on silica gel using CH₂Cl₂-MeOH solvent systems.

Photoreaction of 1 with 2b. Products (**3b** and **4b**) were obtained as an inseparable mixture (74 mg, 59%) of the ratio of 2:1. 4-(2-Hydroxy-3-methoxyphenyl)-1-methyl-2-pyridone (**3b**); GLC-MS *m/z*: 231 (M⁺, 100), 230, 214; ¹H-NMR δ: 3.58 (3H, s, NCH₃), 6.78 (1H, d, *J* = 1.8, 3-H), 6.44 (1H, dd, *J* = 7.3 and 1.8, 5-H), 7.28 (1H, d, *J* = 7.3, 6-H), 10.32 (1H, br, 2'-OH), 3.93 (3H, s, 3'-OCH₃), 6.91 (1H, d, *J* = 8.6, 4'-H), 6.93 (1H, dd, *J* = 8.6 and 8.6, 5'-H), 6.95 (1H, d, *J* = 8.6, 6'-H). Irradiation at the singlet (δ 3.93) due to the OCH₃ group caused enhancement of the doublet at δ 6.91 (4'-H). 4-(4-Hydroxy-3-methoxyphenyl)-1-methyl-2-pyridone (**4b**); GLC-MS *m/z*: 231 (M⁺, 100), 230, 214; ¹H-NMR δ: 3.56 (3H, s, NCH₃), 6.86 (1H, d, *J* = 1.8, 3-H), 6.58 (1H, dd, *J* = 7.3 and 1.8, 5-H), 7.34 (1H, d, *J* = 7.3, 6-H), 7.07 (1H, d, *J* = 1.8, 2'-H), 3.94 (3H, s, 3'-OCH₃), 10.4 (1H, br, 4'-OH), 6.98 (1H, d, *J* = 8.6, 5'-H), 7.13 (1H, dd, *J* = 8.6 and 1.8, 6'-H). Irradiation at the singlet (δ 3.94) due to the OCH₃ group induced enhancement of the doublet at δ 7.07 (2'-H).

Photoreaction of 1 with 2c. Products (**3c**, **3c'** and **4c**) were obtained as an inseparable mixture (40 mg, 28%) of the ratio of 2:3:2. 4-(2-Hydroxy-4-methoxyphenyl)-1-methyl-2-pyridone (**3c**); GLC-MS *m/z*: 21 (M⁺, 100), 230, 214; ¹H-NMR δ: 3.41 (3H, s, NCH₃), 6.42 (1H, d, *J* = 1.8, 3-H), 6.33 (1H, dd,

$J=7.3$ and 1.8 , 5-H), 7.58 (1H, d, $J=7.3$, 6-H), 9.81 (1H, br, 2'-OH), 6.48 (1H, d, $J=1.8$, 3'-H), 3.75 (3H, s, 4'-OCH₃), 6.53 (1H, dd, $J=7.9$ and 1.8 , 5'-H), 7.26 (1H, d, $J=7.9$, 6'-H). 4-(2-Hydroxy-6-methoxyphenyl)-1-methyl-2-pyridone (**3c'**): GLC-MS m/z : 231 (M^+ , 100), 230, 214; $^1\text{H-NMR}$ δ : 3.43 (3H, s, NCH₃), 6.23 (1H, d, $J=1.8$, 3-H), 6.08 (1H, dd, $J=7.3$ and 1.8 , 5-H), 7.56 (1H, d, $J=7.3$, 6-H), 9.55 (1H, br, 2'-OH), 6.57 (1H, d, $J=7.9$, 3'-H), 7.15 (1H, dd, $J=7.9$ and 6.7 , 4'-H), 6.55 (1H, d, $J=6.7$, 5'-H), 3.68 (3H, s, 6'-OCH₃). 4-(4-Hydroxy-2-methoxyphenyl)-1-methyl-2-pyridone (**4c**); GLC-MS m/z : 231 (M^+ , 100), 230, 214; $^1\text{H-NMR}$ δ : 3.38 (3H, s, NCH₃), 6.46 (1H, d, $J=1.8$, 3-H), 6.43 (1H, dd, $J=7.3$ and 1.8 , 5-H), 7.60 (1H, d, $J=7.3$, 6-H), 3.73 (3H, s, 2'-OCH₃), 6.49 (1H, d, $J=1.8$, 3'-H), 9.85 (1H, br, 4'-OH), 6.48 (1H, dd, $J=7.9$ and 1.8 , 5'-H), 7.11 (1H, d, $J=7.9$, 6'-H).

Photoreaction of 1 with 2d. Product (**3d**) (38 mg, 27%) was obtained. 4-(2-Hydroxy-5-methoxyphenyl)-1-methyl-2-pyridone (**3d**); colorless crystals from EtOH-C₆H₆, mp 237-239 °C (Anal. Calcd for C₁₃H₁₃NO₃: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.25; H, 5.61; N, 5.96); MS m/z : 231 (M^+ , 100%), 230, 214; IR (KBr): 3000(OH), 1645(C=O); $^1\text{H-NMR}$ δ : 3.58 (3H, s, NCH₃), 6.82 (1H, d, $J=1.8$, 3-H), 6.42 (1H, dd, $J=7.3$ and 1.8 , 5-H), 7.35 (1H, d, $J=7.3$, 6-H), 10.45 (1H, br, 2'-OH), 6.93 (1H, d, $J=8.3$, 3'-H), 6.85 (1H, dd, $J=8.3$ and 2.9 , 4'-H), 3.78 (3H, s, 5'-OCH₃), 7.30 (1H, d, $J=2.9$, 6'-H). Irradiation of the singlet at δ 3.78 (5'-OCH₃) caused enhancements of the double doublet at δ 6.85 (4'-H) and the doublet at δ 7.30 (6'-H).

Photoreaction of 1 with 2e. An inseparable 1:1 mixture (14 mg, 10%) of **3e** and **4e** was obtained. 4-(3-Cyano-2-hydroxyphenyl)-1-methyl-2-pyridone (**3e**); GLC-MS m/z : 226 (M^+ , 100%), 225, 209; $^1\text{H-NMR}$ δ : 3.47 (3H, s, NCH₃), 6.47 (1H, d, $J=1.8$, 3-H), 6.32 (1H, dd, $J=6.7$ and 1.8 , 5-H), 7.75 (1H, d, $J=6.7$, 6-H), 10.45 (1H, br, 2'-OH), 7.68 (1H, dd, $J=7.3$ and 1.8 , 4'-H), 7.08 (1H, dd, $J=7.3$ and 7.3 , 5'-H), 7.56 (1H, dd, $J=7.3$ and 1.8 , 6'-H). 4-(3-Cyano-4-hydroxyphenyl)-1-methyl-2-pyridone (**4e**); GLC-MS m/z : 226 (M^+ , 100%), 225, 209; $^1\text{H-NMR}$ δ : 3.53 (3H, s, NCH₃), 6.67 (1H, d, $J=1.8$, 3-H), 6.58 (1H, dd, $J=6.7$ and 1.8 , 5-H), 7.77 (1H, d, $J=6.7$, 6-H), 8.03 (1H, d, $J=1.8$, 2'-H), 10.78 (1H, br, 4'-OH), 7.08 (1H, d, $J=7.3$, 5'-H), 7.89 (1H, dd, $J=7.3$ and 1.8 , 6'-H).

Photoreaction of 1 with 2f. An inseparable 5:3 mixture (43 mg, 32%) of **3f** and **4f** was obtained. 4-(4-Cyano-2-hydroxyphenyl)-1-methyl-2-pyridone (**3f**); GLC-MS m/z : 226 (M^+ , 100%), 225, 209; $^1\text{H-NMR}$ δ : 3.40 (3H, s, NCH₃), 6.40 (1H, d, $J=1.8$, 3-H), 6.22 (1H, dd, $J=7.3$ and 1.8 , 5-H), 7.70 (1H, d, $J=7.3$, 6-H), 10.30 (1H, br, 2'-OH), 7.25 (1H, d, $J=1.8$, 3'-H), 7.34 (1H, dd, $J=7.9$ and 1.8 , 5'-H), 7.38 (1H, d, $J=7.9$, 6'-H). 4-(2-Cyano-4-hydroxyphenyl)-1-methyl-2-pyridone (**4f**); GLC-MS m/z : 226 (M^+ , 100%), 226, 209; $^1\text{H-NMR}$ δ : 3.38 (3H, s, NCH₃), 6.57 (1H, d, $J=1.8$, 3-H), 6.42 (1H, dd, $J=7.3$ and 1.8 , 5-H), 7.68 (1H, d, $J=7.3$, 6-H), 7.35 (1H, d, $J=1.8$, 3'-H), 10.70 (1H, br, 4'-OH), 7.23 (1H, dd, $J=7.9$ and 1.8 , 5'-H), 7.48 (1H, d, $J=7.9$, 6'-H). The mixture was treated with an ethereal solution of CH₂N₂ to give an 1:1 mixture of the methyl ethers of **3f** and **4f**, which was also inseparable. 4-(4-Cyano-

2-methoxyphenyl)-1-methyl-2-pyridone; $^1\text{H-NMR}$ (CDCl_3) δ : 3.57 (3H, s, NCH_3), 6.65 (1H, d, $J = 1.8$, 3-H), 6.42 (1H, dd, $J = 6.7$ and 1.8 , 5-H), 7.23 (1H, d, $J = 6.7$, 6-H), 3.88 (3H, s, $2'\text{-OCH}_3$), 7.23 (1H, d, $J = 2.4$, $3'\text{-H}$), 7.36 (1H, dd, $J = 7.3$ and 2.4 , $5'\text{-H}$), 7.43 (1H, d, $J = 7.3$, $6'\text{-H}$). Irradiation of the singlet due to the OCH_3 group at δ 3.88 induced enhancement of the doublet at δ 7.23($3'\text{-H}$). 4-(2-Cyano-4-methoxyphenyl)-1-methyl-2-pyridone; $^1\text{H-NMR}$ (CDCl_3) δ : 3.55 (3H, s, NCH_3), 6.72 (1H, d, $J = 1.8$, 3-H), 6.38 (1H, dd, $J = 6.7$ and 1.8 , 5-H), 7.28 (1H, d, $J = 6.7$, 6-H), 7.19 (1H, d, $J = 2.4$, $3'\text{-H}$), 3.86 (3H, s, $4'\text{-OCH}_3$), 7.17 (1H, dd, $J = 7.3$ and 2.4 , $5'\text{-H}$), 7.41 (1H, d, $J = 7.3$, $6'\text{-H}$). Irradiation of the singlet at δ 3.86 ($4'\text{-OCH}_3$) induced enhancement of the doublet at δ 7.19 ($3'\text{-H}$) and the double doublet at δ 7.17 ($5'\text{-H}$).

Photoreaction of 1 with 2g. Product (**3g**) (24 mg, 18%) was obtained. 4-(5-Cyano-2-hydroxyphenyl)-1-methyl-2-pyridone (**3g**); colorless crystals from $\text{EtOH-C}_6\text{H}_6$, mp $>300^\circ\text{C}$; (M^+ , m/z 226.0743. $\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_2$ requires 226.0743); MS m/z : 226 (100 %), 225, 209; IR (KBr): 3000, 2215(CN), 1650(C=O); $^1\text{H-NMR}$ δ : 3.33 (3H, s, NCH_3), 6.58 (1H, d, $J = 1.8$, 3-H), 6.43 (1H, dd, $J = 8.5$ and 1.8 , 5-H), 7.69 (1H, d, $J = 8.5$, 6-H), 11.20 (1H, br, $2'\text{-OH}$), 7.07 (1H, d, $J = 8.5$, $3'\text{-H}$), 7.68 (1H, dd, $J = 8.5$ and 1.8 , $4'\text{-H}$), 7.78 (1H, d, $J = 1.8$, $6'\text{-H}$).

Photoreaction of 5 with 2a. An inseparable 3:1 mixture (50 mg, 42%) of **6a** and **7a**, and **8a** (12 mg, 10%) was obtained. 6-(2-Hydroxyphenyl)-1-methyl-2-pyridone (**6a**); GLC-MS m/z : 201 (M^+), 200 (100%), 172; $^1\text{H-NMR}$ δ : 3.40 (3H, s, NCH_3), 6.62 (1H, dd, $J = 9.0$ and 1.2 , 3-H), 7.39 (1H, dd, $J = 9.0$ and 6.8 , 4-H), 6.23 (1H, dd, $J = 6.8$ and 1.2 , 5-H), 7.95 (1H, br, $2'\text{-OH}$), 7.05 (1H, d, $J = 8.1$, $3'\text{-H}$), 7.35 (1H, ddd, $J = 8.1$, 7.6 and 1.7 , $4'\text{-H}$), 6.97 (1H, dd, $J = 7.6$ and 7.6 , $5'\text{-H}$), 7.16 (1H, dd, $J = 7.6$ and 1.7 , $6'\text{-H}$). 6-(4-Hydroxyphenyl)-1-methyl-2-pyridone (**7a**); GLC-MS m/z : 201 (M^+), 200 (100%), 172; $^1\text{H-NMR}$ δ : 3.40 (3H, s, NCH_3), 6.58 (1H, dd, $J = 9.0$ and 1.2 , 3-H), 7.39 (1H, dd, $J = 9.0$ and 6.8 , 4-H), 6.13 (1H, dd, $J = 6.8$ and 1.2 , 5-H), 6.95 (2H, d, $J = 8.6$, $2'\text{-H}$ and $6'\text{-H}$), 7.20 (2H, d, $J = 8.6$, $3'\text{-H}$ and $5'\text{-H}$), 7.80 (1H, br, $4'\text{-OH}$). 1-Methyl-6-phenoxy-2-pyridone (**8a**); colorless columns from $n\text{-hexane-C}_6\text{H}_6$, mp $82\text{--}84^\circ\text{C}$ (Anal. Calcd for $\text{C}_{12}\text{H}_{11}\text{NO}_2$: C, 71.62; H, 5.51; N, 6.96. Found: C, 71.83; H, 5.47; N, 6.93); MS m/z : 201 (M^+ , 100%), 108; IR (KBr): 1655(C=O); $^1\text{H-NMR}$ (CDCl_3) δ : 3.63 (3H, s, NCH_3), 6.27 (1H, dd, $J = 9.0$ and 1.2 , 3-H), 7.19 (1H, dd, $J = 9.0$ and 6.8 , 4-H), 5.03 (1H, dd, $J = 6.8$ and 1.2 , 5-H), 7.12 (2H, d, $J = 8.6$, $2'\text{-H}$ and $6'\text{-H}$), 7.44 (2H, dd, $J = 8.6$ and 7.3 , $3'\text{-H}$ and $5'\text{-H}$), 7.29 (1H, dd, $J = 7.3$ and 7.3 , $4'\text{-H}$).

Photoreaction of 5 with 2b. Products (**7b**) (3.5 mg, 3%) and (**8b**) (68 mg, 50%) were obtained. 6-(4-Hydroxy-3-methoxyphenyl)-1-methyl-2-pyridone (**7b**); pale yellow plates from $\text{C}_6\text{H}_6\text{-EtOH}$, mp $194\text{--}196^\circ\text{C}$ (M^+ m/z 231.0893. $\text{C}_{13}\text{H}_{13}\text{NO}_3$ requires 231.0896); MS m/z : 231 (100 %), 230, 214; $^1\text{H-NMR}$ δ : 3.35 (3H, s, NCH_3), 6.58 (1H, dd, $J = 9.2$ and 1.8 , 3-H), 7.37 (1H, dd, $J = 9.2$ and 6.7 , 4-H), 6.10 (1H, dd, $J = 6.7$ and 1.8 , 5-H), 6.80 (1H, d, $J = 1.8$, $2'\text{-H}$), 3.92 (3H, s, $3'\text{-OCH}_3$), 5.86 (1H, br, $4'\text{-OH}$), 6.99 (1H, d, $J = 7.9$, $5'\text{-H}$), 6.85 (1H, dd, $J = 7.9$ and 1.8 , $6'\text{-H}$). Irradiation of the singlet at δ 3.92 ($3'\text{-OCH}_3$)

induced enhancement of the doublet at δ 6.80. **1-Methyl-6-(2-methoxyphenoxy)-2-pyridone (8b)**: colorless columns from hexanes- C_6H_6 , mp 99-101 °C (Anal. Calcd for $C_{13}H_{13}NO_3$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.46; H, 5.63; N, 6.01); m/z 231 (M^+), 108 (100%); IR (KBr): 1660($C=O$); 1H -NMR ($CDCl_3$) δ : 3.19 (3H, s, NCH_3), 6.24 (1H, dd, $J=9.2$ and 1.2, 3-H), 7.20 (1H, dd, $J=9.2$ and 8.0, 4-H), 5.13 (1H, dd, $J=8.0$ and 1.2, 5-H), 3.67 (3H, s, 2'- OCH_3), 7.26 (1H, dd, $J=7.3$ and 1.8, 3'-H), 7.04 (2H, m, 4'-H and 5'-H), 7.15 (1H, dd, $J=7.3$ and 1.8, 6'-H).

Photoreaction of 5 with 2c. Products (**6c**) (28 mg, 20%) and (**8c**) (28 mg, 20%) were obtained. **6-(2-Hydroxy-4-methoxyphenyl)-1-methyl-2-pyridone (6c)**: colorless needles from C_6H_6 -EtOH, mp 250-252 °C (Anal. Calcd for $C_{13}H_{13}NO_3$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.50; H, 5.66; N, 6.01); MS m/z : 231 (M^+ , 100%), 230, 214; IR (KBr): 2950(OH), 1640($C=O$); 1H -NMR δ : 3.25 (3H, s, NCH_3), 6.57 (1H, dd, $J=9.2$ and 1.8, 3-H), 7.37 (1H, dd, $J=9.2$ and 6.7, 4-H), 6.14 (1H, dd, $J=6.7$ and 1.8, 5-H), 7.98 (1H, br, 2'-OH), 6.64 (1H, d, $J=1.8$, 3'-H), 3.78 (3H, s, 4'- OCH_3), 6.55 (1H, dd, $J=8.6$ and 1.8, 5'-H), 7.02 (1H, d, $J=8.6$, 6'-H). Irradiation of the singlet at δ 3.78 (4'- OCH_3) caused enhancements of the signals at δ 6.64 (3'-H) and 6.55 (5'-H). **6-(3-methoxyphenoxy)-1-methyl-2-pyridone (8c)**: colorless plates from C_6H_6 -*n*-hexane, mp 99 °C (Anal. Calcd for $C_{13}H_{13}NO_3$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.74; H, 5.70; N, 6.06); MS m/z : 231 (M^+), 108 (100); IR (KBr): 1660($C=O$); 1H -NMR ($CDCl_3$) δ : 3.60 (3H, s, NCH_3), 6.29 (1H, dd, $J=9.2$ and 1.2, 3-H), 7.20 (1H, dd, $J=9.2$ and 7.3, 4-H), 5.37 (1H, dd, $J=7.3$ and 1.2, 5-H), 6.65 (1H, dd, $J=2.4$ and 2.4, 2'-H), 3.82 (3H, s, 3'- OCH_3), 6.68 (1H, dd, $J=8.5$ and 2.4, 4'-H), 7.32 (1H, dd, $J=8.5$ and 8.5, 5'-H), 6.82 (1H, dd, $J=8.5$ and 2.4, 6'-H).

Photoreaction of 5 with 2d. Products (**6d**) (30 mg, 22%) and (**8d**) (39 mg, 28%) were obtained. **6-(2-Hydroxy-5-methoxyphenyl)-1-methyl-2-pyridone (6d)**: colorless plates from AcOEt-EtOH, mp 196 °C (Anal. Calcd for $C_{13}H_{13}NO_3$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.54; H, 5.67; N, 6.01); MS m/z : 231 (M^+ , 100%), 230, 214; IR (KBr): 3080(OH), 1640($C=O$); 1H -NMR δ : 3.41 (3H, s, NCH_3), 6.58 (1H, dd, $J=8.3$ and 1.2, 3-H), 7.37 (1H, dd, $J=8.3$ and 6.7, 4-H), 6.23 (1H, dd, $J=6.7$ and 1.2, 5-H), 8.95 (1H, br, 2'-OH), 7.02 (1H, d, $J=8.6$, 3'-H), 6.88 (1H, dd, $J=8.6$ and 3.1, 4'-H), 3.81 (3H, s, 5'- OCH_3), 6.68 (1H, d, $J=3.1$, 6'-H). Irradiation of the singlet at δ 3.81 (5'- OCH_3) induced enhancements of the signals at δ 6.88 (4'-H) and 6.68 (6'-H). **6-(4-Methoxyphenoxy)-1-methyl-2-pyridone (8d)**: colorless needles from C_6H_6 -*n*-hexane, mp 132-134 °C (Anal. Calcd for $C_{13}H_{13}NO_3$: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.70; H, 5.68; N, 5.94); MS m/z : 231 (M^+), 108 (100%); IR (KBr): 1655($C=O$); 1H -NMR ($CDCl_3$) δ : 3.63 (3H, s, NCH_3), 6.23 (1H dd, $J=9.2$ and 1.2, 3-H), 7.23 (1H, dd, $J=9.2$ and 7.3, 4-H), 5.23 (1H, dd, $J=7.3$ and 1.2, 5-H), 7.05 (2H, d, $J=9.2$, 2'-H and 6'-H), 6.92 (2H, d, $J=9.2$, 3'-H and 5'-H), 3.83 (3H, s, 4'- OCH_3).

Photoreaction of 5 with 2f. Products (**6f**) (54 mg, 40%) and (**8f**) (34 mg, 25%) were obtained. **6-(2-Cyano-6-hydroxyphenyl)-1-methyl-2-pyridone (6f)**: colorless plates from AcOEt-EtOH, mp 231-233 °C

(Anal. Calcd for $C_{13}H_{10}N_2O_2$: C, 69.01; H, 4.46; N, 12.38. Found: C, 68.72; H, 4.33; N, 12.37); MS m/z : 226 (M^+ , 100%), 225, 209; IR (KBr): 3000(OH), 2220(CN), 1640(C=O); 1H -NMR δ : 3.25 (3H, s, NCH_3), 6.65 (1H, dd, $J=9.2$ and 1.2, 3-H), 7.58 (1H, dd, $J=9.2$ and 6.7, 4-H), 6.38 (1H, dd, $J=6.7$ and 1.2, 5-H), 7.23 (1H, dd, $J=8.6$ and 1.2, 3'-H), 7.51 (1H, dd, $J=8.6$ and 8.6, 4'-H), 7.38 (1H, dd, $J=8.6$ and 1.2, 5'-H), 10.21 (1H, br, 6'-OH). 6-(3-Cyanophenoxy)-1-methyl-2-pyridone (**8f**); colorless plates from C_6H_6 , mp 115-117 °C (Anal. Calcd for $C_{13}H_{10}N_2O_2$: C, 69.01; H, 4.46; N, 12.38. Found: C, 68.86; H, 4.36; N, 12.37); MS m/z : 226 (M^+), 108 (100%); IR (KBr): 1655(C=O); 1H -NMR ($CDCl_3$) δ : 3.53 (3H, s, NCH_3), 6.27 (1H, dd, $J=9.2$ and 1.2, 3-H), 7.16 (1H, dd, $J=9.2$ and 6.7, 4-H), 5.25 (1H, dd, $J=6.7$ and 1.2, 5-H), 7.32 (1H, dd, $J=3.1$ and 2.4, 2'-H), 7.53 (1H, dd, $J=7.5$ and 2.4, 4'-H), 7.35 (1H, m, 5'-H), 7.54 (1H, dd, $J=8.5$ and 3.1, 6'-H).

Photoreaction of 5 with 2g. Product (**6g**) (74 mg, 54%) was obtained. 6-(5-Cyano-2-hydroxyphenyl)-1-methyl-2-pyridone (**6g**); colorless needles from EtOH, mp 280-282 °C (Anal. Calcd for $C_{13}H_{10}N_2O_2$: C, 69.01; H, 4.46; N, 12.38. Found: C, 68.77; H, 4.30; N, 12.41); MS m/z : 226 (M^+ , 100%), 225, 209; IR (KBr): 3000(OH), 2210(CN), 1640(C=O); 1H -NMR δ : 3.32 (3H, s, NCH_3), 6.42 (1H, dd, $J=9.2$ and 1.2, 3-H), 7.24 (1H, dd, $J=9.2$ and 6.7, 4-H), 6.11 (1H, dd, $J=6.7$ and 1.2, 5-H), 11.34 (1H, br, 2'-OH), 7.10 (1H, d, $J=8.6$, 3'-H), 7.79 (1H, dd, $J=8.6$ and 2.4, 4'-H), 7.72 (1H, d, $J=2.4$, 6'-H).

REFERENCES

1. a) K. Konno, K. Hashimoto, Y. Ohfuné, H. Shirahama, and T. Matsumoto, *J. Am. Chem. Soc.*, 1988, **110**, 4807; b) C. A. Veale, P. R. Bernstein, C. Bryant, C. Ceccarelli, J. R. Damewood, Jr., R. Early, S. W. Feeney, B. Gomes, B. J. Kosmider, G. B. Steelman, R. M. Thomas, E. P. Vacek, J. C. Williams, D. J. Wolanin, and S. Woolson, *J. Med. Chem.*, 1995, **38**, 98; c) G. Pattenden and G. F. Smith, *Tetrahedron Lett.*, 1990, **31**, 6557; d) S. Ohmiya, K. Saito, and I. Murakoshi, 'The Alkaloids: Lupine Alkaloids' Vol. 47, ed. by G. A. Cordell, Academic Press, Inc., California, 1995, pp. 1-114.
2. a) C. A. Mirkin, K.-L. Lu, T. F. Snead, G. L. Geoffroy, and A. L. Rheingold, *J. Am. Chem. Soc.*, 1990, **112**, 2809 and references cited therein; b) D. P. Curran and H. Liu, *J. Am. Chem. Soc.*, 1992, **114**, 5863.
3. S. Ohmiya, M. Noguchi, C.-Y. Chen, I. Murakoshi, and H. Otomasu, *Chem. Pharm. Bull.*, 1989, **37**, 2516.
4. S. Ohmiya, M. Noguchi, S. Ina, H. Kubo, and H. Otomasu, *Chem. Pharm. Bull.*, 1992, **40**, 854; N. Sakurai and S. Ohmiya, *J. Chem. Soc., Chem. Commun.*, **1993**, 297.
5. I. Sircar, R. P. Steffen, G. Bobowski, S. E. Burke, R. S. Newton, R. E. Weishaar, J. A. Bristol, and D. B. Evans, *J. Med. Chem.*, 1989, **32**, 342.
6. K. Kubo, N. Ito, Y. Isomura, I. Sozu, H. Arima, M. Homma, and M. Murakami, *Yakugaku Zasshi*, 1979, **99**, 588.
7. R. Beugelmans and M. Bois-Choussy, *Tetrahedron Lett.*, 1988, **29**, 1289; A. B. Pierini, M. T. Baumgartner, and R. A. Rossi, *Tetrahedron Lett.*, 1988, **29**, 3429.
8. N. Alam, C. Amatore, C. Combéllas, A. Thiebault, and J. N. Verpeaux, *Tetrahedron Lett.*, 1987, **28**, 6171.
9. R. Graf, *Ber.*, 1931, **64**, 21.
10. A. R. Katritzky, J. D. Rowe, and S. K. Roy, *J. Chem. Soc. B*, **1967**, 758.
11. B. J. Van Keulen, R. M. Kellog, and O. Piepers, *J. Chem. Soc., Chem. Commun.*, **1979**, 285.