


SENSITIZED PHOTOOXIDATION
OF N-METHYLPYRIDINIUM-3-OXIDE AND ITS RELATED COMPOUNDS

Akira Mori, Shumei Ohta, and Hitoshi Takeshita


Research Institute of Industrial Science, Kyushu University,

Hakozaki, Fukuoka 812


In connection with our previous observation which indicated the involvement of a peroxirane intermediate, a dipolar species, in the sensitized photooxidation of a spirocyclic vinylcyclopropane derivative, we have now carried out the oxidation with N-methylpyridinium-3-oxide (I), a betaine, for the first time. The results, which show that there is a marked solvent effect on the product formation, are presented together with a discussion of the mechanism of their formation. When a chloroform solution of (I) was irradiated by means of a tungsten lamp under an oxygen atmosphere, 5-(N-methylformamido)-2(5H)furanone (II) was obtained along with N-methylmaleimide (III). On the other hand, the oxidation in an ethanol solution of (I) afforded diethyl maleate (IV), diethyl fumarate (V), N-methyl-3-hydroxy- α -pyridone (VI) and ethyl N-methyloxamate (VII). The formation of (II) was explained by lactonization after cleavage of the dioxetane intermediate; while (III) was produced by a further oxidation via the five-membered peroxide. In contrast to this, the reaction in a protic solvent is more complicated; none of the products correspond to an oxidation product formed with one molar unit of singlet oxygen, *i.e.*, (VI) has a lower oxidation state than the normal product, while the others formed by a multistep oxidation. The formation of (VI) can be interpreted in terms of intermolecular intervention of a dioxetane precursor, peroxirane, by reduction with solvent. A tentative pathway (path b) leading to (IV), (V) and (VII) has been proposed after excluding an alternative route by a further irradiation of (VI) which failed to give any isolable product.




(I)



(III)



(II)



(VI)

