

SYNTHESIS OF A BRIDGED HOMO(15)ANNULENIUM ION

Haru Ogawa and Iwao Tabushi

Faculty of Pharmaceutical Sciences, Kyushu University,


Katakasu, Fukuoka, 812

Hidefumi Kato and Yōichi Taniguchi


Department of General Chemistry, Kurume Technical College,

Komorino, Kurume


Ring expansion of 2,15-dimethoxycarbonyl(15)annulenone 4,7 : 10,13-dioxide I by dimethylsulfonium methylide gave rise to a 16-membered homo(15)annulene II, which was transformed to a diol III by a decarboxylation (copperchromite in quinoline)-reduction (NaBH_4) sequence. Compound III is considered to be a possible precursor of homo(15)annulenium ion, a 14π homoaromatic cation IV. As expected, IV in conc. H_2SO_4 at room temperature exhibited a characteristic spectrum, which is very similar to that of aromatic (15)annulenium tetrafluoroborate V (λ_{max} (in 30 % H_2SO_4) at 331 (110,000), 345 nm (10,800)), exhibiting absorption maxima (in conc. H_2SO_4) at 334 and 349 nm (33,000). This may be indicative of the occurrence of IV.




(I)




(II)



(III)



(IV)



(V)