

A NEW SYNTHETIC APPROACH TO THE 1,4,2-OXATHIAZOLE RING SYSTEM BY
THE ACTION OF DIMETHYL ACETYLENEDICARBOXYLATE ON ARYLTHIOHYDROXAMIC ACIDS¹

PARTHASARATHI RAJAGOPALAN*² AND CHANDRAVADAN N. TALATY³

CIBA-GEIGY Research Centre, Bombay 400063, India


The reaction of arylthiohydroxamic
acids with dimethyl acetylenedicarboxylate
yields, in one step, 3-Aryl-5-methoxy-
carbonylmethylene-5-methoxycarbonyl-
1,4,2-oxathiazoles.

Acetylenedicarboxylic acid esters play a unique and important role in
synthetic organic chemistry in that they can undergo, besides simple nucleophilic
additions, cyclizations following nucleophilic additions as well as 1,2-, 1,3-,
1,4-cycloadditions leading to many an interesting carbocyclic and heterocyclic
system.⁴


Over nine years ago we had occasion to examine the reaction of arylthio-
hydroxamic acids (I) with dimethyl acetylenedicarboxylate (II) and would now
like to report our findings.

When methanolic solutions of equimolar quantities of 4-chlorophenylthio-
hydroxamic acid (I, Ar = 4-chlorophenyl)⁵ and II were mixed, set aside at room
temperature for two days, filtered to remove traces of sulfur, the filtrate
stripped of the solvent under reduced pressure, the residual viscous liquid
extracted repeatedly with hot pentane, and the pentane extracts cooled in dry


ice-acetone bath, a solid was obtained which on recrystallization from hexane furnished a crystalline product, m.p. 88-90°, in 25% yield. The molecular formula ($C_{13}H_{12}ClNO_5S$) of the product indicated that it was a 1:1 adduct of I (Ar = 4-chlorophenyl) and II and that, consequently, it could possess any one of the structures represented by IIIa, IV, V⁶ or VI.



I



II




III


III

- | | |
|----------------------------|-----|
| a. Ar = 4-chlorophenyl | 25% |
| b. Ar = 4-fluorophenyl | 40% |
| c. Ar = 4-nitrophenyl | 15% |
| d. Ar = 3,4-dichlorophenyl | 30% |


yield



IV



V



VI

The infrared spectrum of the adduct showed no absorption in the $>\text{NH}$ or $-\text{OH}$ region and a strong and broad carbonyl absorption at 1740 cm^{-1} . Its 100 MHz PMR spectrum carried, besides two singlets (3 protons each) at $\delta 3.73$ and $\delta 3.84$ corresponding, respectively, to the two methyl groups of the two ester functions and an AA'BB' multiplet (4 protons) centered at $\delta 8.02$ corresponding to the 4-chlorophenyl group, a highly significant AB quartet (2 protons, $J = 16\text{ Hz}$) centered at $\delta 3.46$ and no protons that could be exchanged with D_2O even after 15 hours.

Structure IV could be ruled out immediately for, if it were correct, it would give rise to an absorption due the hydroxyl group of the oxime function in its infrared spectrum and its PMR spectrum should exhibit a singlet due to the vinyl proton and a singlet due to the hydroxyl proton which should disappear on exposure to D_2O .

If the adduct possessed either of the structures V or VI, its PMR spectrum should display an AB quartet for the vicinal protons designated A and B respectively, with a coupling constant not exceeding 12Hz at the very extreme irrespective of whether their orientation is cis or trans with respect to each other. The value of the coupling constant of vicinal protons can be derived, within certain limitations, from the Karplus equation⁷ which has been applied effectively by Huisgen⁸ to evaluate the stereochemistry of 1,3-dipolar cycloadditions and which has since been modified by Pachler.⁹ However, structures V (dihedral angle $\Theta = 10^\circ$ for cis and 130° for trans) and VI ($\Theta = 40^\circ$ for cis and 160° for trans) can be excluded on the basis of the observed value for the coupling constant (i.e. 16Hz) which is much larger than that could be expected for the spin-spin coupling of the vicinal protons of either of them.

In fact the observed coupling pattern with the coupling constant of 16Hz

is consistent only with the structure IIIa for the adduct which with its diastereotopic methylene protons would, in its NMR spectrum, be expected to exhibit an AB quartet with a J value of 16-18 Hz.

Compounds IIIb, IIIc, and IIId were also prepared by this one step procedure which constitutes a new route to the 1,4,2-oxathiazole ring system.

Acknowledgement

We thank Dr. S. Selvavinayakam and his associates for the analytical and spectral data, and Dr. R. Zürcher for the 100 Hz PMR spectrum of compound IIIa.

References

1. Dedicated to Dr. T.R. Govindachari, Director, CIBA-GEIGY Research Centre, on his 60th birthday.
2. Present address: Endo Laboratories, Inc., Garden City, NY 11530, U.S.A.
3. Present address: Department of Pharmaceutical Chemistry, University of Maryland, Baltimore, MD, U.S.A.
4. R. Fuks and H.G. Viehe, 'Chemistry of Acetylenes', edited by H.G. Viehe Marcel Dekker, New York, 1969, p. 425.
5. T. Bacchetti and A. Alemagna, Atti acad. nazl. Lincei. Rend., Classe sci. fis. mat. e nat. 1958, 24, 161.. Chem. Abstr., 1958, 52, 18299.
6. We thank Dr. Nagarajan for suggesting this structure as a possibility.
7. M. Karplus, J. Chem. Physics, 1959, 30, 11. J. Amer. Chem. Soc., 1963, 85, 2870.
8. G. Binsch, R. Huisgen, and H. König, Chem. Ber., 1964, 97, 2893.
9. K.G.R. Pachler, J. Chem. Soc. Perkin II, 1972, 1936.

Received, 9th June, 1975