

BIOMIMETIC TRANSFORMATION OF INDOLOQUINOLIZIDINE DERIVATIVE TO 2-ACYLINDOLE

Masako Nakagawa^{*}, Yoko Okajima, Kazumi Kobayashi,

Toshifumi Asaka, and Tohru Hino





Faculty of Pharmaceutical Sciences, Chiba University, Yayoi-cho, Chiba-shi, 280

2-Acylindole 2 has been synthesised by photosensitised oxygenation of indoloquinolizidine 1a, while 1-methyl derivative 1b gave a quinolone 3.

We have previously reported the model reaction for biological oxidation of tryptophan catalyzed by monooxygenase¹ and dioxygenase² to form 3a-hydroxy-pyrroloindoles and oxindoles.

The present paper reports a facile one step synthesis of 2-acylindole 2 from indoloquinolizidine 1³, which provides an implication for a biogenetic formation of 2-acylindole alkaloids from the corresponding indole alkaloids.

A solution of quinolizidinium chloride 1a in water was continuously saturated with oxygen and irradiated with 250 W halogen lamp through a pyrex filter for 20 hr in the presence of rose bengal. Chromatography and elution with CH₂Cl₂ gave 2, mp 134-136°⁴, in 10% yield, whose structure was identified by comparison with an authentic sample. Bubbling oxygen through a solution of 1a in the dark or irradiation of 1a in the absence of the sensitizer yielded no detectable 2-acylindole and the



starting material was recovered. However, if 1a was irradiated in MeOH or 50% aqueous MeOH, 2 was not obtained.

The conversion of simple 2,3-dialkylindoles to 2-acylindoles by autoxidation has many precedents⁵ and the hydroperoxyindolenine has been considered as the primary intermediate. Although the mechanism of formation of 2 is not entirely clear, singlet oxygen is known to add to an enamine system to give an ene reaction product⁶. Thereby, it is conceivable that the reaction from the hydroperoxide to 2 was processed as autoxidation, e.g., the primary intermediate 5 isomerized to the enamine tautomer 6 which was converted into the final product 2.

Support for this idea was given by the observation that photosensitized oxygenation of 1b in H₂O under similar conditions resulted in the formation of 1,2,3,4,6,7,12,12b-octahydroindolizino[1,2-b]quinoline-7-one methochloride 4 in 25% yield which has been derived from cyclization of a cleavage product 3⁷ and the failure to detect 2-acylindole derivative. The structure of 4 was confirmed by direct comparison with an authentic specimen prepared by ozonization of 1b in MeOH. 4; mp 181°(dec.), $\lambda_{\text{max}}^{\text{EtOH}}$ 240, 319, 332 nm⁸, $\gamma_{\text{max}}^{\text{KBr}}$ 1610 cm⁻¹ (C=O)⁸, δ (d₆-DMSO) 3.10-4.0 (2H, m, 4-CH₂), 3.33(3H, s, 5-N-Me), 3.79(3H, s, 12-N-Me), 4.76(2H, AB quartet, 6-CH₂), 5.20(1H, m, 12b-CH), picrate, mp 295-297°(aqueous MeOH)⁹.

Although chemical transformation of indole alkaloids to the corresponding 2-acylindole alkaloids has been reported¹⁰, our preliminary results appear to be a plausible biogenetic route.

We thank Professor L.J.Dolby, University of Oregon, for gift of 2-acylindole 2. We also thank the Ministry of Education for financial support.

REFERENCES

1. M.Nakagawa, T.Kaneko, H.Yamaguchi, T.Kawashima, and T.Hino, Tetrahedron, 1974, 30, 2591.
2. M.Nakagawa, T.Kaneko, K.Yoshikawa, and T.Hino, J.Amer.Chem.Soc., 1974, 96, 624.
3. M.Nakagawa, M.Kiuchi, M.Obi, M.Tonozuka, K.Kobayashi, T.Hino, and Y.Ban, Chem.Pharm.Bull., 1975, 23, 304.
4. L.J.Dolby and G.W.Gribble, J.Org.Chem., 1967, 32, 1391.
5. E.Leete, J.Amer.Chem.Soc., 1961, 83, 3645; H.H.Wasserman and M.B.Floyd, Tetrahedron Lett., 1963, 2009; F.Y.-H. Chen and E.Leete, ibid., 1963, 2013; W.I.Taylor, Proc.Chem.Soc., 1962, 247; S.McLean and G.I.Dmitrienko, Canad.J.Chem., 1971, 49, 3642.
6. I.Saito, M.Imuta, and T.Matsuura, Chem.Lett., Japan, 1972, 1173, 1197 and references are therein.
7. Photosensitised oxygenation of tertiary enamines has been known to undergo oxidative α,β -cleavage via dioxetane intermediates to give carbonyl and amide fragments. cf. C.S.Foote, A.A.Dzakpasu, and J.W.-P.Lim, Tetrahedron Lett., 1975, 1247.
8. B.Witkop and S.Goodwin, J.Amer.Chem.Soc., 1953, 75, 3371.
9. The new compound 4 analysed correctly for C,H,N.
10. L.J.Dolby and S.Sakai, J.Amer.Chem.Soc., 1964, 86, 1890, 5362;

L.J.Dolby and S.Sakai, Tetrahedron, 1967, 23, 1 ; G.H.Foster, J. Harley-Mason, and W.R.Waterfield, Chem Comm., 1967, 21 ; J.D. Albright and L.Goldmann, J.Amer.Chem.Soc., 1969, 91, 4317 ; S.Sakai, A.Kubo, K.Katano, N.Shinma, and K.Sasago, Yakugaku Zasshi, 1973, 93, 1165 ; S.Sakai, E.Yamanaka, K.Katano, and S.Kimura, Abstracts of the 7th. Congress of Heterocyclic Chemistry, Japan, 1974, p 195. cf. Heterocycles, 1975, 3, 94.

Received, 14th August, 1975