

ADDITION REACTIONS OF ACID ANHYDRIDES TO ISOQUINOLINE
AND ITS RELATED COMPOUNDS

Hiroshi Yamanaka, Takayuki Shiraishi, and Takao Sakamoto

Pharmaceutical Institute, Tohoku University

Aobayama, Sendai, 980

Reactions of quinoline, isoquinoline and their N-oxides with acid anhydrides were investigated.

When quinoline N-oxide was treated with diketene in acetic acid, a product 2,6-dimethyl-3-(2-quinolyl)-4-pyrone (39%) was obtained. On the contrary, isoquinoline N-oxide gave three products in the same reaction condition. The structures of these products have been assigned as 2,6-dimethyl-3-(1-isoquinolyl)-4-pyrone (30%), 1-acetyl-2-acetyl-4-acetoxy-1,2-dihydroisoquinoline (13%) and 4-methyl-2-oxo-2H-benzo[a]quinolizine (10%), respectively.

The second product was also obtained from the reaction of 4-acetoxy-isoquinoline with diketene in acetic acid. This fact might suggest that addition of diketene take place to the C=N double bond of the isoquinoline ring. In fact, isoquinoline was transformed to 1-acetyl-2-acetyl-1,2-dihydroisoquinoline (43%) in a manner similar to that given for isoquinoline N-oxide.

As diketene is considered to be an acid anhydride, isoquinoline was heated in acetic anhydride without addition of any other substance to give 2-acetyl-1,2-dihydroisoquinoline-1-acetic acid (43%) whose spectral data are consistent with the structure. This compound was converted to 1-(2-hydroxyethyl)-2-ethyl-1,2,3,4-tetrahydroisoquinoline which was identical with the authentic specimen.

Although such heteroaromatics containing isoquinoline type C=N double bond as phthalazine and 1,6-naphthyridine also reacted with diketene in acetic acid yielding similar adducts, reactions of quinoline and quinoxaline with diketene or acetic anhydride were resulted in the recovery of the starting materials.