

STUDIES ON THE SYNTHESIS OF THE DIMERIC INDOLE ALKALOIDS
THE STEREOSPECIFIC SYNTHESIS OF (±)-DIHYDROCLEAVAMINE AND A NEW
SYNTHESIS OF (±)-QUEBRACHAMINE AND ITS (±)-14, 15-DEHYDRODERIVATIVE

Seiichi Takano, Susumi Hatakeyama, Michiyasu Hirama, Tsutomu Araki, Shin-ichi Yamada,

Masaaki Sato, Tsutomu Sugahara, Kozo Shishido, and Kunio Ogasawara

Pharmaceutical Institute, Tohoku University,

Aobayama, Sendai 980

Novel syntheses of the entitled compounds and their related compounds *via* sulfur containing intermediates have been reported.

Hydroboration followed by oxidation of an epimeric mixture of 2-(2-ethylallyl)-3-thioxo-1,2,3,5,6,11b-hexahydro-11H-pyrrolo[1,2-a]- β -carboline afforded the primary alcohol; 2-(2-ethyl-3-hydroxypropyl)-3-thioxo-1,2,3,5,6,11b-hexahydro-11H-pyrrolo[1,2-a]- β -carboline as a single stereoisomer which was then led to (+)-dihydrocleavamine.

A new synthesis of (+)-quebrachamine was accomplished by two different ways, in which the one employed the thio-Claisen rearrangement to construct crucial intermediate, while the other used the α -diketomonothioketal cleavage reaction and in the latter a introduction of 14,15-double bond of quebrachamine was accomplished by the thermolysis of the corresponding sulfoxide.