

PREPARATION OF N-AMINOHETEROCYCLES AND THEIR REACTIONS

Masanori Somei, Mari Matsubara, and Mitsutaka Natsume

Research Foundation ITSUU Laboratory

Tamagawa 2-28-10, Setagaya, Tokyo

A novel method for the synthesis of N-aminoheterocycles is reported.

- 1) Various reaction conditions were investigated and optimum condition was established. By this method, indoles such as indole, skatole, 2-methylindole, 3-phenylindole, 3-phenylthioindole, 3-isoprenylindole, and indole-acetonitrile were converted to the corresponding 1-aminoindole derivatives.
- 2) The reactivity of 1-aminoindoles were widely investigated and it was shown that they reacted with
 - i) acyl chlorides and sulfonyl chlorides
 - ii) carboxylic acids
 - iii) isocyanates
 - iv) pyrrylium salts
 - v) 1,3-diketones and 1,4-diketones
 - vi) aldehydes, ketones, and α,β -unsaturated ketones
 - vii) epoxide
 - viii) ribose
- 3) Novel pyridazino(2,3-a)indoles and as-triazino(1,6-a)indoles were prepared and some reactions of them were reported.
- 4) Our N-amination method was successfully applied for 1,2,3-benzotriazole and benzimidazole. Interestingly, 6-benzylaminopurine and adenine were aminated and their structures were proved to be 9-aminopurine derivatives. Attempts to prepare nucleoside analogues are now in progress.