

SYNTHESES AND REACTIONS OF 2-INDOL-3-YL-1,3-OXATHIOLIUM SALTS¹⁾

Yoshinori Tominaga, Yoshiro Matsuda, and Goro Kobayashi*

Faculty of Pharmaceutical Sciences, Nagasaki University

1-14 Bunkyo-machi, Nagasaki, Japan 852


The reaction of methyl indole-3-dithiocarboxylates (1a,b,c,d) with phenacyl bromide in acetone gave 2-indol-3-yl-1,3-oxathiolium bromides (2a,b,c,d) which reacted with active methylene compounds to form 2-indol-3-ylthiophene derivatives (3a,b,c,d)

We had previously reported that 3-(α,α -bismethylthiomethylene)-indolenium methyl sulfates, which were prepared from methyl indole-3-dithiocarboxylates, reacted with active methylene compounds to form 3-(methylthio)vinylindole derivatives in good yields²⁾. In our present communication, we report the syntheses and reactions of 2-indol-3-yl-1,3-oxathiolium salts.

These 2-indol-3-yl-1,3-oxathiolium salts were prepared by the following manner: A solution of methyl indole-3-dithiocarboxylates (1a,b,c) and phenacyl bromide in absolute acetone was refluxed for 8 hr on a boiling water bath and then the solvent was evaporated to one-third volume. The concentrated

solution was allowed to stand at room temperature for 4 hr and the precipitated yellow needles were collected on a filter and recrystallized from a mixture of methanol and acetone to give 2-indol-3-yl-1,3-oxathiolium bromide (2a,b,c,d) in 40 - 60% yields. The compounds 2a and 2b were also obtained by reaction of thioamides(1e,f,g) with phenacyl bromide using Hartman's method³⁻⁵).


Although some syntheses of 1,3-oxathiolium salts from thioamides or thiolesters and a few syntheses of 1,3-dithiolium salts from dithiocarboxylates have been reported^{7,8}), there has been to date no reported synthesis of 1,3-oxathiolium salts from dithiocarboxylates.



1a,b,c,d

a; $R^1=H$, $R^2=Me$, b; $R^1=R^2=Me$

c; $R^1=H$, $R^2=Ph$, d; $R^1=Me$, $R^2=Ph$



1e,f,g

e; $R^1=H$, $NR_2=N$ (pyrrolidine ring)

f; $R^1=Me$, $NR_2=N$ (pyrrolidine ring)

g; $R^1=H$, $NR_2=N$ (piperidine ring)


2a,b,c,d

2	R^1	R^2	mp(°C)	UV λ_{max}^{EtOH}	ϵ_{nm} (log ϵ)
a	H	Me	284	280	(4.49)
b	Me	Me	284	285	(4.45)
c	H	Ph	253	247	(4.51), 287(4.32)
d	Me	Ph	274	246	(4.53), 300(4.17)

Reaction of 2b with malononitrile in the presence of powdered potassium hydroxide in dimethyl sulfoxide gave in 72% yield yellow needles of a compound mp 220°, which was shown to be 4-amino-5-benzoyl-3-cyano-2-(1,2-dimethylindol-3-yl)-thiophene from spectral data and elemental analysis.

Similarly, reaction of 2d with active methylene compounds-malononitrile, ethyl acetoacetate, and acetylacetone afforded 5-benzoyl-2-indol-3-ylthiophene derivatives (3b,c,d) in 50 -80% yield.

Recently, Hirai and Ishiba reported the reaction of 1-aryl-1,3-oxathiolium salts with active methylene to form thiophene derivatives⁶⁾. Our results fall into the same category.



3	R ¹	R ²	R ³	R ⁴	mp (°C)	IR (KBr) cm ⁻¹	UV _{max} ^{EtOH} nm (log ε)
a	Me	Me	CN	NH ₂	220	3400 (NH) 3290 (NH) 2210 (CN) 1610 (C=O)	270 (4.14) 352 (4.03) 400 (4.32)
b	Me	Ph	CN	NH ₂	218	3280 (NH) 3300 (NH) 2200 (CN) 1590 (C=O)	294 (4.23) 400 (4.24)
c	Me	Ph	COOEt	Me	154	1700 (C=O) 1620 (C=O)	290 (4.32) 380 (3.96)
d	Me	Ph	Ac	Me	167	1675 (C=O) 1628 (C=O)	290 (4.35) 380 (3.97)

REFERENCES

- 1) This report is "Studies on Indole Derivatives XXVII".
- 2) Y.Tominaga, Y.Matsuda, and G.Kobayashi, J. Pharm. Soc. Japan, 1975, 95, 1073.
- 3) H.Hartman, J. Prakt. Chem., 1971, 313, 730.
- 4) H.Hartman, Z. Chem., 1971, 11, 421.
- 5) H.Hartman, H.Schafer, and K.Gewald, J. Prakt. Chem., 1973, 315, 497.
- 6) K.Hirai and T.Ishiba, Heterocycles, 1975, 3, 217.
- 7) E.Campaigne and N.W.Jacobsen, J. Org. Chem., 1964, 29, 1703.
- 8) M.Ohta and M.Sugiyama, Bull. Chem. Soc. Japan, 1963, 36, 1437.

Received, 8th September, 1975