

SCIADENINE, A NEW BISBENZYLISOQUINOLINE ALKALOID
FROM SCIADOTENIA TOXIFERA

Keiichi Takahashi, Michael J. Mitchell and Michael P. Cava*

Department of Chemistry, University of Pennsylvania

Philadelphia, Pennsylvania 19174, U. S. A.

Sciadenine, a new alkaloid from Sciadotenia toxifera
Krukoff and A. C. Smith (Menispermaceae), has been assigned
the head-to-tail bisbenzylisoquinoline structure I.

The genus Sciadotenia (Menispermaceae) has not hitherto been the subject
of any phytochemical investigation. As part of a broad study of the alkaloids
of the South American Menispermaceae, we now report the isolation and structure
determination of the new bisbenzylisoquinoline alkaloid sciadenine (I) from
Sciadotenia toxifera Krukoff and A. C. Smith.


Countercurrent fractionation of the bases from S. toxifera followed by
recrystallization from chloroform-acetone, gave colorless needles of
sciadenine (I), mp 254-256° (decomp.), $[\alpha]_D^{30} = -43^\circ$ (c = 1.22 pyridine),
 $[\alpha]_D^{25} = +15^\circ$ (c = 1.185, CHCl₃), uv $\lambda_{\max}^{\text{EtOH}}$ (log ϵ) 283 nm (3.47), and 277
(3.48). The mass spectrum of I indicates the composition C₃₇H₄₀N₂O₆,
showing a molecular ion peak at m/e 608, and fragment ions at m/e 607, 312
(base peak), 298, 204, 191 and 190. The fragmentation pattern indicates
that I is derived from coclaurine units joined head to tail¹⁾. The nmr
spectrum of I (CDCl₃) showed three methoxyls at δ 3.40 (3 H), 3.80 (3 H),
and 3.82 (3 H), two N-methyls at δ 2.23 (3 H), and 2.47 (3 H) and ten

aromatic protons in the range δ 5.70 ~ 6.85. Treatment of I with diazomethane afforded the optically inactive O-methyl derivative (II), mp 262-263° (decomp.), which showed the molecular ion peak at m/e 622 and a strong cleavage ion at m/e 312 (base peak) in the mass spectrum; four methoxy groups at δ 3.41 (6 H) and 3.83 (6 H) and two N-methyl groups at δ 2.50 (6 H) were observed in its nmr spectrum.


Reductive cleavage of II with sodium in liquid ammonia gave only dl-armepavine (IV), which was identical with an authentic sample²⁾. Treatment of I with diazoethane afforded the O-ethyl derivative (III), mp 233-234° (decomp.), the nmr spectrum of which showed an O-ethyl group at δ 0.83 (3 H, t, J = 7 Hz) and 3.66 (2 H, q, J = 7 Hz), three methoxy groups at δ 3.41 (3 H) and 3.80 (6 H), and two N-methyl groups at δ 2.47 (6 H). The mass spectrum showed the expected molecular ion peak at m/e 636 and major fragment ions at m/e 326, 312, 204 and 190.

Reductive cleavage of III with sodium in liquid ammonia gave a mixture of R-armepavine (V) and S-7-ethoxy-1,2,3,4-tetrahydro-1-(4-hydroxybenzyl)-6-methoxy-2-methylisoquinoline (VI), which could not be separated by tlc or recrystallization. Treatment of the above mixture with l-(-)-mandelic acid gave colorless needles, mp 172-174° (EtOH), $[\alpha]_D^{25} = +48^\circ$ (c = 0.9, MeOH). Basification of an aqueous solution of the above salt afforded VI, mp 129-130° (acetone-Et₂O-hexane) [Ref.³⁾ mp 130-131°], $[\alpha]_D^{30} = +71^\circ$ (c = 0.62, CHCl₃) [Ref.³⁾ $[\alpha]_D^{17} = +83^\circ$ (c = 0.72, CHCl₃)]. The nmr spectrum showed the O-ethyl group at δ 1.32 (3 H, t, J = 7 Hz), 3.77 (2 H, q, J = 7 Hz), N-methyl group at δ 2.51 (3 H), methoxy group at δ 3.82 (3 H), C₈-proton at δ 6.08, C₅-proton at δ 6.56, C₃' and C₅'-protons at 6.60 (d, J = 8 Hz), and C₂' and C₆'-protons at δ 6.90 (d, J = 8 Hz). The above nmr values indicate


that VI is a 7-ethoxy-6-methoxy substituted isoquinoline derivative⁴⁾. The mother liquor was treated with d-(+)-mandelic acid in the usual way to give a salt which afforded crystalline R-armepavine (V), which was identical with an authentic sample.⁵⁾



- (I) R = H
 (II) R = Me
 (III) R = Et



- (IV) R =  H
 (V) R =  H



(VI)

ACKNOWLEDGEMENT We thank the National Institutes of Health for a grant (CA-11445) in support of this work. We also thank B. A. Krukoff (Honorary Curator, New York Botanical Garden and Consulting Botanist of Merck, Sharp and Dohme Research Laboratories) for the supply of plant material, identification of the voucher sample and botanical consultations. The plant material (stem of the lower part of the vine with bark on) used in this investigation was collected by Sr. Jose Schunke in Peru (San Martin: Mariscal Cáceres, Tocáche Nuevo) in 1971. The vouchers have been deposited in New York Botanical Garden, Field Museum of Natural History, Chicago, and many other institutions. This collection was cited in *Phytologia* 25:35. 1972.

REFERENCES

- 1 J. Baldas, I. R. C. Bick, T. Ibuka, R. S. Kapil, and Q. N. Porter, J. Chem. Soc. Perk. I, 1972, 599.
- 2 M. P. Cava and K. T. Buck, Tetrahedron, 1967, 25, 2795.
- 3 J. A. D. Jeffreys, J. Chem. Soc., 1956, 4451.
- 4 M. Tomita, T. Shingu, K. Fujitani, and H. Furukawa, Chem. Pharm. Bull., 1965, 13, 921.
- 5 M. P. Cava and A. Afzali, J. Org. Chem., 1975, 40, 1553.

Received, 18th December, 1975