

## THE PARTIAL SYNTHESIS OF BURNAMICINE

Shin-ichiro Sakai\* and Etsuji YamanakaFaculty of Pharmaceutical Sciences, Chiba University, Yayoi, Chiba, JapanLloyd J. DolbyDepartment of Chemistry, University of Oregon, Eugene, Oregon

The indole alkaloid burnamicine and its 19,20-dihydro derivative were partially synthesized from geissoschizine methyl ether and hirsutine respectively, using a C/D ring opening reaction with ethyl chloro-carbonate.

Burnamicine<sup>1)</sup> was found as the first cryptopine-type indole alkaloid from Hunteria eburnea Pichon ( Apocynaceae ) in 1963 by Taylor et al.. Subsequently, we prepared dihydroburnamicine (6b) from dihydrocorynantheine with a new C/D ring opening reaction.<sup>2)</sup>


We now report on the partial synthesis of burnamicine (6a) from geissoschizine methyl ether (1a) ( isolated from Uncaria rhynchophylla Miq. by us<sup>3)</sup> ) and of (6b) from hirsutine (1b) with another C/D ring opening reaction. It has recently been reported that chlorocarbonates are good reagents for the C/D ring cleavage of indole alkaloids and for the formation of vobasine type alkaloid from sarpagine type alkaloid.<sup>4)</sup>

Hirsutinol (2b) mp 168-169° and geissoschizol (2a) were formed from (1b) and (1a) respectively, by hydrolysis, decarboxylation and the subsequent NaBH<sub>4</sub> reduction. Reaction of ethyl chlorocarbonate with both alcohols (2a,b) in the presence of Na<sub>2</sub>CO<sub>3</sub> and an excess EtOH in CHCl<sub>3</sub> at room temperature gave rise to neutral 3,4-seco derivatives (3a,b) as epimeric mixtures ( in ca. 70% yield respectively ). The chlorination of (3a,b) with one equivalent

of *t*-BuOCl in  $\text{CH}_2\text{Cl}_2$ , followed by treatment with aq-N HCl gave the 2-acyl-indole derivatives (4a, in 50% yield, amorphous,  $\lambda_{\text{max}}^{\text{EtOH}}$  238, 316 nm,  $M^+$  m/e 384) and (4b, in 67% yield, amorphous,  $\lambda_{\text{max}}^{\text{EtOH}}$  238, 314 nm,  $M^+$  m/e 386). The reduction of (4a,b) with  $\text{LiAlH}_4$  in THF for 5hrs at room temperature yielded corresponding alcohols (5a, in 83% yield, amorphous, uv: indolic,  $M^+$  m/e 328) and (5b, in 97% yield, amorphous, uv: indolic,  $M^+$  m/e 330). The both alcohols (5a,b) were oxidized with active  $\text{MnO}_2$ <sup>5)</sup> in  $\text{CH}_2\text{Cl}_2$  to give burnamicine [ 6a, in 40% yield, mp 193-195°,  $[\alpha]_{\text{D}}^{24}$ : -240° in  $\text{CHCl}_3$ , CD (in dioxane)  $\Delta \epsilon_{\text{max}}$  (nm): -4.6 (333), +1.3 (310), -1.6(290), +8.1(260) ], which was identical spectroscopically with the authentic burnamicine<sup>1)</sup>, and dihydroburnamicine<sup>2)</sup> [ 6b in 38% yield, mp 98-100°,  $[\alpha]_{\text{D}}^{29}$ : +207° in dioxane, +125° in  $\text{CHCl}_3$ , CD (in dioxane)  $\Delta \epsilon_{\text{max}}$  (nm): -0.3(365), +2.0(335), -0.4(311), +1.8 (290), -5.8(261), +3.3(233) ].

Both bases (6a,b) exhibited nearly opposite circular dichroism spectra and this was presumably due to approximately antipodal ten membered ring construction with regard to the plane of 2-acylindole. Though the both bases (6a,b) showed the opposite optical activities, the absolute configuration of  $\text{C}_{15}$  of (6a,b) are the same as evident from the Scheme. The absolute configuration of natural burnamicine has now been determined unambiguously through the present chemical conversion from geissoschizine methyl ether (1a), which has been correlated<sup>3)</sup> with dihydrocorynantheine and corynantheidine with known absolute configurations.

ACKNOWLEDGEMENT We thank Dr. N.Finch, Dr. M.F.Bartlett and Dr. J.Karliner of CIBA-GEIGY Corporation for the identification of burnamicine and are also grateful to The Naito Foundation for financial support.




(1a)  $R_2 = \alpha H$

(2a)  $R_2 = \alpha H$

(3a,b)  $R_2 = \text{mixture } \alpha, \beta \text{ OC}_2\text{H}_5$

(1b)  $R_2 = \beta H$

(2b)  $R_2 = \beta H$



(4a,b)

(5a,b)  $R_2 = \text{mixture } \alpha, \beta \text{ OH}$

(6a,b)

series a,  $R_1 = =\text{CH}-\text{CH}_3$  (E). series b,  $R_1 = \alpha \text{ C}_2\text{H}_5$

i,  $\text{H}_2\text{O}$  ( $\text{H}^+$ ); ii,  $\text{NaBH}_4$ ; iii,  $\text{ClCO}_2\text{C}_2\text{H}_5 / \text{C}_2\text{H}_5\text{OH}, \text{CHCl}_3, \text{Na}_2\text{CO}_3$ ;

iv,  $t\text{-BuOCl}$ ; v,  $\text{H}_2\text{O}$  ( $\text{H}^+$ ); vi,  $\text{LiAlH}_4$ ; vii,  $\text{MnO}_2$ .

Scheme

REFERENCES

- 1 M.F.Bartlett and W.I.Taylor, J.Amer.Chem.Soc., 1963, 85, 1204.  
M.F.Bartlett, R.Sklar, A.F.Smith and W.I.Taylor, J.Org.Chem. 1963, 28, 2197.
- 2 L.J.Dolby and S.Sakai, J.Amer.Chem.Soc., 1964, 86, 5362; idem., Tetra-  
hedron, 1967, 23, 1.
- 3 E.Yamanaka, M.Fujiu, N.Aimi, S.Sakai and J.Haginiwa, unpublished data.
- 4 S.Sakai, A.Kubo, K.Katano, N.Shinma and K.Sasago, Yakugaku Zasshi, 1973,  
93, 1165.
- 5 J.Attenburrow, A.F.B.Cameron, J.H.Chapman, R.M.Evans, B.A.Hems,  
A.B.A.Jansen and T.Walker, J.Chem.Soc., 1952, 1094.

Received, 21st February, 1976