

A NEW SYNTHESIS OF
BENZO[C]PHENANTHRIDINE DERIVATIVE

Hideo Iida^{*}, Kohkichi Takahashi and Toyohiko Kikuchi

Tokyo College of Pharmacy

1432-1 Horinouchi, Hachioji, Tokyo Japan

4-Substituted isocarbostyryl derivative (V)
was reduced with lithium aluminum hydride to
give the 1,2-dihydro-4-substituted isoquinoline
which was heated with hydrochloric acid to give
two cis-hexahydrobenzo[c]phenanthridine deriva-
tives (VI) and (VII or VIII).

The benzo[c]phenanthridine ring system is found in a few
alkaloids¹. Several methods have been described^{2,3,4} for the
construction of the ring system.

In connection with our studies on reaction of isocarbostyryl
derivatives⁵, we have investigated the synthesis of dibenzo-
[c]phenanthridine derivatives, which involve new synthesis
of 4-substituted isocarbostyryl derivative (V) from 4-sub-
stituted homophthalimide derivative (III).

4-(3,4-Dimethoxyphenacyl)-2-methylhomophthalimide (III),
m.p. 155-156° ; ir $\nu_{\max}$ (nujol) 1700 and 1650 cm^{-1} ; mass m/e


353 (M^+) and nmr δ ($CDCl_3$) 3.42 (3H, s, N- CH_3), 3.84 (3H, s, O- CH_3), 3.91 (3H, s, O- CH_3) and 4.01 (2H, d, $J=4Hz$, CH_2) was prepared in 70-80% yield by treating 2-methylhomophthalimide (I) with 3,4-dimethoxyphenacyl bromide (II) in the presence of sodium ethoxide.

Catalytic reduction of this compound (III) on 5% palladium charcoal in acetic acid containing perchloric acid at 60° and atmospheric pressure gave 4-(3,4-dimethoxyphenethyl)-2-methylhomophthalimide (IV) in 90% yield as an oil, ir ν_{max} (film) 1700 and 1650 cm^{-1} ; mass m/e 339 (M^+) and nmr δ ($CDCl_3$) 2.48 (4H, m, CH_2-CH_2), 3.30 (3H, s, N- CH_3), 3.85 (6H, s, O- $CH_3 \times 2$) and 4.00 (1H, t, $J=4Hz$, C_4-H).

Treatment of the imide (IV) with sodium borohydride afforded the 3,4-dihydro-3-hydroxyisocarbostyryl, which was then acidified with 10% hydrochloric acid to give the isocarbostyryl (V), m.p. 133-134°; ir ν_{max} (nujol) 1650 cm^{-1} ; mass m/e 323 (M^+) and nmr δ ($CDCl_3$) 2.90 (4H, s, CH_2-CH_2), 3.50 (3H, s, N- CH_3), 3.84, 3.86 (6H, s, O- $CH_3 \times 2$) and 6.78 (1H, s, C_3-H).

4-(3,4-Dimethoxyphenethyl)-2-methylisocarbostyryl (V) was treated with lithium aluminum hydride to give 1,2-dihydro-4-substituted isoquinoline, which was heated with concentrated hydrochloric acid to give two compounds, which were separated by column chromatography on silica gel with chloroform as eluent.

The first compound was obtained in 40% yield, m.p. 120-122°;



ir $\nu_{\max}$ (nujol) 1600 cm^{-1} ; mass m/e 309 (M^+) and nmr $\delta(\text{CDCl}_3)$ 2.40 (3H, s, N-CH₃), 3.58, 4.00 (2H, ABq, $J=20\text{Hz}$, C₆-H), 3.60 (1H, d, $J=4.5\text{Hz}$, 4b-H), 3.85, 3.90 (6H, s, O-CH₃x2) and 6.60, 6.93 (2H, s, C₁-H and C₄-H).

The nmr spectrum of the product (VI) also supported the benzo[c]phenanthridine structure and the coupling constant shown by the 4b-proton suggested that BC ring junction was *cis*.

A similar result of the coupling constant of 4b-proton in the benzo[c]phenanthridine was reported by Ninomiya et al.⁶

The compound (VI) was determined as *cis*-4b,5,6,10b,11,12-hexahydro-2,3-dimethoxy-5-methylbenzo[c]phenanthridine.

The second compound was obtained in 50% yield, a single product, m.p. 250-252° ; ir $\nu_{\max}$ (nujol) 3450 and 1600 cm^{-1} ; mass m/e 295 (M^+) and nmr $\delta(\text{CDCl}_3)$ 2.30 (3H, s, N-CH₃), 3.68 (1H, d, $J=4.5\text{Hz}$, 4b-H), 3.70, 3.97 (2H, ABq, $J=20\text{Hz}$, C₆-H₂), 3.80 (3H, s, O-CH₃) and 6.60, 6.93 (2H, s, C₁-H and C₄-H).

The mass and nmr spectra of the second product were indicated demethylation of the benzo[c]phenanthridine (VII or VIII).

The position of demethylation was not determined yet.

REFERENCES

- 1 R.H.F. Manske, and H.L. Holmes, The Alkaloids Vol IV; Chap 35, Academic Press, New York (1954).

- 2 A.S. Bailey, and R. Robinson, J. Chem. Soc., 1950, 1375.
A.S. Bailey, R. Robinson, and R.S. Staunton, J. Chem. Soc.,
1950, 2277.
- 3 S.F. Dyke, M. Sainsbury, and B.J. Moon, Tetrahedron, 1968,
24, 1467.
- 4 W.J. Gensler, S.F. Lawless, A.L. Bluhm, and H. Dertouzos,
J. Org. Chem., 1975, 40, 733.
- 5 H. Iida, K. Kawano, T. Kikuchi, and F. Yoshimizu,
J. Pharm. Soc., Japan, 1967, 96, 176.
- 6 I. Ninomiya, T. Naito, T. Kiguchi, and T. Mori, J. Chem. Soc.,
Perkin 1, 1973, 1696.

Received, 13th July, 1976