

MICHAEL ADDITIONS OF INDOLES TO 2-OXOINDOLIN-
3-YLIDENE KETONES

Akinori Kubo^{*}, Tatsuya Nakai, and Toshikazu Nozoye

Meiji College of Pharmacy, Setagaya, Tokyo, Japan

Michael-type additions to (E)-2-oxoindolin-3-ylidene ketones(1) and indoles(2) gave rise to C-1'(3) and/or C-3(4) adducts depending on the oxindolic N-substituents and the C-1' adducts(3) were converted into new 9H-pyridazino[3,4-b]indole derivatives(6).

2-Oxoindolin-3-ylidene ketones(1) are of considerable interests in their reactivities because of the presence of two α,β -unsaturated carbonyl systems in the molecule. Tacconi and co-workers¹⁾, in a series of investigations, have studied the reactions of (E)-2-oxoindolin-3-ylideneacetophenone(1a) with enamines and ethyl vinyl ethers undergoing heterodiene cycloadditions. It has been concluded that (E)-2-oxoindolin-3-ylideneacetophenone(1a) underwent 1,2 or 1,4-cycloaddition with aldo-enamines depending on the oxindolic nitrogen substituents. These results led us to study the effect of N-substituents on Michael-type additions.

We wish to report here the influence of N-substituents in (E)-

2-oxoindolin-3-ylidene ketones(1) on Michael-type additions²⁾ with indole derivatives as typical enamine and the structures of the unusual minor products in these reactions.

Indoles were found to react readily with (E)-2-oxoindolin-3-ylideneacetophenone(1a) in a refluxing acetic acid solution³⁾ for 3 hr in N₂ to afford the corresponding adducts in good yields⁴⁾ (Table 1). In a typical experiment evaporation of the solvent and subsequent chromatographic separation on silica gel gave C-1' adduct(3a), 75.5%, as the mixture of *dl*-threo and *dl*-erythro isomers and C-3 adduct(4a), mp 251-254°, 2.8%, $\lambda_{\max}^{95\% \text{EtOH}}$ nm(log ϵ): 221(4.65), 246(4.36), 281(3.98), 291(3.91); $\nu_{\max}^{\text{KBr}}$ cm⁻¹: 3310, 1721; δ (DMSO-d₆): 4.26 and 4.51(d, J=18Hz, 1H), 10.38 and 10.90(s, 1H, NH). Recrystallization resulted successfully in isolation of each *dl*-diastereomer, 3a-I, mp 165-179°, 71%, $\lambda_{\max}^{95\% \text{EtOH}}$ nm(log ϵ): 218(4.66), 245(4.36), 283(3.94), 291(3.88); $\nu_{\max}^{\text{KBr}}$ cm⁻¹: 3360, 1705; δ (DMSO-d₆): 4.49 and 5.59(d, J=6Hz, 1H), 10.09 and 10.69(s, 1H, NH) and 3a-II, mp 224-226°, 4.5%, δ (DMSO-d₆): 4.16 and 5.78(d, J=6Hz, 1H), 10.19 and 10.80(s, 1H, NH). The gross structures of C-1' and C-3 adducts were determined easily by their mass spectra, 3a showed intense peaks at M-105(COPh)(A-fission) along with m/e 234(base), 206(B-fission) and 4a showed the only intense peak at m/e 247(M-CH₂COPh)(base).

These results(Table 1) indicate that if R₁ is an electron-donating substituent the 3-position of the oxindole becomes more activated and the addition leads to C-3 adduct(4) in a moderate yield together with C-1' adduct(3), whereas, for the electron-attracting substituent the addition leads to C-1' adduct(3) exclusively.





Table 1 Addition of Indoles to 2-Oxoindolin-3-ylidene Ketones


	1		2		Reaction	3	4	5
	R ₁	R ₂	R ₃	R ₄	Conditions	Yield (%)	Ratio of Diastereomers	Yield (%)
a)	H	Ph	H	H	AcOH, reflux 3hr	75.5	15.8:1	2.8
b)	H	Ph	H	CH ₃		91	2:1	—
c)	CH ₃	Ph	H	H		45	3:1	31
d)	CH ₃	Ph	H	CH ₃		71	1.7:1	12
e)	COCH ₃	Ph	H	CH ₃		38+47(3b)	—	—
f)	COPh	Ph	H	H		—	—	—
g)	H	CH ₃	H	H		76	—	11
h)	H	CH ₃	CH ₃	H		65	—	—
a)	H	Ph	H	H	BF ₃ ·Et ₂ O	52	1.5:1	5
g)	H	CH ₃	H	H	90°, 10min	30	—	—
h)	H	CH ₃	CH ₃	H		40	—	—

Table 2 9H-Pyridazino[3,4-b]indole Derivatives(6)


	R ₁	R ₂	R ₃	mp(°C)	Yield(%)
6a	H	Ph	H	>300	33
6c	CH ₃	Ph	H	291.5	29
6g	H	CH ₃	H	262-6	35
6h	H	CH ₃	CH ₃	>300	17



Under similar conditions, (E)-2-oxoindolin-3-ylideneacetone (1g) afforded C-1' adduct(3g), mp 205-208°, 76%, m/e 304 (M^+ , 21), 261 (45, A) and 172 (base), 144 (49, B); δ (DMSO- d_6): 2.16(s, 3H), 4.37 and 4.7: (d, J=6Hz, 1H), together with by-product(5g), mp >300°, 11%, whose structural assignment was based on the following evidences, $C_{27}H_{21}ON_3$, m/e 403 (M^+ , 52), 388 (M-15, base); $\nu_{max}^{KBr} cm^{-1}$: 1714, 1619; δ (DMSO- d_6) 2.03(s, 3H), 2.94 and 3.53(d, J=14Hz, 1H), 10.35, 10.50 and 10.55(s, 1H, NH); positive to Ehrlich reaction.



5g



5a

Furthermore, we examined this Michael-type addition in $BF_3 \cdot Et_2O$ at 90° for 10 min⁵⁾. Thus 1a gave the C-1' adduct(3a), 52%, along with a minor product(5a), mp 284-285.5°, 12%, whose structure was determined by the following data, $C_{24}H_{18}ON_2$, m/e 350 (M^+ , base); $\nu_{max}^{KBr} cm^{-1}$: 1712, 1620; δ (C_5D_5N): 2.95(dd, J=6, 14Hz, 1H), 3.93(dd, J=8, 14 Hz, 1H), 5.39(dd, J=6, 8Hz, 1H); negative to Ehrlich reaction.

Formation of 5a and 5g can reasonably involve 'C-3 adduct' intermediate followed by intramolecular cyclization during the reaction. Investigation of this point is continuing in our laboratory.

Finally 3a, 3c, 3g and 3h were converted into new 9H-pyridazino-[3,4-b]indole derivatives(6) on treatment with hydrazine hydrate⁶⁾

in acetic acid (Table 2).

As the result, we demonstrate that (E)-2-oxoindolin-3-ylidene ketones(1) are the excellent Michael acceptors under acidic conditions and the influence of the oxindolic N-substituents can be regarded as the determinant of the mode of Michael-type addition.

REFERENCES

- 1 a)G.Tacconi,A.Gamba,F.Marinone and G.Desimoni, Tetrahedron, 1971, 27, 561. b)G.Tacconi,F.Marinone,A.Gamba and G.Desimoni, ibid., 1972, 28, 1517. c)G.Tacconi,P.Iadarole,F.Marinone, P.P.Righetti and G.Desimoni, ibid., 1975, 31, 1179. d)G.Desimoni,A.Gamba,M.Onticelli,M.Nicola and G.Tacconi, J.Am.Chem.Soc., 1976, 98, 2947. e)G.Tacconi,A.C.Piccolini,P.P.Righetti,E.Selva and G.Desimoni, J.Chem.Soc.Perkin I, 1976, 1248. f)G.Desimoni and G.Tacconi, Chem.Rev., 1975, 75, 651.
- 2 R.L.Autrey and F.C.Tahk, Tetrahedron, 1968, 24, 3337.
- 3 W.E.Noland and M.R.Venkiteswaran, J.Org.Chem., 1961, 26, 4263.
- 4 Satisfactory elemental analytical, mass spectral and nmr data have been obtained for all new compounds.
- 5 M.Ando,G.Bölchi and T.Ohnuma, J.Am.Chem.Soc., 1975, 97, 6880.
- 6 G.Kobayashi and S.Furukawa, Chem.Pharm.Bull.(Tokyo), 1964, 12, 1129.

Received, 16th August, 1976