

A SIMPLE SYNTHESIS OF 2,2',5',2''-TERTHIENYL

Takashi Asano, Shota Ito, Norio Saito and Kiyotaka Hatakeda

Government Industrial Research Institute, Tohoku



Nigatake, Haranomachi, Sendai 983, Japan

The synthesis of 2,2',5',2''-terthienyl (IV) is reported. A key intermediate in this synthesis was 1,4-di(2-thienyl)-1,4-butanedione (III), which was prepared from trimethylsilyl enol ether (II) of 2-acetylthiophene (I). Corresponding furyl derivatives were also synthesized by the similar procedure.

Since 2,2',5',2''-terthienyl (IV) was isolated from *Tagetes* plants¹, considerable attention have been given to synthesize it and some methods have been reported in literatures², because of its high nematocidal activity against nematodes such as *Ditylenchus dipsaci* and *Heterodea rostochiensis*³.

However those methods are laborious and/or the yields are not satisfactory. In this paper, we wish to report a simple synthesis of IV from commercially available 2-acetylthiophen (I) in three steps.

Our synthetic scheme is based on preparation of 1,4-di-(2-thienyl)-1,4-butanedione (III) as a key intermediate derived from 2-acetylthiophene (I). Recently attention has been paid to the preparation of 1,4-butanedione derivatives⁴, and as a new synthetic method, it was reported that trimethylsilyl enol ether of acetophenone gave 1,4-diphenyl-1,4-butanedione⁵. We prepared III by a modification of the method.



The reaction of I with trimethylchlorosilane (1.3 equiv) and triethylamine (2.5 equiv) in dimethylformamide gave trimethylsilyl enol ether (II) in an excellent yield (90%). The physical data of II are as follows: bp 73°/2 Torr: M⁺ 198

(C₉H₁₄OSSi): IR (liquid); no carbonyl, 1617 (shoulder) and 1595 cm⁻¹: nmr (CCl₄); δ 0.36 (s, 3H, Si-Me), 0.41 (s, 6H, Si-Me), 4.23 and 4.71 (d, each 1H, J=1.7, =CH₂), and 6.81-7.58 (m, 3H, ring H). Stirring of II with silveroxide (2 equiv) in dimethylformamide on a steam bath afforded pale yellow crystals of III in 71% yield: mp 130° (130-131° in lit.^{2(b)}): M⁺ 250 (C₁₂H₁₀O₂S₂): nmr (CDCl₃); 3.32 (s, 4H, COCH₂), 7.04-7.82 (m, 6H, ring H): IR (CHCl₃); 1638 cm⁻¹ (C=O). Treatment of III with hydrogen chloride and hydrogen sulfide in ethanol gave a greenish yellow substance (IV) in 66% yield: mp 92-93° (94-95.5° in lit.¹). Its elemental analysis (C₁₂H₈S₃) and UV spectrum data confirm that the structure of IV is 2,2',5',2"-terthienyl¹.

Similarly, when 2-acetylfuran (V) was used as a starting material, trimethylsilyl enol ether (VI) and 1,4-di(2-furyl)-1,4-butanedione (VII) were obtained by the same procedure as above, and treatment of VII with acetic anhydride and hydrochloric acid gave pale yellow crystals of 2,2',5',2"-terfuryl (VIII) (overall yield 65%). These substances possess following properties. VI, bp 48-49°/3 Torr: IR (liquid); 3120, 2950, 1635 (shoulder) and 1618 cm⁻¹: M⁺ 182 (C₉H₁₄O₂Si). VII, mp 128-130° (132 in lit.⁶): M⁺ 218 (C₁₂H₁₀O₄). VIII, mp 62-63°: IR (CHCl₃); 3130, 1580, 1522, 1430 and 998 cm⁻¹: M⁺ 200 (C₁₂H₈O₃): nmr (CDCl₃) δ 6.33-6.66 (m, 6H) and 7.33-7.48 (m, 2H).

REFERENCES

- 1 L. Zechmeister and J. W. Sease, J. Amer. Chem. Soc., 1947, 69, 273.
- 2 (a) J. W. Sease and L. Zechmeister, J. Amer. Chem. Soc., 1947, 69, 270; (b) H. J. Kooremen and J. Wynberg, Rec. Trav. Chim. Pays-Bas, 1967, 86, 37; (c) M. Yamazaki, N. Onodera, Y. Kurosu and H. Kawada, Japan Kokai, 75 49,273.
- 3 (a) T. Yokoo, Öyokenkyu Hokoku, 1958, 1; (b) J. H. Uhlenbroek and J. D. Bijloo, Rec. Trav. Chim. Pays-Bas, 1958, 77, 1004.
- 4 (a) E. J. Corey and L. S. Hegedus, J. Amer. Chem. Soc., 1964, 91, 4926; (b) J. E. McMurry and J. Melton, J. Amer. Chem. Soc., 1971, 93, 5309; (c) J. Ficini and J. P. Genet, Tetrahedron Letters, 1971, 1565; (d) J. E. Richman and J. L. Herrmann, Tetrahedron Letters, 1973, 3267; (e) E. Ghera, D. H. Perry and S. Shoua, J. Chem. Soc., Chem. Commun., 1973, 858; (f) M. Araki, S. Sakata, H. Takei and T. Mukaiyama, Chem. Lett., 1947, 687.
- 5 Y. Ito, T. Konoike and T. Saegusa, J. Amer. Chem. Soc., 1975, 97, 649.
- 6 E. Niwa and M. Miyake, Chem. Ber., 1969, 102, 1443.

Received, 11th January, 1977