


PHOTOCHEMISTRY OF BRIDGED STILBENE OXIDES

Kazuyoshi Nishiyama, Kiyoyasu Ishikawa, Ila Sarkar, David C. Lankin, and
Gary W. Griffin*

Department of Chemistry, University of New Orleans, New Orleans, Louisiana
 70122


The photochemistry of a pair of stilbene oxides in which the 2,3-positions are bridged by three and four methylene groups has been studied and compared with a lower homolog. Evidence is provided that cyclic ylides are produced upon photolysis of these oxiranes in solution at 254 nm. The primary photoproducts derived from the ylides in turn are β -styrylpropio- and β -styrylbutyrophenones, respectively. In the latter case Norrish type II cleavage then occurs to give as final products, 1-phenylbutadiene and acetophenone. The bridged oxiranes were shown to be more photostable than their open chain analogs and on the basis of double irradiation experiments it was established that a carbene rather than a conventional diradical process is operative or competes in the ylide photocleavage and ketone formation.

The photochemistry of aryl (1) and arene oxides (2) has been the subject of extensive study in our laboratories. Among common reactions of the former are [3+2+1]cycloeliminations to give carbonyl compounds and carbenes (1a-c) presumably through carbonyl ylides which have been intercepted in several cases with alkenes (1c-e, 1g-h) and nucleophiles (1d) (Eq. 1). Recently, we have observed that epoxydiphenylmaleic anhydride,



which forms a remarkably stable cyclic ylide, does not photofragment in the conventional manner (1f). In contrast carbon dioxide is lost upon photolysis in solution (25°) and, unlike the closely related dicarboxy- or dicarbomethoxystilbene oxide analogs, no carbenes are detected as primary photoproducts (3,4). This observation rekindled our interest in the photochemical behavior of bridged stilbene oxides and we wish to summarize our previously unpublished as well as supplementary work in this area.


While our investigations in this area were in progress it was established that the bridged oxides 1,4-diphenyl-2,2,3,3-tetramethyl-5-oxabicyclo-[2.1.0]pentane (**1a**) (as well as the 4,4'-dibromodiphenyl analog **1b**) and the related monoaryl glycidic ester **1c** undergo thermolysis to give stable ylides **2** which, at least in the case of **1a** and **1c**, may be trapped by dipolarophiles including benzonitrile which gives the unstable adducts **3** (Eq. 2) which in turn undergo retrograde cleavage to oxazoles and 2,3-dimethyl-2-butene (**5**).



While ylide formation is also observed upon photolysis of **1a-c** (254 nm) no evidence for photocleavage of the oxiranes **1** or ylides **2** to give keto-carbenes **4** and/or **5** is presented. In fact, reportedly, the oxide **1a** is remarkably stable. Repetitive cycles of thermolysis (or photolysis) and constrained recyclization in the disrotatory fashion (dictated by steric factors) leads only to limited decomposition (5,6).

Although the photostability of the higher homologs of **1a**, namely **6** and **7** exceeds that of *cis*-dimethylstilbene oxide (**1j**), which photolyzes to an acyclic ylide (*vide infra*), the dramatic photostability attributed to **1a** is absent although in both cases photochromic behavior is observed at 77°K. Typically a benzene solution (0.1 M) of the oxide **6** (mp 104°C) (**7**) prepared by peracetic acid oxidation of 1,2-diphenylcyclopentene was irradiated


(254 nm)(8) in a quartz vessel for 5 h. The major products formed and isolated by preparative glc were identified as *cis*- and *trans*-1,5-diphenyl-4-penten-1-one (**8a** and **8b**, 2:1, respectively) (Eq. 4). The identity of **8b** [nmr CCl_4) τ 2.0-2.8 (m, aromatic and vinyl protons), 3.75 (t, vinyl protons), and 6.8-7.5 (m, methylene protons)] was confirmed by comparison of spectral data with that of an authentic sample (mp 58-59°C) prepared in an alternate fashion (9).



In addition, ozonolysis of **8a** [nmr (CCl_4) τ 2.0-2.7 (m, aromatic protons), 3.58 (d, vinyl protons), 4.1-4.6 (m, vinyl protons), 7.0-7.4 (m, methylene protons)] and subsequent sequential treatment of the products with hydrogen peroxide and diazomethane gave as expected, dimethyl succinate and methyl 3-benzoylpropionate. Finally, the thermodynamically less stable *cis* isomer **8a** was converted, as expected, to the *trans* isomer **8b** upon treatment with iodine in benzene at the reflux temperature (80°C). A minor (> 1%) crystalline photoproduct (mp 280°C) is formed upon prolonged irradiation of **6** (26 h) which was assigned structure **9a** or **9b** on the basis of ultraviolet and infrared spectral as well as combustion analytical data (10,11). That the thermodynamically less stable isomer **8a** is the predominant photoproduct may only reflect the fact that secondary photoequilibration of the primary products is relatively rapid. This contention is supported by the observation that direct irradiation of a benzene solution of **8b** under conditions simulating those employed with **6** gives rise to the same product mixture of **8a** and **8b** (2:1, respectively).


Photolysis (254 nm) (8) of the oxide **7** (mp 77-78°C) prepared by peracetic acid oxidation of 1,2-diphenylcyclohexene (**12**), under conditions identical to those employed for preparation of **6**, gives four products (**10a**, **10b**, **12**, and **13**), which were separated and isolated by preparative glc (Eqs. 5 and 6). In addition to the isomeric unsaturated ketones **10a** and **10b** [**10a**, nmr (CCl_4) τ 2.5-2.7 (m, aromatic and vinyl protons), 3.69 (t, vinyl protons) and 7.0-8.0 (m, methylene protons)] the secondary photoproducts acetophenone (**12**) and

trans-1-phenylbutadiene (13) prepared independently (14) are also formed and were identified by comparison with authentic samples. It is reasonable to assume that these products arise by γ Norrish Type II cleavage *via* the






diradical intermediate **11**, derived from **10a** and/or **10b**, which have allylic gamma hydrogens (Eq.6) (15). In the case of **8a** and **8b** much less reactive vinylic hydrogens appear at the γ -position and only a trace of the fragmentation product **12** was detected by glc.

A priori at least two viable mechanisms may be invoked to rationalize the conversion of **6** and **7** to the isomeric pairs of unsaturated ketones **8** and **10**, respectively. Extensive precedent may be found in the literature (2d) (16) for C-O bond photolysis to give diradical(s) such as **14**. 1,2-Migration is also recognized as a reaction characteristic of excited state processes (17a) if not their ground state counterparts (17b) and in the case of **14** would afford the 1,4-diradical **15** which in turn would undergo fragmentation to the observed products **8** and **10** (Eq. 7). This mechanism, it is true, suffers from the disadvantage that a less stable diradical **15** must be formed from **14** although the intramolecular transfer of hydrogen from even methyl substituents is a well-documented photoreaction first recognized in our laboratories (17a).



An alternative mechanistic pathway in which the photogenerated carbonyl ylides **16** are implicated may be operative. Subsequent photocleavage to the ketocarbene **17** (**1c**) followed by 1,2-hydrogen migration (**17**) (Eq. 8) would also account for the observed conversion. If, in fact, the ylide mechanism is operative (despite our inability to observe luminescence and epr spectra anticipated for **17**), then enhancement of the conversion should be observed upon simultaneous irradiation of **6** and **7** with radiation in the visible and ultraviolet regions [i.e., through double irradiation studies (**1c**) *vide infra*]. This assertion is dependent upon the validity of our contention that the ylide **16** will absorb in the visible and the reasonable assumption while the 1,4-diradical **15** should be transparent in this spectral region. A comparison of the absorption spectral data for the intermediates (presumed to be carbonyl ylides) (**1**) formed upon irradiation of **6** and **7** with those formed from *cis*-dimethylstilbene oxide are presented in Table 1 and attest to the fact that bridging does not greatly affect the nature of the intermediate.

Table 1
Spectral Properties of the Species Generated Upon Photolysis of the
Oxides 6 and 7a)

	λ_{max}	405,	456,	528
	λ_{max}	399,	418 (shoulder)	525
	λ_{max}	405,	457	537

a) Photolyses (254 nm) (8) were conducted in 2-methyltetrahydrofuran at 77°K.

That the proposed ylides 16 are formed upon photolysis of oxides 6 and 7 was confirmed by interception to give the expected cycloadducts 18, 19, 20, and 21 among the photoproducts obtained from 6 and 7 using the dipolarophiles fumaronitrile (Eqs. 9 and 10) and maleonitrile in acetonitrile. The results of the photocycloaddition of 6 and 7 to fumaro- and maleonitrile are presented in Table 2. Structures 18 and 20 may be assigned to the major photoadducts obtained from fumaronitrile and the oxides 6 and 7, respectively. On the other hand, while the


Table 2
Cycloadducts Obtained from 6 and 7 and Maleo- and Fumaronitrile

Oxide	Dipolarophile	% Cycloadduct ^{a,b}			
		<u>18</u>	<u>20</u>	<u>19</u>	<u>21</u>
6	Fumaronitrile	40.5		4.7	
	Maleonitrile	5.6		33.0	
7	Fumaronitrile		38.1		4.5
	Maleonitrile		9.1		34.3


a) Irradiation (254 nm) (8) for 5 h at 40° in acetonitrile as a solvent.

b) The values reported were determined by actual isolation using thick-layer and column chromatographic techniques.

cis adducts 19 and 21, like the *trans* adducts 18 and 20, have been characterized spectroscopically (Table 3), the structures of the former pair remain to be elucidated. Clearly, however, the stereospecificity of the cycloaddition



reactions of 6 and 7 is high although precisely how high must yet be established and is complicated by competitive photoisomerization of the dipolarophile. Furthermore, it should not be overlooked that these cycloaddition reactions, although photoinitiated, generally occur in the ground state (Ic-h).






The utility of frontier orbital theory in selecting the most favorable dipolarophiles (electron-rich or electron-poor) should be noted. On the basis of frontier orbital energies, Houk (18) suggested that ylides substituted with alkyl or conjugating substituents should react readily with electron-deficient dipolarophiles and less readily or not at all with electron-rich adducts. The reaction of electron-deficient dipolarophiles with carbonyl ylides of the conjugated type 16 described herein are classified as HOMO controlled; i.e., the lowest unoccupied orbitals of the dipolarophiles interact with the highest occupied orbitals (HOMO) of the ylide. Our inability as well as that of others (1g) to detect adducts of 6 with those of electron-rich dipolarophiles and *cis*- and *trans*-butene-2 supports the predictions.

Table 3
Spectral Data and Physical Properties of Adducts from 18 - 21

Adduct	mp	
18 (trans)	178°C	Mass Spectrum M ⁺ 314 IR (Nujol) 2250 cm ⁻¹ (-CN) NMR (CDCl ₃) τ 2.2-2.8 (m, aromatic, protons) 6.37 (q, methine protons), 7.2-8.4 (m, methylene protons)
19 (cis)	176°C	Mass Spectrum M ⁺ 314 IR (Nujol) 2240 cm ⁻¹ (-CN) NMR (CDCl ₃) τ 2.2-2.8 (m, aromatic, protons) 6.46 (s, methine protons), 7.2-8.4 (m, methylene protons)
20 (trans)	182°C	Mass Spectrum M ⁺ 328 IR (Nujol) 2245 cm ⁻¹ (-CN) NMR (CDCl ₃) τ 2.2-2.8 (m, aromatic protons) 6.27 (q, methine protons), 7.1-8.4 (m, methylene protons)
21 (cis)	183°C	Mass Spectrum M ⁺ 328 IR (Nujol) 2245 cm ⁻¹ (-CN) NMR (CDCl ₃) τ 2.4-2.8 (m, aromatic protons) 6.23 (s, methine protons), 6.9-8.3 (m, methylene protons)

While spectral and chemical evidence has been provided for the generation of the cyclic carbonyl ylides 16 under photolytic conditions where conversion of 6 and 7 to 8 and 10, respectively, occurs, additional evidence is required to confirm that the ylides are indeed intermediates. To this end a comparison of the stabilities of 6 and 7 relative to that of *cis*-dimethylstilbene oxide was conducted (Table 4).

Table 4
The Photostability of the Oxides 6 and 7 Relative to that of *cis*-Dimethyl-Stilbene Oxide

Oxide	Reaction Time (h)	% Recovery of Starting Material ^a
	1	72
	3	41
 6	1	80
	3	61
 7	1	84
	3	68

^aIrradiations (254 nm) (8) were conducted under similar conditions (40°) in cyclohexane as a solvent and the residual oxide was isolated by tlc.

Inspection of the data shown in Table 4 indicates that the cyclic ylides 6 are indeed somewhat more stable under the conditions utilized than the corresponding open chain ylide derived from *cis*-dimethylstilbene oxide; however the marked photostability reported for 1a (5a) is not an intrinsic property of the next higher homologs 6 and 7. The significant feature, however, is that simultaneous irradiation with visible light (double irradiation) accelerates the conversion of starting material in the case of 6 and 7. For example, when the substrate 7 was subjected to double irradiation (1c) only 34% was recovered relative to 44% upon direct irradiation for 3 h (40°) in cyclohexane. This result supports the contention that the conversion of 6 and 7 to 8 and 10 may occur at least in part, according to the ketocarbene mechanism depicted in Eq. 8.

ACKNOWLEDGMENT.

The authors wish to acknowledge support by National Cancer Institute (DHEW) (grant CA-18346-01,2) and the National Science Foundation (grant CHE14831) for this work. They also wish to thank Dr.E.Elder and Ms.J.Thompson for aid in the preparation of the manuscript.

REFERENCES

- ¹ (a) Presented in part at the 28th SE American Chemical Society Regional Meeting, Gatlinburg, Tenn. Oct. 27-29, 1976; (b) N.R.Bertoniere and G.W. Griffin, Chap. 3 in "Carbenes", I, M.Jones, Jr. and R.A.Moss, Eds., John Wiley and Sons, New York, N.Y., 1973, pp 305-359; (c) G.W.Griffin, K.Ishikawa, and I.J.Lev, J.Am.Chem.Soc., 1976, 98, 5697; (d) I.J.Lev, K.Ishikawa, N.S. Bhacca, and G.W.Griffin, J.Org.Chem., 1976, 41, 2654; (e) K.Ishikawa, G.W. Griffin, and I.J.Lev, ibid., 1976, 41, 3747; (f) G.W.Griffin, K.Nishiyama, and K.Ishikawa, ibid., 1977, 42, 180; (g) G.A.Lee, ibid., 1976, 41, 2656; (h) V. Markowski and R.Huisgen, Tetrahedron Lett., 1976, 4643. It should be noted that our original proposal that carbonyl ylides were implicated in carbene formation from oxiranes was made in 1967 (1h) and which has been overlooked by these and other investigators (1i). See also Ref. 16, p 327. Furthermore, a widely circulated abstract (G.W.Griffin, I.Lev, and K.Ishikawa, Vth IUPAC Symposium on Photochemistry, Enschede, The Netherlands, July 22-26, 1973, contr. paper No.20, p 62) submitted prior to publication of Dr.Markowski's thesis (1h) addresses problems related to the stereochemistry of ylide opening and adduct formation. (i) P.Petrellis and G.W.Griffin, J.Chem.Soc.,Chem.Commun., 1967, 691; ibid., 1968, 1099; (j) H.Kristinsson, Tetrahedron Lett., 1966, 2343.
- ² (a) N.E.Brightwell Ph.D.Dissertation, University of New Orleans, New Orleans La., 1974; (b) N.E.Brightwell and G.W.Griffin, J.Chem.Soc.,Chem.Commun., 1973, 37; (c) B.J.Dowty, N.E.Brightwell, J.L.Laseter, and G.W.Griffin, Biochem. Biophys.Res.Commun., 1974, 57, 452; (d) G.W.Griffin and A.Padwa, chap.2 in "Photochemistry of Heterocyclic Compounds", O.Buchardt, Ed., John Wiley and Sons, Inc., New York, N.Y., 1976; (e) G.W.Griffin, K.Ishikawa, and S.K.Satra, J.Heterocycl. Chem., 1976, 13, 1369; (f) K.Ishikawa, H.C.Charles, and G.W.Griffin, Tetrahedron Lett., 1977, 427; (g) K.Ishikawa and G.W.Griffin, Angew.Chem., 1977, 89, 181; Angew.Chem.Int.Ed.Engl., 1977, 16, 171.
- ³ G.W.Griffin, D.M.Gibson, and K.Ishikawa, J.Chem.Soc.,Chem.Commun., 1975, 595.
- ⁴ K.Nishiyama and G.W.Griffin, unpublished results.
- ⁵ (a) D.R.Arnold and L.A.Karnischky, J.Am.Chem.Soc., 1970, 92, 1404; (b) D.R. Arnold and Y.C.Chang, J.Heterocycl.Chem., 1971, 8, 1097; (c) D.R.Arnold, A.B.Evvin, and L.A.Karnischky, Pure Applied Chem., 1970, 24, 523.

- 6 See E.F.Ullman and J.E.Milks, J.Am.Chem.Soc., 1964, 86, 3814 and references cited therein, as well as H.E.Zimmerman and R.D.Simkin, Tetrahedron Lett., 1964, 1847, for additional examples of photogenerated cyclic carbonyl ylides; however, in these cases of substituted indenone oxides and related compounds the lowest excited state is probably n,π^* rather than π,π^* in character.
- 7 R.Criegee, A. Kerckow and H.Zinke, Chem.Ber., 1955, 88, 1878.
- 8 Irradiations were conducted in serum-capped quartz or Pyrex test tubes as indicated and a Rayonet RPR-100 chamber reactor equipped with 16 8-w 254 nm or 350 nm lamps was employed as a light source.
- 9 (a) We are indebted to Prof.A.Padwa for providing us with an authentic sample of 8b to employ for purposes of comparison. (b) C.G.Overberger and J.J. Monagle, J.Am.Chem.Soc., 1956, 78, 4470; (c) For a study of intramolecular triplet energy transfer in the bichromophoric systems 8 and 10 see D.O.Cowan and A.A.Baum, ibid., 1970, 92, 2153.
- 10 The intramolecular cyclization of γ,δ -unsaturated ketones to give bicyclic oxetanes of the types 9a and 9b is not without precedent. (a) N.C.Yang, M.Nussim and D.R.Coulson, Tetrahedron Lett., 1965, 1525; (b) R.Srinivasan, J.Am.Chem.Soc., 1960 82, 775; (c) S.R.Kurowsky and H.Morrison, ibid., 1972, 94, 507.
- 11 Satisfactory combustion analyses were obtained on all new compounds by Galbraith Laboratories, Inc., Knoxville, Tenn. or Integral Microanalytical Laboratories, Inc., Raleigh, N.C.
- 12 P.Tomboulia, J.Org.Chem., 1961, 26, 2652.
- 13 While an emission "similar" to that of acetophenone was observed upon photolysis of 7 (77°K) which might be attributed to a ketocarbene, it is now clear that acetophenone itself is indeed produced as a secondary photolysis product. See R.S.Becker, R.O.Bost, J.Koic, N.R.Bertoniere, R.L.Smith, and G.W.Griffin, J. Am.Chem.Soc., 1970 92, 1302.
- 14 "Organic Synthesis", N.Rabjohn, Ed.-in-Chief, Coll.Vol.IV, John Wiley and Sons, Inc., New York, N.Y., 1963, p 771.
- 15 J.G.Calvert and J.N.Pitts, "Photochemistry", John Wiley and Sons, Inc., New York, N.Y., 1966, pp 379-385.

16 N .R.Bertoniere and G.W.Griffin, chap. 2 in "Organic Photochemistry", III, O.L.Chapman, Ed., Marcel Dekker, Inc., New York, N.Y., 1973, pp 115-195.

17 (a) H.Kristinsson and G.W.Griffin, J.Am.Chem.Soc., 1966, 88, 378; idem., Tetrahedron Lett., 1966, 3259; E.W.Valyocsik and P.Sigal, J.Org.Chem., 1971 36, 66; K.Salisburg, J.Am.Chem.Soc., 1972, 94, 3707; P.H.Mazzocchi and K.S. Lustig, ibid., 1975, 97, 3707; ibid., 1975, 97, 3714; (b) R.P.Story, Tetrahedron Lett., 1962, 414; C.Walling and A.A.Zavitsas, J.Am.Chem.Soc., 1963, 85, 2084.

18 K.N.Houk, J.Sims, R.E.Duke, Jr., R.W.Strozier, and J.K.George, J.Am.Chem.Soc., 1973, 95, 7287; K.N.Houk, J.Sims, C.R.Watts, and L.J.Luskus, ibid., 1973, 95, 7301.

Received, 11th June, 1977