

PO 5

SYNTHESES IN THE FURAN SERIES

J. Šrogl, M. Jonda, I. Stibor

Department of Organic Chemistry, Prague Institute of Chemical Technology, 166 28 Prague 6, Czechoslovakia

PO 6

STUDIES IN PYRROLE SERIES

M. Jonda, J. Šrogl, P. Holý and P. Duchek,

Department of Organic Chemistry, Institute of Chemical Technology, Prague

The investigation of anodic behaviour of pyrroles bears a specific feature of this type of heterocycle. In contrary to furans and thiophenes a four-electron oxidation takes place in case of 1-methylpyrrole leading to 5,5-dimethoxy-1-methyl-pyrrolin-2-on as the main product.

The oxidation has been carried out in a two- or three-electrode system using platinum anode in methanolic solutions of potassium hydroxide or tetramethylammonium perchlorate.

Under similar conditions the electrochemical oxidation of some pyrrole derivatives has been studied. Thus, 1,2,4-trimethylpyrrole and 1,2,5-trimethylpyrrole, respectively, afforded products of oxidation attack on C-methyl groups, i.e. the corresponding methoxymethyl derivatives. Without obligation to protect the nitrogen atom an analogous course of reaction has been observed in case of diethyl 3,5-dimethyl-2,4-pyrrole dicarboxylic acid.

Using ammonium bromide as electrolyte a mixture of products has been observed in oxidation of 1-methylindole among them 3,3-dibromo-1-methyl-2-indalinon, 3,3,5-tribromo-1-methyl-2-indolinon and 5-bromoindatin have been identified.

The reaction mechanism is discussed.

PO 7


MIXED HYDRIDE HYDROGENOLYSIS OF AROMATIC ALIPHATIC ACETALS. THE REACTION CONTROLLING STEP

J. Dušek and J. Jonas*


Department of Organic Chemistry, Faculty of Sciences, J. E. Purkyně University, Kotlářská 2, 611 37 Brno, Czechoslovakia

Generally, mixed hydride hydrogenolysis of acetals is supposed to proceed via the following steps:

- 1) formation of a complex between the acetal and the hydride,
- 2) formation of a stable oxycarbenium ion,
- 3) rapid hydride ion transfer to the oxycarbenium ion



With R_1 and R_2 = alkyl, the reaction controlling step (RCS) is supposed to be the step 2). However, there have been discussions concerning the RCS in cases where R_1 or R_2 = aryl, and step 1) has been repeatedly suggested as the RCS. We have tried to clarify this situation by preparing mixed aromatic aliphatic acetals of the types (I), (II), (III), and (IV), measuring their k_{rel} in the reaction with chloroalane, and using the data obtained in LFER correlations (which should give reaction constants ρ with opposite signs for RCS being 1) and RCS being 2)).



where for all I–IV, X = H, CH₃, OCH₃, Cl, and *tert*-C₄H₉. The acetals I, II, III, and IV were prepared from *iso*-butyl vinyl ether, cyclohexyl vinyl ether, 2,3-dihydrofuran, 2,3-dihydropyran, and the corresponding *p*-substituted phenols, respectively. The relative rate constants k_{rel} were measured using both HPLC and ¹H NMR.

The correlations obtained were all satisfactory and gave positive ρ s for reactions of acetals I, II, and III, and a negative ρ for the reaction of IV. The obtained ρ s differ considerably in their absolute values, too.

The significance of both differences is discussed.

PO 8

CYCLISATION OF 2-NITROARYLHYDRAZO COMPOUNDS TO 2-PHENYLBENZOTRIAZOLE-1-OXIDES


M. Potáček*, K. Picka, and J. Švaříček

Department of Organic Chemistry, Faculty of Sciences, J. E. Purkyně University, Brno

Research Institute of Pure Chemicals, Lachema, Brno

The aim of our study was elucidation of the pathway by which 2-nitroazocompounds are reductively converted into the corresponding 2-phenylbenzotriazoles.

We have shown earlier¹ that 2-nitrohydrazocompounds and 2-phenylbenzotriazole-1-oxides are intermediates in this reaction (see Scheme 1).



Scheme 1