

PO 20

SYNTHETIC APPLICATION OF PYRIDINIUM ARYLSULFONYLMETHYLIDES

R. A. Abramovitch

Department of Chemistry, Clemson University, Clemson S.C. 29631 USA

E. Floch*

Department of Organic Chemistry, Faculty of Chemical Technology, Slovak Technical University, 880 37 Bratislava, CSSR


The 1,3 dipolar cycloaddition of pyridinium arylsulfonylmethylides (1) with dimethyl acetylenedicarboxylate has been described recently¹ and led the formation of 1,2-dimethoxycarbonylindolizines (2). Reaction with methyl propiolate gave 1-methoxy carbonylindolizines. 1-Pyridinium arylsulfonylmethylides were generated by deprotonation of the corresponding pyridinium salts. Pyridinium p-toluenesulfonylmethylides also reacted with maleic anhydride in the presence of alcohols to give indolizine-2-carboxylates in a process involving selective decarboxylation and aromatization, and with phenylcyanooctene to give 1-cyano-2-phenylindolizines.²

Rather than expected indolizine derivatives as products of the 1,3 dipolar cycloaddition, reaction of ylide (1), (R = H with N-substituted maleimides in the presence of nucleophiles R'XH gave³ compounds (3) involving transfer of a carbon atom from (1) to the maleimide.

The preparation and the structure of compounds (3) will be discussed in light of spectral measurements and of the physicochemical properties of these compounds.

REFERENCES

- 1) R. A. Abramovitch and V. Alexanian, *J. Org. Chem.*, **41**, 2144 (1976)
- 2) R. A. Abramovitch and S. S. Mathur, *Heterocycles*, **5**, 91 (1976)
- 3) R. A. Abramovitch and D. P. Vanderpool, unpublished results.




PO 21

SYNTHESIS OF 1H, 9H-PYRULO[2',3':4,5]FURO[3,2-b]INDOLE DERIVATIVES

A. Kratoščíková*, J. Kováč, M. Dandárová

Department of Organic Chemistry, Slovak Technical University, 880 37 Bratislava

By condensation of 5-(2-nitrophenyl)-2-furaldehyde (I) with ethylester azidoacetic acid was obtained ethyl 3-[5-(2-nitrophenyl)-2-furyl]-2-azidoacrylate (II) which by a thermal ring closure gave ethyl 2-(2-nitrophenyl)-4H-furo-[3,2-b]pyrrole-5-carboxylate (III). Triethyl phosphite deoxygenation of the compound III rendered a derivative of a new heterocyclic system: 1H, 9H-pyrulo[2',3':4,5]furo[3,2-b]-indole (IV).




Other convenient way for obtaining of the compound IV is discussed also spectral data of the synthesised compounds are interpreted.

PO 22

SYNTHESIS AND TAUTOMERIC EQUILIBRIA OF 2-METHYLENE-3-OXO-1,2,3,4-TETRAHYDROQUINOXALINE DERIVATIVES

Jaromír Toman, Jiří Kličnar and Vladimír Macháček
Department of Organic Chemistry, University of Chemical Technology, 532 10 Pardubice — Czechoslovakia

It was found¹ ketimine-enamine tautomeric isomerisation (1) of 2-ethoxycarbonylmethylene-3-oxo-1,2,3,4-tetrahydroquinoxaline (I, R¹ = CO₂C₂H₅, R² = H) is catalysed with acids. In this paper the measurements of tautomeric equilibria by means of ¹H NMR (2H₂O-DMSO, 30–120 °C) was extended to further derivatives of the mentioned ester with substituents in the benzene ring (CH₃, Cl, NO₂, OCH₃) and to the derivatives with another groups activating the tautomeric isomerisation (CO, CN, NO₂).



The individual 6- and 7-substituted esters I were prepared² using the synthesis via appropriate oxides II (R² = 6- or 7-CH₃, Cl, NO₂, OCH₃), whereas the described³ 6-chloro- and 6-nitro-derivatives I (R¹ = CO₂C₂H₅, R² = 6-Cl or 6-NO₂) were found to be mixtures of both 6- and 7-isomers, their relative amounts were estimated. The cyanoderivative (I, R¹ = CN, R² = H) was obtained using a new synthesis starting from 1,2-diaminobenzene and ethyl isoxazole-5-carboxylate as potential α-dicarbonyl compound.

