

PO 20

SYNTHETIC APPLICATION OF PYRIDINIUM ARYLSULFONYLMETHYLIDES

R. A. Abramovitch

Department of Chemistry, Clemson University, Clemson S.C. 29631 USA

E. Floch*

Department of Organic Chemistry, Faculty of Chemical Technology, Slovak Technical University, 880 37 Bratislava, CSSR


The 1,3 dipolar cycloaddition of pyridinium arylsulfonylmethylides (1) with dimethyl acetylenedicarboxylate has been described recently¹ and led the formation of 1,2-dimethoxycarbonylindolizines (2). Reaction with methyl propiolate gave 1-methoxy carbonylindolizines. 1-Pyridinium arylsulfonylmethylides were generated by deprotonation of the corresponding pyridinium salts. Pyridinium p-toluenesulfonylmethylides also reacted with maleic anhydride in the presence of alcohols to give indolizine-2-carboxylates in a process involving selective decarboxylation and aromatization, and with phenylcyanooctene to give 1-cyano-2-phenylindolizines.²

Rather than expected indolizine derivatives as products of the 1,3 dipolar cycloaddition, reaction of ylide (1), (R = H with N-substituted maleimides in the presence of nucleophiles R'XH gave³ compounds (3) involving transfer of a carbon atom from (1) to the maleimide.

The preparation and the structure of compounds (3) will be discussed in light of spectral measurements and of the physicochemical properties of these compounds.

REFERENCES

- 1) R. A. Abramovitch and V. Alexanian, *J. Org. Chem.*, **41**, 2144 (1976)
- 2) R. A. Abramovitch and S. S. Mathur, *Heterocycles*, **5**, 91 (1976)
- 3) R. A. Abramovitch and D. P. Vanderpool, unpublished results.




PO 21

SYNTHESIS OF 1H, 9H-PYRULO[2',3':4,5]FURO[3,2-b]INDOLE DERIVATIVES

A. Kratoščíková*, J. Kováč, M. Dandárová

Department of Organic Chemistry, Slovak Technical University, 880 37 Bratislava

By condensation of 5-(2-nitrophenyl)-2-furaldehyde (I) with ethylester azidoacetic acid was obtained ethyl 3-[5-(2-nitrophenyl)-2-furyl]-2-azidoacrylate (II) which by a thermal ring closure gave ethyl 2-(2-nitrophenyl)-4H-furo-[3,2-b]pyrrole-5-carboxylate (III). Triethyl phosphite deoxygenation of the compound III rendered a derivative of a new heterocyclic system: 1H, 9H-pyrulo[2',3':4,5]furo[3,2-b]-indole (IV).




Other convenient way for obtaining of the compound IV is discussed also spectral data of the synthesised compounds are interpreted.

PO 22


SYNTHESIS AND TAUTOMERIC EQUILIBRIA OF 2-METHYLENE-3-OXO-1,2,3,4-TETRAHYDROQUINOXALINE DERIVATIVES

Jaromír Toman, Jiří Kličnar and Vladimír Macháček
Department of Organic Chemistry, University of Chemical Technology, 532 10 Pardubice — Czechoslovakia

It was found¹ ketimine-enamine tautomeric isomerisation (1) of 2-ethoxycarbonylmethylene-3-oxo-1,2,3,4-tetrahydroquinoxaline (I, R¹ = CO₂C₂H₅, R² = H) is catalysed with acids. In this paper the measurements of tautomeric equilibria by means of ¹H NMR (2H₂O-DMSO, 30–120 °C) was extended to further derivatives of the mentioned ester with substituents in the benzene ring (CH₃, Cl, NO₂, OCH₃) and to the derivatives with another groups activating the tautomeric isomerisation (CO, CN, NO₂).



The individual 6- and 7-substituted esters I were prepared² using the synthesis via appropriate oxides II (R² = 6- or 7-CH₃, Cl, NO₂, OCH₃), whereas the described³ 6-chloro- and 6-nitro-derivatives I (R¹ = CO₂C₂H₅, R² = 6-Cl or 6-NO₂) were found to be mixtures of both 6- and 7-isomers, their relative amounts were estimated. The cyanoderivative (I, R¹ = CN, R² = H) was obtained using a new synthesis starting from 1,2-diaminobenzene and ethyl isoxazole-5-carboxylate as potential α-dicarbonyl compound.



The 1,4-dimethyl derivative III ($R^1 = \text{CO}_2\text{Et}$, CN) with fixed enamine structure were prepared and their electronic spectra were discussed in comparison with those of parent derivatives I without the methyl groups.

The action of phosphorus pentasulfide on 4-methyl derivative of ester I ($R^1 = \text{CO}_2\text{C}_2\text{H}_5$, $R^2 = \text{H}$) in chlorobenzene gave 3-thion derivative IV ($R = \text{CO}_2\text{C}_2\text{H}_5$) with the ketimine structure. The thionation in pyridine solution afforded 1,3-dimethyl-1,2-dihydroquinoline-2-thione (IV, $R = \text{H}$).

From the values of relative amounts of both tautomers it follows, that electron-withdrawing substituents decrease the value $K_T = \text{ketimine/enamine}$, the electron donors having destabilisation effect on enamine. The effect of groups activating the tautomeric isomerisation and the effect of intramolecular H-bonding are discussed.

LITERATURE:

1. Macháček V., Toman J., Kličnar J.: Coll. Czech. Chem. Commun. — in press.
2. Toman J., Kličnar J., Macháček V.: Coll. Czech. Chem. Commun. — in press.
3. Iwanami Y.: Nippon Kagaku Zosshi 83, 316 (1962); Chem. Abstr. 59, 3919 (1963).

PO 23

SYNTHESIS OF SOME 2-ARYL-2,3-DIHYDRO-1,2,4-TRIAZINO [6,5-b]-INDOL-3-ONES

Jan Slouka, Vojtěch Bekárek and Václav Štemberk

Faculty of Sciences, Palacký University, Olomouc, Czechoslovakia

Coupling of diazonium salts with ethyl 3-indolylcarbamate gives high yields of the corresponding 2-aryloxy-3-ethoxy carbonylaminindoles, which undergo easily thermal cyclization to give the respective 2-aryl-2,3-dihydro-9H-1,2,4-triazino [6,5-b] indol-3-ones or the corresponding 4H-tautomers. The starting carbamate has been prepared by the Curtius rearrangement of 3-indolecarboxylic acid azide. Structure of the prepared 2-aryloxy-3-ethoxycarbonylaminindoles and of 2-aryl-2,3-dihydro-9H-1,2,4-triazino [6,5-b] indol-3-ones were studied by means of IR and ^1H -NMR spectroscopy with the use of ^{15}N -labeled derivatives.

PO 24


1,3-DIPOLAR CYCLOADDITION REACTION OF C-BENZOYL-N-PHENYLNITRONE WITH FURAN AND ITS DERIVATIVES

L. Fišera and J. Kováč

Department of Organic Chemistry, Slovak Institute of Technology, 880 37 Bratislava, Jánová 1, Czechoslovakia

A frontier orbital treatment (FMO) of furan suggests its possible reactivity in 1,3-dipolar cycloadditions with dipoles possessing low lying LUMO's, such as nitrones, especially those with electronwithdrawing substituents.

In this lecture is reported a detailed study of the cycloaddition of C-benzoyl-N-phenylnitron with furan, with particular attention directed at the regiochemistry of the reaction and to the detection of substitution products. On performing the reaction following products were obtained:



The structures were assigned on the basis of chemical and NMR evidence. The cycloaddition with 2-methylfuran, 2-methoxycarbonylfuran, 2-furancarbaldehyde and cycloadditions of some other nitrones to furan are also described.

PO 25


SYNTHESIS AND ANTITUBERCULOTIC ACTIVITY OF S-ALKYL 2-(6-X-BENZOTHIASOZOLYL)-DITHIOCARBAMATES

A. Gvozdková*, Z. Odlerová*, T. Gögh*

*Department of Organic Chemistry, Faculty of Science, Komenský University, 816 31 Bratislava


*Research Institute of Epidemiology and Microbiology, Department of Tuberculosis, 881 27 Bratislava

The relation between structure and tuberculostatic activity of S-Alkyl 2-(6-X-benzothiazolyl)dithiocarbamates




against M. tuberculosis H₃₇Rv and M. tuberculosis INH resistant has been studied

It has been proved that biological activity of dithiocarbamic groups isn't influenced by the structure of used amine. This fact has been also demonstrated on 2-(6-X-benzothiazolyl)thiuram disulphides



It has been stated, that tuberculostatic activity is increased by the substituents in position 6 in this order:



In this connection the influence of length of alkyl R has been studied. The most effective have been found the alkyls $\text{C}_2 - \text{C}_9$.