

REACTIONS OF DI-SUBSTITUTED CYCLOIMMONIUM YLIDES WITH CYCLOPROPENES

Kiyoshi Matsumoto*, Yukio Ikemi-Kono*, and Takane Uchida**

*College of Liberal Arts and Science, Kyoto University, Sakyo-ku,
Kyoto 606, Japan

**Faculty of Education, Fukui University, Bunkyo, Fukui 910, Japan


As a part of our studies on the reactions of strained cyclic olefines, reactions of pyridinium dicyanomethylides and dialkoxycarbonylmethylides with 1,2,3-triphenylcyclopropene (TPP) and diphenylcyclopropenethione (DPT) were examined.

Pyridinium and mono- or di-substituted-pyridinium dicyanomethylides undergo 1,3-dipolar cycloadditions to TPP in refluxing DMF producing the corresponding 1,2,3-triphenylindolizines (I) and 5-cyano-6,7,8- or 4-cyano-1,2,3-triphenyl-4H-quinolizines (II or III). The formation of 4H-quinolizines may predominate depending on the structures of ylides and the 1:1 adducts (IV) are also obtained in some cases.


The similar reactions of the dialkoxycarbonylmethylides with TPP give the indolizines and/or the 1:1 adducts only in poor yield.

On the other hand, with DPT the pyridinium dicyano-, dimethoxycarbonyl-, diethoxycarbonyl-, and methoxycarbonylcyanomethylides react in acetonitrile at room temperature as a nucleophile to afford 7-azaindenethione (V), 7-oxaindenethione (VI), 2-pyranethione (VII), 2-thiopyranone (VIII), or 2-pyranone derivatives (IX).


The structures of the products and the possible mechanisms for thier formations are discussed.



(I)




(II)




(III)




(IV)




(V)




(VI)



(VII)



(VIII)



(IX)

