

SEPTENTRIONINE AND SEPTENTRIODINE: TWO NEW C<sub>19</sub>-DITERPENOID  
ALKALOIDS FROM ACONITUM SEPTENTRIONALE KOELLE

S. William Pelletier\* and Rajinder S. Sawhney  
Institute for Natural Products Research and the Department of Chemistry,  
University of Georgia, Athens, Georgia 30602, U. S. A.

Arne Jørgen Aasen  
Department of Pharmacy, University of Oslo  
Oslo 3, Norway


Investigation of the alkaloids of Aconitum septentrionale has led to the isolation and structure elucidation of two new diterpenoid alkaloids, septentrionine (1) and septentriodine (2). These are the first reported examples of C<sub>19</sub>-diterpenoid alkaloids bearing a carboxy ester group on the amide side chain. <sup>13</sup>C nmr chemical shifts and assignments for these and related alkaloids are presented.

In 1957 Marion et al<sup>1</sup> reported the isolation and characterization of seven alkaloids from the roots of Aconitum septentrionale Koelle. Two alkaloids, lappaconitine and deacetylappaconitine, were identified, but the structures of five other alkaloids were not elucidated. We have reinvestigated this plant (collected in Norway) and now report the complete structures of two new alkaloids, septentrionine (1) and septentriodine (2).

Septentrionine, C<sub>38</sub>H<sub>54</sub>N<sub>2</sub>O<sub>11</sub>, mp 123-125° (corr.), [α]<sub>D</sub><sup>26</sup> + 21.2° (c 1.0, CHCl<sub>3</sub>), was isolated from the 85% ethanol extract of the root by a combination of gradient pH separation, thick-layer, and column chromatographic techniques. The alkaloid showed ir absorption at 3540 (OH), 3275 (NH), 1750, 1708, 1688 (carbonyls) and 1608, 1590 (aromatic) cm<sup>-1</sup>. The 100 MHz <sup>1</sup>H nmr spectrum of septentrionine exhibited absorption characteristic of N-CH<sub>2</sub>-CH<sub>3</sub> (3H, t, J=6.8Hz) centered at δ 1.08; -CO-CH<sub>2</sub>-CH<sub>2</sub>-COOCH<sub>3</sub> (4H, s) at δ 2.80; and six methoxyl groups at δ 3.25 (3H), 3.37 (6H), 3.47 (6H), 3.70 (3H). A group of signals appearing in the region δ 7.04-8.74 (two d, two t) ppm are indicative of the aromatic protons of the anthranoyl group.<sup>2</sup> In the downfield region, the spectrum also showed a signal at δ 11.15 attributable to a secondary amide proton.

The <sup>13</sup>C nmr spectrum showed 38 signals corresponding to 38 carbon atoms in the molecule. These signals have been assigned on the basis of noise-decoupled spectra, direct analysis of non-protonated carbons, and by comparison with <sup>13</sup>C nmr spectra of other related C<sub>19</sub>-diterpenoid alkaloids.<sup>3</sup> The <sup>13</sup>C nmr spectrum of septentrionine indicates the presence of a lycocotnine-type skeleton. The pattern of chemical shifts of the aromatic

carbons in septentrionine is similar to that of ajacine (3) and delsemine (4), indicating that the alkaloid has a straight chain rather than an imide moiety.<sup>3</sup> The singlets at 90.4 and 80.9 ppm (Table 1) clearly demonstrate the presence of C(7)-OH and C(8)-OCH<sub>3</sub> substitution in septentrionine.<sup>4</sup>



- |   |                                                                                                                                                          |   |                                                                                           |
|---|----------------------------------------------------------------------------------------------------------------------------------------------------------|---|-------------------------------------------------------------------------------------------|
| 1 | R <sup>1</sup> = CH <sub>3</sub> ; R <sup>2</sup> = -(CH <sub>2</sub> ) <sub>2</sub> - COOCH <sub>3</sub>                                                | 5 | R <sup>1</sup> = R <sup>3</sup> = CH <sub>3</sub> ; R <sup>2</sup> = H                    |
| 2 | R <sup>1</sup> = H; R <sup>2</sup> = -(CH <sub>2</sub> ) <sub>2</sub> - COOCH <sub>3</sub>                                                               | 6 | R <sup>1</sup> = R <sup>2</sup> = CH <sub>3</sub> ; R <sup>3</sup> = COCH <sub>3</sub>    |
| 3 | R <sup>1</sup> = H; R <sup>2</sup> = CH <sub>3</sub>                                                                                                     | 7 | R <sup>1</sup> = R <sup>3</sup> = CH <sub>3</sub> ; R <sup>2</sup> = COCH <sub>3</sub>    |
| 4 | R <sup>1</sup> = H; R <sup>2</sup> = mixture of:                                                                                                         | 8 | R <sup>1</sup> = H; R <sup>2</sup> = COCH <sub>3</sub> ; R <sup>3</sup> = CH <sub>3</sub> |
|   | $\begin{array}{c} \text{CH}_3 \\   \\ -\text{CH}-\text{CH}_2-\text{CONH}_2 \\   \\ -\text{CH}_2-\text{CH}-\text{CONH}_2 \\   \\ \text{CH}_3 \end{array}$ | 9 | R <sup>1</sup> = R <sup>2</sup> = H; R <sup>3</sup> = CH <sub>3</sub>                     |

Hydrolysis (5% methanolic KOH, 25°C) of septentrionine afforded a new alkamine (5) and an amino acid, C<sub>11</sub>H<sub>11</sub>NO<sub>5</sub>, mp 183-184°. The <sup>1</sup>H nmr spectrum of the alkamine shows the presence of N-CH<sub>2</sub>-CH<sub>3</sub> (3H, t, J=7Hz) centered at δ 1.05 and five aliphatic methoxyl groups at δ 3.25 (3H), 3.37 (6H) and 3.47 (6H).


Obviously, the methoxyl group which is lost during hydrolysis of septentrionine must be present in the amino acid moiety. The <sup>13</sup>C nmr spectrum of the alkamine shows 26 carbons which are interpretable on the basis of structure (5). This structure (5) was confirmed by comparison of its <sup>13</sup>C nmr spectrum with that of ambigua (6), a new alkaloid recently isolated from *Consolida ambigua*.<sup>4</sup> The alkamine (5) was relatively unstable and decomposed when attempts were made to crystallize it. On acetylation with Ac<sub>2</sub>O/Py, 5 gave a monoacetate (7) whose <sup>13</sup>C nmr spectrum was similar to that of 5 except for some differences in the chemical shifts of C(4) and C(18) (Table 1). The additional signals at 171.5 and 21.0 ppm in compound 7 must be assigned to the acetyl group.

The amino acid, mp 183-184°, was identified as N-succinoylanthranilic acid by comparison of the ir and <sup>13</sup>C nmr spectra with those of an authentic sample prepared from anthranilic acid and succinic anhydride.<sup>5</sup>

On the basis of the above data, septentrionine is assigned structure 1, and is probably identical with alkaloid "A" (mp 123-129°) of Marion *et al.*<sup>6</sup>

Table 1. Carbon-13 Chemical Shifts and Assignments for  
Septentrionine (1), Septentriodine (2) and Related Alkaloids<sup>a</sup>

Carbon <sup>c</sup>	1	5	7	6	2	8
1	83.1	83.2	82.9 <sup>b</sup>	83.4	84.0	84.0
2	25.6	25.5	25.3	25.3	26.1	26.1
3	31.9	31.2	30.9	31.5	31.6	31.9
4	37.7	38.7	37.2	38.3	37.6	37.2
5	40.6	40.8	40.2	40.9	43.3	43.3
6	91.5	91.4	91.3	91.1	91.1	90.9
7	90.4	89.7	90.1	90.0	88.7	88.5
8	80.9	80.6	81.0	80.4	77.6	77.5
9	51.9	52.1	51.9	52.0	50.4	50.4
10	37.7	37.9	37.9	35.9	38.1	38.1
11	47.6	47.2	47.7	47.2	49.1	49.0
12	27.9	27.9	28.1	27.4	28.7	28.7
13	46.7	46.7	46.5	46.1	46.1	46.1
14	83.5	83.6	83.6 <sup>b</sup>	75.0	84.0	84.0
15	27.9	27.9	28.1	28.5	33.7	33.7
16	82.8	82.9	82.9	81.9	82.7	82.6
17	66.2	66.9	66.7	66.7	64.6	64.6
18	70.6	68.8	69.7	79.3	69.9	69.1
19	53.2	53.5	53.5	53.6	52.4	52.4
N-CH <sub>2</sub>	51.9	52.2	51.9	52.6	51.0	51.0
CH <sub>3</sub>	14.8	15.0	14.5	14.9	14.1	14.1
1'	55.7	55.6	55.7	55.6	55.9	55.7
6'	60.0	59.8	60.0	59.4	57.9	57.8
8'	54.4	53.6	53.5	53.7	-	-
14'	57.7	57.7	57.7	-	58.1	58.0
16'	56.5	56.5	56.6	56.3	56.4	56.3

Carbon	1	7	6	2	8
	168.4	(172.6) <sup>d</sup>		168.3	
1	115.1	(117.2)		114.7	
2	141.8	(142.3)		141.9	
3	120.6	(121.3)		120.8	
4	134.9	(135.1)		135.2	
5	122.7	(123.8)		122.8	
6	130.8	(132.4)		130.5	
R = NH-CO	170.6	(171.3)		170.6	
CH <sub>2</sub>	29.0	(29.8)		28.9	
CH <sub>2</sub>	32.7	(33.5)		32.7	
CO	173.3	(176.1)		173.3	
OCH <sub>3</sub>	51.9			51.9	
-CO		171.5	171.3		170.9
CH <sub>3</sub>		21.0	21.4		20.8

<sup>a</sup> In ppm downfield to Me<sub>4</sub>Si; solvent deuteriochloroform.

<sup>b</sup> These assignments may be interchanged in any vertical column.

<sup>c</sup> Primed numbers correspond to CH<sub>3</sub>O- groups attached to non-primed C-atoms.

<sup>d</sup> Values in parentheses are those of N-succinylanthranilic acid.

The second alkaloid, septentriodine (2),  $C_{37}H_{52}N_2O_{11}$ ,  $[\alpha]_D^{25} + 36.2^\circ$  (c, 1.0,  $CHCl_3$ ) could not be induced to crystallize. The 100 MHz  $^1H$  nmr spectrum of septentriodine is similar to that of septentrionine except for the absence of one methoxyl absorption. It reveals the presence of  $N-CH_2-CH_3$  (3H, t, J 7.1 Hz) centered at  $\delta$  1.07;  $-COCH_2-CH_2-COOCH_3$  (4H, s) at  $\delta$  2.80; five aliphatic methoxyl groups at  $\delta$  3.26, 3.34, 3.38, 3.42 and 3.71. The presence of four aromatic protons is shown by a group of signals in the region  $\delta$  7.0-8.7. The secondary amide ( $NH-CO$ ) proton was located at  $\delta$  11.10. The alkaloid showed ir bands at 3470 ( $-OH$ ), 3310 ( $NH$ ), 1740, 1705, 1687 (carbonyls), 1605, 1590 (aromatic) and 1085 ( $C-O$ )  $cm^{-1}$ . The base peak at  $m/e$  669 ( $M^+ - 31$ ) in the mass spectrum of septentriodine confirms the presence of a methoxyl group at C(1).<sup>7</sup>

The  $^{13}C$  nmr spectrum of septentriodine resembles that of septentrionine. The singlets at 88.7 and 77.6 ppm (Table 1) show the presence of a hydroxy group at C(8) in septentriodine. The chemical shifts of the alkamine part of septentriodine are very close to those of the corresponding carbons of tricornine (8). The structure (2) for septentriodine was confirmed by its hydrolysis (5% methanolic KOH) to lycoctonine (9) and N-succinoylanthranilic acid.

Septentrionine and septentriodine represent the first examples of  $C_{19}$ -diterpenoid alkaloids bearing a carboxy ester group in the amide side chain.

**Acknowledgment:** We thank Mr. Howard C. Higman, Tobacco Research Laboratory, Richard B. Russell Center, for the mass spectrum of septentriodine, Dr. Naresh V. Mody for the  $^{13}C$  nmr spectra, and Mr. Courtney Pape for the 100 MHz  $^1H$  nmr spectra.

#### REFERENCES AND NOTES

1. Léo Marion, L. Fonzes, C. K. Wilkins, Jr., J. P. Boca, F. Sandberg, R. Thorsen and E. Lindén, Can. J. Chem. **45**, 969 (1967).
2. S. W. Palletier and R. S. Sawhney, Heterocycles, **9**, 463 (1978).
3. S. W. Palletier, N. V. Mody, R. S. Sawhney and J. Bhattacharyya, Heterocycles, **7**, 327 (1977).
4. S. W. Palletier, R. S. Sawhney and N. V. Mody, Heterocycles, **9**, 1241 (1978).
5. K. Auwers and J. Harger, Ann. **292**, 187 (1896).
6. The molecular composition,  $C_{43}H_{59}N_3O_{12}$ , reported for alkaloid "A" is in error.
7. M. S. Yunusov, Ya. V. Rashkes, V. A. Tel'nov and S. Yu. Yunusov, Khim. prirod. Soedinenii, **5**, 515 (1969).

Received, 11th December, 1978