

$\gamma$ -BROMINATION OF QUINOLINE AND PYRIDINE N-OXIDES

Hirohisa Saito and Masatomo Hamana\*

Faculty of Pharmaceutical Sciences, Kyushu University

Maidashi 3-1-1, Higashi-ku, Fukuoka 812, Japan

Abstract—Treatment of quinoline 1-oxide (1) with bromine (2 equiv.) and thallium triacetate (3 equiv.) in acetic acid at 50° for 29 hr produces 4-bromoquinoline 1-oxide (2) in 65.8 % yield. Similarly, quinaldine and 3-bromoquinoline 1-oxides (5 and 12) give the corresponding  $\gamma$ -bromo derivatives (6 and 13) in good yields. From the reaction of 2-cyanoquinoline 1-oxide (8), 2-cyano-4-bromoquinoline 1-oxide (9) and 2-carbamoyl-4-bromoquinoline 1-oxide (10) are obtained. The reactivity of N-oxides of pyridine series is somewhat lower and pyridine and 2-picoline 1-oxides resist bromination under similar conditions, but 2,6-lutidine, 3-picoline and 3,5-lutidine 1-oxides (14, 16 and 18) afford the corresponding  $\gamma$ -bromo derivatives (15, 17 and 19).

Contrary to the facile  $\gamma$ -nitration of pyridine and quinoline 1-oxides, the corresponding  $\gamma$ -halogenation has long been rather difficult<sup>1,2</sup>. For instance, H. den Hertog and his co-workers obtained 2-bromo- and 4-bromo-pyridine 1-oxides in a total yield of 10 % on heating pyridine 1-oxide with excess bromine in 90 % sulfuric acid at 200° in the presence of silver sulfate<sup>3</sup>. Ochiai and Okamoto<sup>4</sup> have reported that 4-bromoquinoline 1-oxide is formed in 32 % yield when the perbromide of quinoline 1-oxide ( $C_9H_7ON \cdot 1/2HBr_3$ ) was shaken with bromine water in a sealed tube at room temperatures, however its reproducibility is rather poor for the synthetic procedure.

We now wish to report the successful  $\gamma$ -bromination of quinoline and pyridine 1-oxides under rather mild conditions by means of bromine and thallium triacetate. A solution of quinoline 1-oxide (1, 0.8 g), bromine (1 g, 1.1 equiv.) and thallium triacetate (2.3 g, 1.1 equiv.) in acetic acid (10 ml) was stirred at room tempera-

tures. While thallium triacetate dissolved in a little while, no reaction occurred and 1 was recovered almost quantitatively after 24 hr's reaction. However when the same reactants were warmed at 70° for 3 hr, 4-bromoquinoline 1-oxide (2), colorless needles, mp 127-128.5°, and quinoline (4) were obtained in 14.5 and 4.2 % yields, respectively, accompanied by recovery of 1 in 45 % (Exp. 3 in Table I). Representative results obtained under various conditions are given in Table I.

Table I. Bromination of Quinoline 1-oxide (1)

C1=CC=C2C(=C1)N(O)C=C2
 $\xrightarrow[\text{AcOH}]{\text{Br}_2, \text{Tl}(\text{OAc})_3}$ 
BrC1=CC=C2C(=C1)N(O)C=C2
+ 
BrC1=CC=C2C=CN1C=C2
+ 
C1=CC=C2C=CN1C=C2

1
2
3
4

Exp. NO.	Br <sub>2</sub> (equiv.)	Tl(OAc) <sub>3</sub> (equiv.)	Reaction		Product, Yield (%)			Recov. (%)
			Temp. (°C)	Time (hr)	<u>2</u>	<u>3</u>	<u>4</u>	
1	1.1	1.1	50	4	9.7	-	8.2	41.1
2	1.1	1.1	60	4	21.4	-	7.7	36.0
3	1.1	1.1	70	3	14.5	-	4.2	45.0
4	1.1	1.1	80	4	11.8	-	6.6	33.5
5	2.0	1.0	60	4	minute	4.8	4.4	
6	1.0	2.0	50	4	27.4	-	2.2	41.4
7	2.0	3.0	50	4	46.4	3.9	-	23.5
8	2.0	4.0	50	4	34.8	1.9	-	34.5
9	2.0	3.0	50	29	65.8	6.0	13.6	12.1

These results indicate that the reaction is fairly affected by the reaction temperature and the amount of reagents. While the reaction is very slow at room temperatures, warming at 50-60° substantially promotes the reaction to give 2 in 10-22 % yields even when molar equivalent amounts of bromine and thallium triacetate were used, but higher reaction temperatures (over 70°) conversely decrease the yield of 2. The use of two or three molar excess of the reagents was apparently favorable for the  $\gamma$ -bromination, and the best yield (65.8 %) was obtained from a run under the conditions of Exp. 9 shown in Table I. A small amount of 4-bromoquinoline (3) was also formed as a by-product in reactions using excess reagents. Subsequently, bromination of some derivatives of 1 was carried out (Chart 1).

While lepidine 1-oxide gave no bromo derivative, quinaldine 1-oxide (5) underwent  $\gamma$ -bromination more readily than 1 to afford 4-bromoquinaldine 1-oxide (6), pale yellow needles, mp 141-143°, and 4-bromoquinaldine (7), bp 130°/4 mm. (picrate, mp

246-247°) in 63 and 0.9 % yields, respectively. In view of its high susceptibility to bromination, it is very noteworthy that the active 2-methyl group was inert to bromination under the above conditions. The structures of 6 and 7 were confirmed by their independent syntheses from 4-nitroquinoline N-oxide<sup>5</sup>.




Chart 1.

The reaction of 2-cyanoquinoline 1-oxide (8) resulted in the formation of 2-cyano-4-bromoquinoline 1-oxide (9), pale yellow needles, mp 188-189°, 2-carbamoyl-4-bromoquinoline 1-oxide (10), colorless needles, mp 278-279°, and 2-carbamoylquinoline 1-oxide (11)<sup>6</sup>, colorless plates, mp 223°, in the respective yields of 22.5, 11.2 and 5.4 %. It was clarified by separate reactions that 8 was convertible into 11 by the action of thallium triacetate in acetic acid, and 10 originated only from 9 but not from 11. Bromination of 3-bromoquinoline 1-oxide (12) produced 3,4-dibromoquinoline 1-oxide (13), colorless needles, mp 146-147°, in a high yield of 95.7 %.

The reactivity of N-oxides of pyridine series was found to be considerably lower compared to that of quinoline 1-oxides. Thus any brominated products could not be isolated from reactions of pyridine and 2-picoline 1-oxides in spite of many at-

tempts under various conditions, the N-oxide being recovered in each case. However, 2,6-lutidine 1-oxide (14) afforded 4-bromo-2,6-lutidine 1-oxide (15), colorless needles, mp 67-68°, under the conditions shown in Chart 2. 3-Picoline 1-oxide (16) and 3,5-lutidine 1-oxide (18) were more reactive as expected, and the corresponding 4-bromo derivatives 17<sup>7</sup>, colorless pillars, mp 112-113° (59.2 %) and 19, colorless needles, mp 199° (72.8%), were obtained, respectively. The structures of 15, 17 and 19 were unambiguously established by elemental analyses, the mass and NMR spectrometry.




Chart 2.

The orienting effect of N-oxide function apparently operates not only in the formation of 15 but also in that of 17, because 3-picoline is known to give not 4-bromo derivative by the usual bromination procedure<sup>8</sup>. Although no information is available on bromination of 3,5-lutidine, it may be reasonably assumed that the directing effect of N-oxide function appears also in the formation of 19. Further studies are in progress to explore the scope of the reaction and elucidate the mechanism.

#### ACKNOWLEDGEMENT

We are grateful both to a Grant-in Aid for Chemical Research in Development and Utilization of Nitrogen-Organic Resources from the Ministry of Education, Science and Culture, Japan, and to Takeda Science Foundation for partial financial support of this work.

## REFERENCES

1. E. Ochiai, 'Aromatic Amine Oxides', Elsevier Publishing Co., Amsterdam, 1967, pp. 210 - 246.
2. A. R. Katritzky and J. M. Lagowski, 'Chemistry of the Heterocyclic N-Oxides', Academic Press, London and New York, 1971, pp. 231 - 258.
3. H. C. van der Plas, H. J. den Hertog, M. van Ammers, and B. Haase, Tetrahedron Letters, 1961, 32.
4. E. Ochiai and T. Okamoto, Yakugaku Zasshi, 1947, 67, 87.
5. E. Ochiai and K. Satake, Yakugaku Zasshi, 1951, 71, 1078.
6. T. Takahashi and Y. Hamada, Yakugaku Zasshi, 1955, 75, 1434.
7. I. Itai and H. Ogura, Yakugaku Zasshi, 1955, 75, 242 ; They reported that compound 17 melted at 72-73°. However, we confirmed that their sample contained a small amount of water and the pure crystals of 17 melted at 112-113°.
8. L. van der Does and H. J. den Hertog, Rec. Trav. Chim. Pays-Bas, 1965, 84, 951.

Received, 21st December, 1978