

SYNTHESIS OF SPIROFLUORENES via KETENE ADDUCTS

Magid A. Abou-Gharbia and Madeleine M. Joullie<sup>\*\*</sup>


Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104 U.S.A.

Abstract - New classes of spirofluorenes were prepared via simple transformations of ketene adducts.


We have recently reported the synthesis of spirofluorenes of biological interest by conventional synthetic routes.<sup>1,2</sup> As a continuation of our investigations of the biological activity of spirofluorenes, we now wish to report the synthesis of novel spirofluorenes using ketene adducts as precursors.

The cycloadditions of N-fluorenylideneanilines **1** with 2-oxathirane 1,1-dioxide **2** afforded spirothiazolidinones **3a-c** in 70-80% yields. The same compounds were also prepared by an alternate synthetic route involving the condensation of imines **1a-c** with mercaptoacetic acid and subsequent oxidation of the intermediate spirothiazolidinones (**4a-c**) with sodium perchlorate (Scheme 1).<sup>3,4</sup> The yields for both methods were comparable.

Scheme 1



Another class of spirofluorenes (**5** and **6**) were prepared via the simple and mild transformation of ketene adducts **3a** and **3c**.



Compound **5** was prepared by heating **3a** with a strong non-nucleophilic base such as sodium hydride in DMF at 100° (½ hr.), followed by alkylation with 1,2 dibromopropane. In the presence of an excess of sodium hydride, the generated tertiary anion displaces the remaining bromine of the substituted bromopropane to afford **5** (8 hr). Compound **6** was obtained by heating **3c** with p-N,N-dimethylaminonitrosobenzene and a catalytic amount of ammonium acetate in dry benzene and removing the water formed in the reaction by azeotropic distillation. Previous investigators had described the reactions of ketenes with N-benzylidenealkylamine or - arylamine N-oxides.<sup>5,6</sup> We have extended these investigations to N-fluorenylidenealkylamine or - arylamine N-oxides.<sup>7,8</sup> Although the reaction of diphenylketene and N-fluorenylidene-N-methylamine N-oxide was reported by Taylor et al.<sup>6</sup> to afford, among other products, spirooxazolidinone **7**, only the melting point (145°) of this compound was given. Since no spectral or analytical data was available at the time, we reinvestigated this reaction.<sup>9</sup> When this condensation was carried out in benzene,

Scheme 2




Table 1. Physical and Spectral Properties of Spirofluorenes.

Compound No.	Yield, %	mp, °C	Spectral Data
3a	81	260-262	IR $\nu_{\text{max}}^{\text{KBr}}$ 1680 (C=O), 1140 and 1335 $\text{cm}^{-1}$ ( $\text{SO}_2$ ); $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 5.0(s, 2H), 6.65-7.0(m, 5H), and 7.3-8.0(m, 8H); m/e 361 ( $\text{M}^+$ ).
3b	76	237-239	IR $\nu_{\text{max}}^{\text{KBr}}$ 1680 (C=O), 1130 and 1325 $\text{cm}^{-1}$ ( $\text{SO}_2$ ); $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 5.0(s, 2H) and 6.85-7.95(m, 12H); m/e 315 ( $\text{M}^+ - \text{SO}_2$ ).
3c	79	240-241	IR $\nu_{\text{max}}^{\text{KBr}}$ 1700 (C=O), 1140 and 1325 $\text{cm}^{-1}$ ( $\text{SO}_2$ ), $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 4.95(s, 2H), 6.70-7.1(m, 4H) and 7.20-7.9(m, 8H).
4a	71	225-227	IR $\nu_{\text{max}}^{\text{KBr}}$ 1650 (C=O), $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 4.25(s, 2H) and 6.55-7.8(m, 13H); $^{13}\text{CNMR}(\text{CDCl}_3)$ 78.27 ppm(C-9).
4b	82	130-132	IR $\nu_{\text{max}}^{\text{KBr}}$ 1650 (C=O), $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 4.20(s, 2H), and 6.75-7.95(m, 13H).
5	63	223-225	IR $\nu_{\text{max}}^{\text{KBr}}$ 1675 (C=O), 1115 and 1320 $\text{cm}^{-1}$ ( $\text{SO}_2$ ); $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 2.1-2.5 (m, 2H), 2.9-3.2(t, 4H), and 6.8-7.5(m, 13H); m/e 401 ( $\text{M}^+$ ).
6	82	138-140	IR $\nu_{\text{max}}^{\text{KBr}}$ 1680 (C=O), 1600 (C=N), 1120 and 1320 $\text{cm}^{-1}$ ( $\text{SO}_2$ ); $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 3.1(s, 6H), 6.6-6.8 (m, 4H), and 7.1-7.8(m, 12H).
7	72	148-150	IR $\nu_{\text{max}}^{\text{KBr}}$ 1775 $\text{cm}^{-1}$ (C=O), $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 2.15(s, 3H) and 7.1-7.6(m, 18H); $^{13}\text{CNMR}(\text{CDCl}_3)$ 102.7 ppm (spiro C-9); m/e 403 ( $\text{M}^+$ ).
9	59	237-239	IR $\nu_{\text{max}}^{\text{KBr}}$ 1775 and 1840 $\text{cm}^{-1}$ (C=O); $^1\text{HNMR}(\text{CDCl}_3)$ $\delta$ 2.25 (s, 3H), 3.41-4.0 (m, 2H), and 6.4-7.9 (m, 18H); m/e 457 ( $\text{M}^+$ ).

spirofluorene 7 was obtained as the sole product. Attempts to condense diphenylketene with N-fluorenylideneaniline N-oxide resulted in the formation of an unstable adduct which decomposed with gas evolution.

When 7 was heated in benzene, loss of carbon dioxide occurred to afford the highly colored nitrogen ylid (8) which was trapped with maleic anhydride to afford spirofluorene 9 (Scheme 2). Attempts to form a similar ylid from the adduct obtained from tert-butylcyanoketene and N-fluorenylidene-methylamine N-oxide were unsuccessful.

#### REFERENCES AND NOTES

1. M. A. Abou-Gharbia and M.M. Joullie, J. Pharm. Sci., 56, 1977, 1653.
2. M. A. Abou-Gharbia, T. T. Su and M. M. Joullie, J. Pharm. Sci., 67, 1978, 953.
3. A. Gomes and M. M. Joullie, J. Heterocycl. Chem., 6, 1969, 729.
4. Sodium permanganate afforded a poor yield of the sulfone and m-chloroperbenzoic acid gave no yield.
5. R. N. Pratt, D. P. Stokes, G. A. Taylor and P. C. Brookes, J. Chem. Soc. (C), 1968, 2086; D. P. Stokes and G. A. Taylor, ibid., 1971, 2334.
6. A. F. Gettins and G. A. Taylor, J. Chem. Soc. Chem. Comm., 1972, 1146.
7. M. A. Abou-Gharbia and M. M. Joullie, Synthesis, 5, 1977, 318.
8. M. A. Abou-Gharbia, I. Miura and M. M. Joullie, Heterocycles, 9, 1978, 457.
9. A subsequent publication, A. F. Gettins, D. P. Stokes, and G. A. Taylor, J. Chem. Soc. Perkin I, 1977, 1849, reports additional data for 7, however, the melting point given, 138° still disagrees with our value (148-150°) and their previous results. (145°).

Received, 5th April, 1979