

STUDIES ON HETEROCYCLIC ANALOGUES OF AZULENE. PART 5.<sup>1</sup>  
 REARRANGEMENT OF 7H-6a-AZACYCLOBUTA[j]CYCLOPENT[1,2,3-cd]AZULENE  
 RING SYSTEM ON SILICA GEL

Noritaka Abe and Tarozaemon Nishiwaki


Department of Chemistry, Faculty of Sciences, Yamaguchi University,  
 Yoshida, Yamaguchi 753, Japan

Abstract -- Rearrangement of 7H-6a-azacyclobuta[j]cyclopent[1,2,3-cd]-azulene ring system to 3H-2a-azacyclopenta[ef]heptalene ring system by contact with silica gel was reported.


It is known that contacts of some organic substances with adsorbents (i.e. alumina or silica gel) cause structural changes.<sup>2</sup> Sometimes silica gel displays as catalyst in the isomerization reactions of olefines<sup>3</sup> or bicyclic compounds.<sup>4</sup> We now report that 7H-6a-azacyclobuta[j]cyclopent[1,2,3-cd]azulene ring system rearranges to 3H-2a-azacyclopenta[ef]heptalene ring system on silica gel.

Contact of 5-ethyl 8,9-dimethyl 6-chloro-7-phenyl-7H-6a-azacyclobuta[j]cyclopent[1,2,3-cd]azulene-5,8,9-tricarboxylate (1a)<sup>5</sup> with silica gel<sup>6</sup> for 5 days at room temperature gave 1-ethyl 4,5-dimethyl 2-chloro-3-phenyl-3H-2a-azacyclopenta[ef]heptalene-1,4,5-tricarboxylate (2a)<sup>7,8</sup> as brown needles; mp 148-149° (decomp.); 56 %; IR (nujol) 1725, 1705, and 1700 cm<sup>-1</sup> (C=O). In <sup>1</sup>H nmr spectrum (CDCl<sub>3</sub>), two 1H singlets to be assignable to H-3 and H-6 are seen at δ 5.31 and 6.11, respectively. Another signals are seen at δ 1.36 (3H, t, J 7 Hz, CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 3.59 (3H, s, CO<sub>2</sub>CH<sub>3</sub>), 3.84 (3H, s, CO<sub>2</sub>CH<sub>3</sub>), 4.23 (2H, q, J 7 Hz, CO<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 5.95-6.2 (1H, m, H-9), 6.3-6.5 (2H, m, H-7,8), 7.30 (5H, m, phenyl), and 7.38 (1H, d, J 11 Hz, H-10). <sup>13</sup>C nmr spectrum (CDCl<sub>3</sub>) of 2a exhibits signal assignable to sp<sup>3</sup> carbon atom at δ 68.59 (d, C-3). The uv spectrum [ $\lambda_{\max}^{\text{EtOH}}$  228 (log ε 4.54), 260 (4.28), 434 (4.38), 492sh (3.95), 530 (3.74), and 570 nm (3.43)] resembles that of 3H-2a-azacyclopenta[ef]heptalene ring system.<sup>9</sup> From these results, we assigned the structure.<sup>10</sup>

In a similar manner, 1b gave 2b as brown needles, mp 116-118°, in 60 % yield.<sup>10</sup> On more active adsorbent (alumina), 1a underwent decomposition and gave no



1a: E=CO<sub>2</sub>Me, R=H  
 1b: E=CO<sub>2</sub>Me, R=Me




2a: E=CO<sub>2</sub>Me, R=H  
 2b: E=CO<sub>2</sub>Me, R=Me

obvious products.

Formation of the products (2) can be accommodated by a mechanism started by the scission of the bond between nitrogen and benzylic-carbon atoms of 1, followed by formation of 3 and intramolecular cyclization of 3 with association of silica gel.<sup>11</sup> It is considered that adsorption of 1 on silica gel plays important roles which activates the bond between nitrogen and benzylic-carbon atoms of 1 and arranges to the suitable orientation for rearrangement to 3.<sup>11</sup>

ACKNOWLEDGMENT We thank Dr. M. Yasunami (Tohoku Univ.) and Dr. A. Mori (Kyushu Univ.) for nmr spectral determination.



#### References and notes

- Part 4, T. Nishiwaki, N. Abe, K. Ishida, and K. Miura, Bull. Chem. Soc. Japan, 1978, 51, 667.
- L. Novotný and K. Kotva, Collection Czechoslov. Chem. Commun., 1974, 39, 2949; H. J. Seebald and W. Schunack, J. Chromatog., 1972, 74, 129; A. Szabó and E. Tömörkény, ibid., 1978, 151, 90.
- B. B. West, G. L. Haller, and R. L. Burwell, Jr., J. Catal., 1973, 29, 486 and references cited therein.
- I. I. Khochenko, E. M. Mil'vitskaya, A. F. Plate, and A. M. Pryanishnikova, Zh. Org. Khim., 1973, 9, 2307.
- N. Abe and T. Nishiwaki, J. C. S. Chem. Commun., in press.
- Merck Kieselgel 60 was used.
- 1 was stable at room temperature but gave another rearranged products at reflux in benzene or xylene. These results were delineated in a separate paper.<sup>5</sup>
- 2 was slightly unstable at room temperature in air or by prolonged contact with silica gel, and gave unidentified red substances.
- N. Abe, Y. Tanaka, and T. Nishiwaki, J. C. S. Perkin I., 1978, 429.
- Satisfactory elemental analyses and spectroscopical data other than partially given in this paper were obtained for all new compounds here described.
- According to reference 3, catalytic activity of the commercial gels may result from traces of transition metals, most probably iron.

Received, 26th May, 1979