


SYNTHESIS OF PIMPRININE AND RELATED OXAZOLYLINDOLE ALKALOIDS
FROM N-ACYL DERIVATIVES OF TRYPTAMINE AND TRYPTOPHAN METHYL ESTER
BY DDQ OXIDATION

Yuji Oikawa, Tadao Yoshioka, Kunihiko Mohri, and Osamu Yonemitsu*
Faculty of Pharmaceutical Sciences, Hokkaido University
Sapporo 060, Japan

Abstract — N-Acyl derivatives of tryptamine and L-tryptophan methyl ester were treated with DDQ under anhydrous conditions directly to give pimprinine and related oxazoly lindole alkaloids. Under aqueous conditions, the N-acyl derivatives were readily oxidized to the corresponding β -keto compounds, which were also converted to the oxazoly lindoles by the treatment with POCl_3 .

Pimprinine (3b), an alkaloid having antiepileptic¹ and monoamine oxidase inhibitory activities,² was first isolated from Streptomyces pimprina³ and synthesized by Joshi et al⁴ through the dehydrative cyclization of 1-acetyl-3-acetamidoacetyl-indole, though somewhat difficult to synthesize the starting material, 3-aminoacetylindole. Recently, Koyama et al isolated two very close alkaloids, pimprine (3c) and pimprinaphine (3h), as well as pimprinine (3b) from Streptoverticillium olivoreticuli and synthesized all the three compounds by an improved method via 3-(5-oxazoly)indole (3a).⁵

We recently reported that 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ) is the only reagent for the selective oxidation of side chains at C-3 of indoles.⁶ Almost at the same time, the biological oxidation of this type was interestingly found, that is, Hayaishi et al isolated a new enzyme, tryptophan side chain oxidase from Pseudomonas, which oxidizes tryptophan derivatives to their β -hydroxy, β -keto, and α, β -dehydro compounds.⁷ In the present paper, we wish to report an application of the DDQ oxidation for a biomimetic synthesis of pimprinine and a series of related compounds.⁸



When N-acetyltryptamine (1b) and DDQ (2 equiv) in THF was refluxed under argon for 50 min, pimprinine (3b) was formed directly and isolated quite easily, though in a very poor yield (10%), after passing through a short alumina column. The N-benzoyl compound (1e) similarly gave the corresponding oxazolyndole derivative (3e), but the yield was still unsatisfactory. Better results were obtained in the oxidation of N-acyl derivatives of tryptophan methyl ester (2), especially 2f having an electron-donative methoxybenzoyl group gave 4f in fairly good yield, whereas the nitro derivative (2g) gave again a poor result (Table I).

Table I. Oxazolyndoles (3, 4) from N-Acyl Derivatives of Tryptamine (1) and Tryptophan Methyl Ester (2) by DDQ Oxidation in Anhydrous THF.

<u>1</u> , <u>2</u>	R'	<u>3</u>	<u>4</u>
		Yield(%) mp°C	Yield(%) mp°C
<u>b</u>	CH ₃	10 201-202	36 239-240
<u>d</u>	CH(CH ₃) ₂		55 194-195
<u>e</u>	C ₆ H ₅	39 224-226	73 266-267
<u>f</u>	p-CH ₃ OC ₆ H ₄		79 245-246
<u>g</u>	p-NO ₂ C ₆ H ₄		22 296-298

When N-acyl tryptamines (1) were treated with DDQ (2 equiv) in THF-H₂O (9:1) under argon at room temperature for 1-2 hr, the corresponding β -keto products (5) were readily isolated in good yields regardless of the N-acyl group after passing through a short alumina column in EtOAc.⁹ Similarly, N-acyltryptophan methyl esters (2), except 2e and 2f, were converted to 6 (Table II) with the concomitant formation of slight amounts of oxazolyliindoles (4) having strong fluorescences, though only detected by tlc and uv spectra. Interestingly, 2e and 2f gave only the corresponding oxazolyliindoles (4e, 4f), not β -keto derivatives (6e, 6f), even in the aqueous solvents.¹¹ No trace of 4g, on the contrary, was detected in the oxidation of 2g having an electron-withdrawing nitrophenyl substituent.

Table II. β -Keto Derivatives (5, 6) from 1 and 2 by DDQ
Oxidation in THF-H₂O (9:1).

<u>1</u> , <u>2</u>	R'	<u>5</u> Yield(%) mp°C	<u>6</u> Yield(%) mp°C
<u>a</u>	H	86 194-195	69 214(dec)
<u>b</u>	CH ₃	73 216-218	84 202-205
<u>c</u>	CH ₂ CH ₃	89 218-220	79 145-146
<u>d</u>	CH(CH ₃) ₂	78 212-213	74 157-159
<u>e</u>	C ₆ H ₅	quant 213-214	72 ^a 266-267
<u>f</u>	p-CH ₃ OC ₆ H ₄	85 207-208	75 ^a 245-246
<u>g</u>	p-NO ₂ C ₆ H ₄	84 239-241	43 oil
<u>h</u>	CH ₂ C ₆ H ₅	quant 215-217	80 150-152
<u>i</u>	OCH ₂ C ₆ H ₅	86 230-232	71 oil

^a Yield of 4, not 6.

Since 3-acylamidoacetylindoles (5) are known to give oxazolyliindoles (3) by the dehydrative cyclization with POCl₃,^{4,5} 5b-f,h were subjected to this reaction and the results are shown in Table III. This indirect synthesis from 1 is obviously superior at least in yield to the direct synthesis by the oxidation under anhydrous conditions. Practically, 3 were synthesized from 1 as follows: 1 and DDQ (2 equiv) in THF-H₂O (9:1) under argon were allowed to stand at room temperature for 1-2 hr. After evaporation of the THF, the residues were treated with 0.1 N NaOH to recover 2,3-dichloro-5,6-dicyanohydroquinone (DDHQ) and the precipitated solids were collected by filtration and washed with H₂O and EtOAc (decolorization). The dried solids (5) were refluxed in POCl₃ under argon to give 3.


Table III. Oxazolyllindoles (3) from 5 by
Dehydrative Cyclization with POCl₃.

	R'	Yield(%)	mp°C
<u>b</u>	CH ₃	82	202-203
<u>c</u>	CH ₂ CH ₃	86	161-163
<u>d</u>	CH(CH ₃) ₂	85	135-136
<u>e</u>	C ₆ H ₅	80	225-226
<u>f</u>	p-CH ₃ OC ₆ H ₄	83	205-206
<u>h</u>	CH ₂ C ₆ H ₅	92	198-200

Although the exact mechanism of this oxidation is still not clear because of no detailed mechanistic studies, the data in Tables I and II leads us to assume that the principal processes can be described as shown in the following scheme. Under the anhydrous conditions, four consecutive reactions, dehydrogenation, intramolecular nucleophilic addition, another dehydrogenation, and isomerization presumably occur to give the oxazolyllindole derivatives (3, 4). The second step (i→ii) must be affected by R', i.e., accelerated by electron-donative substituents, whereas retarded by electron-withdrawing ones. Actually, the oxidation of 2e and 2f was completed within 10 min, whereas a few hours were required in the cases of 2b, 2d, and 2g. The route via vii is improbable because no traces of vii were detected in all cases even in the oxidation with equimolar DDQ. The difference in yield between 3 and 4 owing to R can be explained in terms of the reactivity difference in the final step (iii→3, 4).

Under the aqueous conditions, all the starting materials (1, 2), except 2e and 2f, gave readily the corresponding β-keto derivatives (5, 6). In these cases, the second step must have changed from the intramolecular nucleophilic addition to the intermolecular addition of H₂O (i→iv), followed by another dehydrogenation and isomerization. However, in the oxidation of 2e and 2f, again the cyclization (i→ii) rather than the addition of H₂O must have occurred even in aqueous solvents because of electron-donative substituents (R') and then formed 4e and 4f via iii just as in the anhydrous solvents. If this is the case, 1e and 1f having the same electron-donative groups must also give the intermediates iii, followed by the addition of H₂O to give another intermediates vi, which readily changed to the β-keto compounds (5e, 5f), probably because the isomerization (iii→3) without assistance of carbomethoxy groups must be difficult to occur.¹²

Improvements of the direct synthesis of oxazolylindoles by the DDQ oxidation are now in progress.



ACKNOWLEDGEMENTS

This work was supported by a Grant-in Aid for Special Project Research (Chemical Research in Development and Utilization of Nitrogen-Organic Resources) from the Ministry of Education, Science and Culture.

We thank Professor K. Koyama for helpful advice and discussions.

REFERENCES AND NOTES

1. M. J. Narasimhan and V. G. Ganla, Hindustan Antibiot. Bull., 9, 138 (1967); C. A., 67, 20358j (1967).
2. T. Takeuchi, K. Ogawa, H. Iinuma, H. Suda, K. Ukita, T. Nagatsu, M. Kato, H. Umezawa, and O. Tanabe, J. Antibiot., 26, 162 (1973).
3. D. S. Bhate, R. K. Hulyalkar, and S. K. Menon, Experientia, 16, 504 (1960).
4. B. S. Joshi, W. I. Taylor, D. S. Bhate, and S. S. Karmarker, Tetrahedron, 19, 1437 (1963).
5. K. Koyama, K. Yokose, and L. J. Dolby, J. Org. Chem., submitted.
6. Y. Oikawa and O. Yonemitsu, J. Org. Chem., 42, 1213 (1977).
7. K. Takai, H. Ushiro, Y. Noda, S. Narumiya, T. Tokuyama, and O. Hayaishi, J. Biol. Chem., 252, 2648 (1977); Y. Noda, K. Takai, T. Tokuyama, S. Narumiya, H. Ushiro, and O. Hayaishi, ibid., 253, 4819 (1978); J. Roberts and H. J. Rosenfeld, ibid., 252, 2640 (1977); H. J. Rosenfeld, K. A. Watanabe, and J. Roberts, ibid., 252, 6970 (1977).
8. The biosynthesis of pimprinine is presumably initiated by the side chain oxidation of tryptamine and/or tryptophan derivatives, though no reports have been published.
9. For the removal of 2,3-dichloro-5,6-dicyanohydroquinone (DDHQ) the following method is more practically useful than the alumina column method: After evaporation of the THF, the residues were treated with 0.1 N NaOH and the precipitated solids were collected by filtration, washed with H₂O and a small amount of EtOAc to decolorize, and then dried to give the products in almost pure states. DDHQ recovered from the alkaline solutions can be readily reverted to rather expensive DDQ.¹⁰
10. D. Walker and T. D. Waugh, J. Org. Chem., 30, 3240 (1965).
11. Oxidation of 2f in more aqueous THF (THF:H₂O = 3:1) under dropwise addition of DDQ gave the β -keto product (6f) in 34% yield as well as 4f (20%). This result shows that the formation of 4f and 6f is competitive depending on the concentration of H₂O.
12. The formation of 4e and 4f via the corresponding β -keto products, 6e and 6f, (6→vi→4) is very improbable, because the isolated 6f was completely stable at room temperature even in the presence of TsOH.

Received, 13th August, 1979