

PHOTOCHEMISTRY OF CONJUGATED NITROGEN-CARBONYL SYSTEMS

PHOTOADDITION REACTION OF HYDROXYLAZOBENZENES AND THE RELATED THIONE ANALOGS WITH OLEFINS


Yuichi Kanaoka, Masato Hasebe and Eisuke Sato

Faculty of Pharmaceutical Sciences, Hokkaido University

Sapporo, 060, Japan


As part of systematic photochemical research on conjugated nitrogen-carbonyl systems, which stemmed from our studies on nitrogen-carbonyl systems such as imides,¹⁾ amides and nucleotides, we have started exploring photoreactions of a variety of azobenzenes. In the present work photoaddition reactions of hydroxyazoaromatic compounds and their sulfur analogs with olefins were investigated.

On irradiation in acetonitrile, 3(2H)-pyridazinones 1a undergo [2+2]cycloaddition with olefins such as 2-methyl-2-butene 2 to afford cyclobutane derivatives 3 in a stereoselective manner. By contrast, the photoreaction of their sulfur analog 1b took a quite different course; namely, irradiation of 1b in the presence of 2 mainly gave rise to 4.



In the case of 2-thiouracil, photoaddition of 2 occurred both at the C₅-C₆ double bond and the thione group leading to 5 and 6, respectively. From the parent member of the family, 2-mercaptopyridine and 2-mercaptopyrimidine, the rearranged thiol products 7 were formed under similar conditions.

Finally, 2,4-pyridinedione 8, one of the simplest conjugated nitrogen-carbonyl systems, was examined. As expected, this compound underwent either cyclobutane formation or dimerization depending upon reaction conditions.



REFERENCES

- 1) Y.Kanaoka, Acc.Chem.Res. 11, 407 (1978).