

THE REACTION OF 4-METHOXYQUINOLINE 1-OXIDE WITH DIMETHYL
ACETYLENEDICARBOXYLATE

Yasuhisa Ishiguro, Kazuhisa Funakoshi, Seitaro Saeki, and
Masatomo Hamana*

Faculty of Pharmaceutical Sciences, Kyushu University,
Maidashi, Higashi-ku, Fukuoka 812, Japan

Ikuhiko Ueda


College of General Education, Kyushu University, Ropponmatsu,
Chuo-ku, Fukuoka 810, Japan

Abstract — 4-Methoxyquinoline 1-oxide reacts with dimethyl acetylenedicarboxylate in boiling dioxane affords N-quinolinium ethylide (1), 2-substituted quinoline (2), furo[3,2-c]quinoline (3) and 9-oxo-tetrahydro-cis-furo[3,2-b]quinoline (4). Similar reactions readily proceed at low temperatures in dichloromethane, acetonitrile and DMF to give 2, 3 and 4.

We have recently studied the 1,3-dipolar cycloaddition of some 3-monosubstituted and 3,4-disubstituted quinoline 1-oxides, and have obtained many interesting results¹. This paper mainly deals with our observations on the reaction of 4-methoxyquinoline 1-oxide with dimethyl acetylenedicarboxylate, which was carried out in connection with the above studies.

Treatment of 4-methoxyquinoline 1-oxide with dimethyl acetylenedicarboxylate (1 equiv.) in boiling dioxane for 1 hr gave α -[N-(4-methoxyquinolinium)]- α,β -bis-methoxycarbonyl- β -oxo-ethylide (1) (8.8%), methyl α -methoxycarbonyl- α -[2-(4-methoxyquinolyl)]pyruvate (2) (31.5%), 2,3-bismethoxycarbonylfuro[3,2-c]quinoline (3) (12.2%) and 3a,4,9,9a-tetrahydro-2,3-bismethoxycarbonyl-9-oxo-cis-furo[3,2-b]quinoline (trace). While the reaction at room temperatures in dioxane was very slow, similar reactions readily proceeded even at lower temperatures when dichloromethane, acetonitrile or DMF was used as a solvent instead of dioxane, and products 2, 3 and 4 were obtained, but no formation of N-ylide 1 was noticed in these cases. Table I summarizes these results.

Table I. The Reaction of 4-Methoxyquinoline 1-Oxide
with Dimethyl Acetylenedicarboxylate



Solvent	Reaction		Product, Yield(%)			
	temp. (°C)	time (hr)	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>
dioxane	101	1	8.8	31.5	12.2	trace
CH ₂ Cl ₂	-10	1	-	1.7	9.8	4.6
CH ₂ Cl ₂	0	1	-	22.1	9.2	8.1
MeCN	0	0.5	-	18.8	-	4.6
DMF	0	1	-	34.0	19.5	5.9

Structure assignments of the products are based on the satisfactory elemental analyses, and the IR and PMR spectra shown in Table II. Further, the structure of the 2,3-dihydroquinoline 4 was unambiguously established by an X-ray diffraction study².

Canonne *et al.*³ have recently reported that the reaction of 4-chloroquinoline 1-oxide (5) with diethyl acetylenedicarboxylate in boiling toluene gave the 2-substituted product (6) (mp 91-92°; 3.1%) and the furo[3,2-*c*]quinoline (7) (3.7%). While the PMR spectra of furoquinolines, 3 and 7, were closely similar to each other, the spectral pattern of 2 was found to be considerably different from that of 6 reported as the 2-substituted product by them.

In order to explore the structure of 2, 2 was oxidized by heating with 30% hydrogen peroxide and acetic acid to give 4-methoxyquinaldic acid 1-oxide (8), from which methyl 4-methoxyquinaldate (9) was obtained upon successive treatment with

Table II. The Some Physical Properties of the Products

Products	IR (cm ⁻¹)	PMR (CDCl ₃) δ (ppm)
$\underline{1}$ Yellow needles mp 209-211°	νC=O : 1680, 1725	3.60 (3H, s, COOCH ₃), 3.94 (3H, s, COOCH ₃), 4.28 (3H, s, 4-OCH ₃), 7.16 (1H, d, J _{2,3} =7 Hz, C ₃ -H), 7.68-8.44 (4H, m, Ar-H), 8.64 (1H, d, J _{2,3} =7 Hz, C ₂ -H)
$\underline{2}$ Colorless scales mp 163-165°	νC=O : 1690, 1725	3.76 (3H, s, COOCH ₃), 3.88 (3H, s, COOCH ₃), 4.12 (3H, s, 4-OCH ₃), 7.32-7.80 (3H, m, Ar-H), 8.05 (1H, dd, J _{7,8} =8 Hz, J _{6,8} =2 Hz, C ₈ -H), 8.10 (1H, s, C ₃ -H), 15.40-16.00 (1H, br s, NH)
$\underline{3}$ Colorless needles mp 148-149°	νC=O : 1725	4.06 (3H, s, COOCH ₃), 4.07 (3H, s, COOCH ₃), 7.58-8.44 (4H, m, Ar-H), 9.40 (1H, s, C ₂ -H)
$\underline{4}$ Yellow prisms mp 142-145°	νNH : 3320 νC=O : 1645, 1710, 1740	3.80 (3H, s, COOCH ₃), 3.87 (3H, s, COOCH ₃), 5.17 (1H, d, J _{2,3} =10 Hz, C ₂ -H), 5.35 (1H, d, J _{2,3} =10 Hz, C ₃ -H), 6.68 (2H, m, Ar-H), 7.36 (1H, m, Ar-H), 7.78 (1H, dd, J _{5,6} =8 Hz, J _{5,7} =2 Hz, C ₅ -H)
$\underline{6'}$ Yellow needles mp 135-136°	νC=O : 1690, 1725	1.36 (6H, m, two CH ₃), 4.30 (4H, m, two CH ₂), 7.44-7.88 (3H, m, Ar-H), 8.09 (1H, dd, J _{7,8} =8 Hz, J _{6,8} =2 Hz, C ₈ -H), 16.20-17.00 (1H, br s, NH)

phosphorus trichloride and methanol⁴. Compound $\underline{9}$, colorless needles, mp 146-147°, was proved identical with an authentic sample prepared in the same way from ethyl 4-methoxy-2-quinolinecyanoacetate⁵. Thus, product $\underline{2}$ was established as the 2-substituted quinoline.

This result prompted us to re-examine the reaction described by Canonne *et al.*³ Treatment of $\underline{5}$ with diethyl acetylenedicarboxylate (1 equiv.) for 1 hr in boiling toluene gave not the reported $\underline{6}$ but instead another product ($\underline{6'}$) (4.6%) together with $\underline{7}$ (9.6%). Product $\underline{6'}$, yellow needles, mp 135-136°, has the empirical formula C₁₇H₁₆ClNO₅, and the IR and PMR spectra are consistent with the 2-substituted quinoline structure. Heating $\underline{6'}$ with 30% hydrogen peroxide and acetic acid afforded 4-chloroquinaldic acid 1-oxide ($\underline{10}$), yellow needles, mp 164-165°. Compound $\underline{10}$ was identical with a sample prepared by a similar oxidation of ethyl 4-chloro-2-quinolinecyanoacetate easily obtainable from $\underline{5}$ and ethyl cyanoacetate by means of acetic anhydride. Accordingly, the structure proposed by Canonne *et al.*³ for $\underline{6}$

should be assigned to product 5' isolated by us. Comparison of the PMR spectra of 2 and 6' (Table II) further supports this conclusion; it seems likely that product 6 obtained by Canonne et al. is a different type of compound.

These reactions are formulated in Chart 1.




Chart 1

The primary 1,3-cycloadduct (A) initially formed from 4-methoxyquinoline 1-oxide and dimethyl acetylenedicarboxylate is not stable enough to be isolated, and readily undergoes the N-O bond fission. The formation of the N-ylide 1, the 2-substituted quinoline 2 and the furo[3,2-c]quinoline 3 can be explained by courses a, b and c, respectively, as formulated in Chart 2⁶. However it is difficult at present to rationalize the formation of 9-oxo-tetrahydro-furo[3,2-b]quinoline 4. Course a involves the rearrangement of A to the aziridine intermediate (B), which

isomerizes to 1. The concerted loss of the α -proton with the N-O bond fission in A gives 2 (course b). The formation of 3 follows course c which involves the successive formation of the 2,3-dihydroquinoline (C) and the 3,4-dihydroquinoline (D), and elimination of methanol from D.

Although the de-methylation of the 4-methoxy group of the quinoline ring is inevitable, the formation of 4 is very much noticeable because 4 is the first example of 2,3-dihydroquinolines obtained from the 1,3-dipolar cycloaddition of quinoline 1-oxides. The reaction seems likely to proceed through the 2,3-dihydroquinoline intermediate (E) different from D, however the details of the mechanism is not clear at all, particularly with respect to the de-methylation process. Furthermore, the intermediacy of E postulated above suggests the possibility that the formation of the 2-substituted quinoline 2 follows an alternate path involving E as a precursor, instead of course b^{6,7}.




Chart 2

Further studies are in progress in order to clarify the essential features of the formation of 4 by using various 4-substituted quinoline 1-oxides and dipolarophiles.

ACKNOWLEDGEMENT

We are grateful to a Grant-in Aid for Chemical Research in Development and Utilization of Nitrogen-Organic Resources from the Ministry of Education, Science and Culture, Japan, for partial financial support of this work.

REFERENCES

1. M. Yoshida *et al.*, Abstract Paper of the 11th Congress of Heterocyclic Chemistry, Kanazawa, 1978, 168-172; *Idem*, Heterocycles, 1979, 12, 167; M. Hamana, H. Noda, and M. Aoyama, Heterocycles, 1974, 2, 167; H. Noda, T. Yamamori, M. Yoshida, and M. Hamana, Heterocycles, 1976, 4, 453.
2. I. Ueda, Y. Ishiguro, K. Funakoshi, S. Saeki, and M. Hamana, Acta Crystallographica, 1980 in press.
3. P. Canonne, G. Lemay, and R. A. Abramovitch, Heterocycles, 1978, 9, 1217.
4. Since spectral examinations of compound 8 were rather difficult because of its sparing solubility in organic solvents, it was transformed into 9 as a reference compound.
5. K. Funakoshi, H. Sonoda, Y. Sonoda, and M. Hamana, Chem. Pharm. Bull., 1978, 26, 3504.
6. R. A. Abramovitch and I. Shinkai, Accounts Chem. Res., 1976, 9, 192, and references quoted therein.
7. T. Hisano, T. Matsuoka, M. Ichikawa, and M. Hamana, Heterocycles, 1980, 14,

Received, 24th November, 1979