

## HYDROLYTIC RING CLEAVAGE OF THIADIAZOLO[3,2-a]PYRIMIDINONES


Tadakazu Tsuii\* and Yoshimi Otsuka

Department of Chemistry, Japan Women's University,


Bunkyo-ku, Tokyo, Japan

**Abstract** — The alkali and acid hydrolysis of 2-substituted 7-methyl-5H-thiadiazolo[3,2-a]pyrimidin-5-ones gave 3-amino-6-methyluracil, 3-amino-6-methyl-2-thiouracil and 2-amino-1,3,4-thiadiazole derivatives according to the kind of substituents. While, the treatment of 7-methyl-5H-thiadiazolopyrimidin-5-one with methanolic hydrochloric acid gave 6-methyl-2-thiouracil.

Thiadiazolo[3,2-a]pyrimidine derivatives were synthesized by two routes starting from 3-amino-2-thiouracils<sup>1</sup> and 2-amino-1,3,4-thiadiazoles<sup>2</sup>, and some of them were noted the selective herbicidal activity<sup>2</sup>, but the chemical properties including the stability were little known. Previously, the authors described the alkaline hydrolysis of thiadiazolo[3,2-a]pyrimidines to form 2-thiocyanatopyrimidine derivatives as a result of N-N bond cleavage<sup>3</sup>. Subsequently, Okabe, et al.<sup>4</sup> reported that 7-methyl-5H-thiadiazolo[3,2-a]pyrimidin-5-one (Ia) was hydrolyzed with alkali to 6-methyl-2-thiouracil (III) via 2-thiocyanato derivative (II), while the 2-methyl and 2-benzylthio derivatives of Ia (Ib and If) were converted to 3-acetamido- (IV) and 3-amino-6-methyl-2-thiouracil (V) by C<sub>2</sub>-S




bond cleavage, respectively. Now, the authors found that the acid-catalyzed hydrolysis of I afforded 3-amino- and 3-substituted amino-6-methyluracils (VI and VII) by C<sub>9</sub>-S bond cleavage together with other products, and further revealed that the alkaline hydrolysis of I gave V and/or VII according to the kind of substituents. When a solution of Ib (200 mg) in 5% HCl (2 ml) was refluxed for 2 hr, VIIIb.HCl (R=Me) was obtained in 63% yield. VIIIb.HCl; mp 260° (dec), UV  $\lambda_{\text{max}}^{\text{H}_2\text{O}}$  211, 253, 326 nm; NMR (DMSO-d<sub>6</sub>)  $\delta$  2.26 (9-OH), 2.68 (7-Me), 2.91 (2-Me), 7.45 (ring C-H); IR (KBr) 2640, 1592 cm<sup>-1</sup> (N<sup>+</sup>H<sub>2</sub>). On prolonged reflux with HCl, VIIIb was converted to VII, which was identical with the authentic sample<sup>5</sup>. On similar treatment, Ic (R=SMe) afforded the mixture of 3-(methyldithiocarbamido)-6-methyluracil (VIc, 61%) and VIIIc.HCl (13%). VIc<sup>6</sup>; mp 230° (dec), UV  $\lambda_{\text{max}}^{\text{H}_2\text{O}}$  248, 273 nm (pH 1). VIIIc.HCl; mp 192° (dec); UV  $\lambda_{\text{max}}^{\text{H}_2\text{O}}$  256, 332 nm; IR (KBr) 2590, 1593 cm<sup>-1</sup> (N<sup>+</sup>H<sub>2</sub>). Both VIc and VIIIc.HCl were converted to VII by further treatment with HCl, indicating that the reaction course to VII from I involves the successive formations of the HCl salt of I (IX) and VIII. IX was obtained by the treatment of I with HCl at room temperature. The reaction of Ia (R=H) and Ie (R=NH<sub>2</sub>) with 5% HCl gave VII in 53 and 61% yields, respectively, and that of Id (R=SH) produced the thiadiazole compound (Xd) in 24% yield along with the formation of VII in 64% yield. Xd, mp 245°, was identical with the authentic sample<sup>7</sup>. Although the decomposition of pyrimidine moiety in I was unexpected, the formation of Xd would involve the intermediacy of hydrate compound (XId).




Scheme 2.

When Ic was allowed to stand with HCl-saturated MeOH at room temperature for 2 weeks, VIc was obtained by C<sub>9</sub>-S bond cleavage. On the other hand, the similar treatment of Ia with HCl-saturated MeOH afforded the unexpected 6-methyl-2-thiouracil, III, exclusively. Since the treatment of 2-thiocyanato compound, II, with HCl-saturated MeOH gave III, it was assumed that the reaction involves the intermediacy of II accompanying the N-N bond cleavage.




Scheme 3.

Next, the alkaline hydrolysis of I was attempted. The reaction of 2-methylthio derivative (Ic) (300 mg) with 1% NaOH (10 ml) for 10 days at room temperature gave the mixture of VIc (62%) and V (20%). The reflux of Id (R=SH) with 1% NaOH for 2 hr afforded VII, and that of Ie (R=NH<sub>2</sub>) gave the mixture of V (31%) and VII (23%), which were separated by the preparative thin layer chromatography on silica gel using MeOH-CHCl<sub>3</sub> (3:17).




Scheme 4.

The initial step in  $C_2$ -S bond cleavage of Ib and If might be the attack by hydroxide ion on the  $C_2=N$  double bond as shown in Scheme 5 (XII). On the other hand, the  $C_2$ -S bond cleavage was not observed in the hydrolysis of Id. Such a formation of I-hydroxide adduct (XIII) as illustrated in Scheme 5 might not occur in Id, since the electron-withdrawing action of 2-S<sup>-</sup> substituent would be



Scheme 5.

unfavorable for pushing a  $\pi$ -electron of C<sub>2</sub>=N double bond to the N atom in XII. As depicted in Scheme 6, the C<sub>9</sub>-S bond cleavage in the hydrolysis of Id is presumably initiated by the attack of hydroxide ion on the C<sub>9</sub> atom. The weak electron-with-



Scheme 6.

drawing action of 2-methylthio and 2-amino substituents in Ic and Ie would cause the attack by hydroxide ion on both C<sub>2</sub> and C<sub>9</sub> atoms.

#### REFERENCES

1. T. Tsuji and T. Ueda (née Yamaka), Chem. Pharm. Bull. (Tokyo), 1971, 19, 2530.
2. T. Okabe, M. Maekawa, and E. Taniguchi, Agric. Biol. Chem., 1973, 37, 1197.
3. T. Tsuji and Y. Kamo, Chem. Lett., 1972, 641.
4. T. Okabe, E. Taniguchi, and K. Maekawa, Bull. Chem. Soc. Japan, 1974, 47, 2813.
5. T. Tsuji and Y. Otsuka, Chem. Pharm. Bull. (Tokyo), 1978, 26, 2765.
6. All new compounds gave carbon, hydrogen and nitrogen analysis within normal limits.
7. P.C. Guha, J. Amer. Chem. Soc., 1922, 44, 1504.

Received, 8th November, 1979