

A NEW METHOD FOR SYNTHESIS OF LACTAM

Yoshihisa Kurasawa* and Atsushi Takada

School of Pharmaceutical Sciences

Kitasato University, Shirokane, Minato-ku, Tokyo 108, Japan

Abstract — 2-Substituted 3-oxo-3,4-dihydroquinoxalines were oxidized with H_2O_2 to 2,3-dioxo-1,2,3,4-tetrahydroquinoxaline. This oxidation of the imino bonding in the heterocycles to the lactam was assumed to proceed via the various peroxides, but not via the oxaziranes.

The oxidation of Schiff bases (imines, $R_1R_2C=NR_3$) with peracid gives oxaziranes,^{1,2,3} which are further converted to amides with ferrous ion and by pyrolysis (at 300°C).² The oxaziranes are also obtained from hydroperoxides, which are produced by the reaction of Schiff bases with H_2O_2 ⁴ (Chart 1-a). We now report a novel one-step oxidation of imino bonding in heterocycles to lactam with H_2O_2 (Chart 1-b). 2-Substituted 3-oxo-3,4-dihydroquinoxalines were used as the heterocycles (Chart 2).

A solution of 2-methyl-3-oxo-3,4-dihydroquinoxaline (I)⁵ (3.1 mmol) and H_2O_2 ⁶ (31 mmol) in AcOH




Chart 1

(30 ml) was heated on a water bath to afford 2,3-dioxo-1,2,3,4-tetrahydroquinoxaline (V),⁷ in 6% yield. A similar reaction of 4-(3'-oxo-3',4'-dihydroquinoxaline-2'-yl)pyrazolones (IIa-c)⁸ (3.3 mmol) with H₂O₂ (165 mmol) in AcOH (100 ml) brought about V, in 40% yield. 2-Ethoxycarbonylmethylene-3-oxo-1,2,3,4-tetrahydroquinoxaline (III)⁹ (4.3 mmol) and 3-(N,N-dimethylaminocarbonyl)furo-[2,3-b]quinoxaline (IV)⁸ (3.6 mmol) also gave V in the presence of 10-fold molar amount of H₂O₂, in 60% and 65% yields, respectively. From the above reaction mixture, the oxaziranes and a moiety other than quinoxaline were not obtained.

As shown in Table I, it is evident that the yield of V is low when 2-substituent in the quinoxaline ring is methyl group, but its yield is improved when the 2-substituent includes α,β -unsaturated bonding (C=C). This may indicate that the addition of H₂O₂ to I, II, IIIa,⁹ and IV⁸ produces the peroxides (II),¹⁰ (IIIi), (IIIj), and (IVi) as the respective intermediates, as shown in Chart 3. However, these peroxides were not isolated, presumably due to ease of decomposition. Compound V may be formed from these peroxides in a radical or concerted mechanism.^{11,12}

On the other hand, the mechanism via the oxaziranes should be eliminated in the above reactions, since ferrous ion or heating at a higher temperature was necessary for the formation of the amide from the oxazirane, even if the oxazirane was produced. Furthermore, when the solution of the oxazirane was heated near 100°C for a long time, the nitrone was brought about.²

In conclusion, it is of interest that the oxidation of the imino bonding in the heterocycles with H₂O₂ generates the lactam via the peroxides, but not via the oxaziranes.





Chart 2

Table I

| Compound | Molar ratio of H ₂ O ₂ | Reaction time | Yield of V (%) |
|----------|--|---------------|----------------|
| I | 10-fold | 1 hr | 6 |
| II | 50-fold | 3 hr | 40 |
| III | 10-fold | 1 hr | 60 |
| IV | 10-fold | 1 hr | 65 |



* reference No.8

Chart 3

References and Footnotes

1. W. D. Emmons, J. Amer. Chem. Soc., 1956, 78, 6208.
 2. Idem, ibid., 1957, 79, 5739.
 3. E. Schmitz, Angew. Chem. internat. Edit., 1964, 3, 333.
 4. E. Hoefl and A. Rieche, Angew. Chem. internat. Edit., 1965, 4, 524.
 5. C. L. Lees and H. N. Rydon, J. Chem. Soc., 1955, 303.
 6. 30% solution of H_2O_2 in H_2O was employed.
 7. Spectral and physical data of V were identical with the sample prepared by the reaction of o-phenylenediamine with diethyl oxalate.
 8. Y. Kurasawa and A. Takada, Heterocycles, 1980, 14, 281.
 9. R. Mondelli and L. Merlini, Tetrahedron, 1966, 22, 2356.
 10. A. Rieche, E. Schmitz, and E. Beyer, Chem. Ber., 1959, 92, 1212.
 11. C. F. Cullis, A. Fish, and D. W. Turner, Proc. Roy. Soc. (London), 1961, A262, 318; Idem, ibid., 1962, A267, 433.
 12. The mechanism like the Baeyer-Villiger reaction was denied, since it did not lead to the formation of compound V.
- H. Ishii, T. Ishikawa, S.-T. Lu, and I.-S. Chen, Tetrahedron Letters, 1976, 1203.

Received, 22th December, 1979