

STEREOSELECTIVE REDUCTION OF 1,2,3,4,6,7,12,12b-OCTAHYDROINDOLO-
[2,3-a]QUINOLIZINE WITH SODIUM BOROHYDRIDE IN TRIFLUOROACETIC ACID¹


Gordon W. Gribble*, Jerry L. Johnson, and Mark G. Saulnier
Department of Chemistry, Dartmouth College, Hanover, New
Hampshire 03755 USA

Abstract - The sodium borohydride/trifluoroacetic acid reduction of the title alkaloid (1) is highly stereoselective, giving the all-cis product 2 in excellent yield.



Recent years have seen great improvements in the methodology for reducing indoles to dihydroindoles (indolines). These newer methods, generally under acidic conditions, with borane,² amine-boranes,³ boron hydrides,⁴ and trialkylsilanes,⁵ supersede the older methods (i.e., metals in mineral acid, catalytic hydrogenation)⁶ for generality, convenience, and efficiency. Furthermore, the use of trifluoroacetic acid in several of these methods^{2,3d,4b,4c,4e,5b} allows one to reduce 1,2,3,4-tetrahydro- β -(and γ)-carbolines and other indoles containing a second, basic nitrogen. These particular indoles can be difficult to reduce under weaker acid conditions because protonation of the basic nitrogen suppresses protonation of the indole double bond, which is presumed necessary for successful reduction.

For example, sodium borohydride (NaBH_4) in trifluoroacetic acid ($\text{CF}_3\text{CO}_2\text{H}$) has been used to reduce several indole[2,3-a]quinolizidines and other indole alkaloids (i.e., yohimbine,⁷ reserpiline,^{7b} ibogaine,^{7b} tetraphylline,^{7b} raubasine^{7b}). Although these reductions were reported to be stereoselective, no attempt was made to ascertain the stereochemistry of the major product.^{7b} To answer this question we have studied the reaction of the readily available⁸ alkaloid 1,2,3,4,6,7,12,12b-octahydroindolo[2,3-a]quinolizine (1)⁹ with $\text{NaBH}_4/\text{CF}_3\text{CO}_2\text{H}$.^{10,11}

Treating a solution of 1 in neat $\text{CF}_3\text{CO}_2\text{H}$ at -5° to 0° with NaBH_4 (pellets) gave 1,2,3,4,6,7,7a,12,12a,12b-decahydroindolo[2,3-a]quinolizine (2) as one major stereoisomer, mp $75-75.5^\circ$, in 90% purified yield.^{12,13} The ^1H -nmr spectrum of 2 reveals one distinguishing feature: a one-proton doublet of doublets at 3.55 ppm



($J=2.1$ and 6.6 Hz), suggestive of H_A .¹⁴ To identify this signal unambiguously and to assign the two associated coupling constants, we prepared the deuterated derivatives 4-6 as shown below.




Treating 1 with $\text{NaBD}_4/\text{CF}_3\text{CO}_2\text{H}$ produced 4 in 66% yield whose ^1H -nmr spectrum showed a greatly diminished signal at 3.5 ppm, confirming this as arising from H_A . Treating 1 with $\text{NaBH}_4/\text{CF}_3\text{CO}_2\text{D}$ produced 5 in 88% yield which displayed H_A as a

doublet (2.1 Hz), and treating 3¹⁵ with $\text{NaBH}_4/\text{CF}_3\text{CO}_2\text{H}$ produced 6 in 65% yield which displayed H_A as a doublet (6.6 Hz). Thus, the two couplings involving H_A are assigned $J_{AC}=2.1$ Hz and $J_{AB}=6.6$ Hz.

Using Dreiding stereomodels¹⁶ we have examined all of the reasonable conformations of the four diastereomers of 2, and measured the appropriate dihedral angles in each with a Dreiding torsionmeter¹⁶ to estimate J_{AB} and J_{AC} from the Karplus equation.¹⁷⁻¹⁹ The only pair of dihedral angles ($\text{H}_B\text{C}_{7a}\text{C}_{12a}\text{H}_A$ and $\text{H}_A\text{C}_{12a}\text{C}_{12b}\text{H}_C$) that gives reasonable agreement between the predicted and observed coupling constants J_{AB} and J_{AC} belongs to a conformation having the cis-cis configuration for 2 (shown below). The value $J_{AB}=6.6$ Hz (dihedral angle $25-30^\circ$) is in excellent agreement with the comparable couplings reported for cis-1,2,3,4,4a,9a-hexahydrocarbazole (6 Hz)²⁰ and for a cis-2,3,4,4a,5,9b-hexahydro-1H-pyrido[4,3-b]indole derivative (6.6 Hz),^{4c} both of which have a cis-fused five-six ring system. In contrast, a five-six trans-ring fusion is known to produce larger coupling (e.g., 11.8 Hz in trans-hexahydrocarbazole²⁰). Similarly, the ^{13}C chemical shifts for C-7a (40.3 ppm) and C-12a (63.9 ppm) in 2 (signals which respectively disappear in the ^{13}C spectra of 5 and 4) match well with the corresponding chemical shifts reported for the cis-fused pyrido[4,3-b]indole derivative cited above (41.1 and 57.5 ppm) but not for the trans-fused isomer (47.3 and 68.4 ppm).

The value $J_{AC}=2.1$ Hz (dihedral angle $40-45^\circ$) is smaller than the Karplus equation would generally predict, and most likely is a consequence of the two electronegative nitrogens attached to the coupling fragments, since this is known to decrease vicinal coupling, sometimes substantially.^{18,19} A trans-stereochemistry of H_AH_C , regardless of the H_AH_B stereochemistry, would have given rise to a large J_{AC} (8-12 Hz²⁰). Therefore, we assign the cis-cis configuration to 2.



This stereochemistry is reasonable mechanistically since one anticipates stereo-electronic axial protonation²¹ of 1 giving cis-indolenium ion 7, followed by hydride delivery by the bulky bis- or tris(trifluoroacetoxy)borohydride species¹⁰ from an equatorial direction²² giving the generally presumed^{2a,3b,6,14} more stable five-six cis-fused indoline ring.²³ Indeed, treating a solution of 1 and NaBH₄ in tetrahydrofuran with CF₃CO₂H, conditions which produce the more reactive and less selective reducing agent sodium mono(trifluoroacetoxy) borohydride,¹⁰ gives 2 in only 20% yield along with at least two of the other diastereomers.

Acknowledgments

This investigation was supported in part by Grant CA-24422, awarded by the National Cancer Institute, DHEW, and by Merck Sharp and Dohme Research Laboratories. We also thank Dr. Catherine E. Costello (Massachusetts Institute of Technology) for the high resolution mass spectra (National Institutes of Health Resource Grant FR00317 from the Division of Research Facilities and Resources), and the Colorado State University Regional NMR Center, funded by National Science Foundation Grant No. CHE 78-18581, for high-resolution NMR spectra.

References and Notes

1. This paper is Part X in the series "Reactions of Sodium Borohydride in Acidic Media." For Part IX, see G. W. Gribble, W. J. Kelly, and M. P. Sibi, Synthesis, submitted.
2. (a) B. E. Maryanoff and D. F. McComsey, J. Org. Chem., 1978, 43, 2733; (b) B. E. Maryanoff, D. F. McComsey, and S. D. Nortey, J. Org. Chem., 1981, 46, 355.
3. (a) J. G. Berger, Synthesis, 1974, 508; (b) J. G. Berger, S. R. Teller, C. D. Adams, and L. J. Guggenberger, Tetrahedron Lett., 1975, 1807; (c) Y. Kikugawa, J. Chem. Res., 1977, 212; (d) Y. Kikugawa, J. Chem. Res., 1978, 184.
4. (a) G. W. Gribble and J. H. Hoffman, Synthesis, 1977, 859; (b) N. F. Kuchero-ova, N. N. Novikova, N. M. Sharkova, I. D. Silenko, and V. A. Zagorevskii, Khim. Geterostsikl. Soedin., 1977, 995; C.A., 1977, 87, 167920; (c) J. G. Berger, F. Davidson, and G. E. Langford, J. Med. Chem., 1977, 20, 600; (d) T. Wakamatsu, H. Inaki, A. Ogawa, M. Watanabe, and Y. Ban, Heterocycles, 1980, 14, 1441; (e) N. F. Kuchero-ova, N. M. Sipilina, N. N. Novikova, I. D. Silenko, S. G. Rozenberg, and V. A. Zagorevskii, Khim. Geterostsikl. Soedin.,

- 1980, 1383; C.A., 1981, 94, 121372.
5. (a) (review) D. N. Kursanov, Z. N. Parnes, and N. M. Loim, Synthesis, 1974, 633; (b) A. E. Lanzilotti, R. Littell, W. J. Fanshawe, T. C. McKenzie, and F. M. Lovell, J. Org. Chem., 1979, 44, 4809.
 6. B. Robinson, Chem. Rev., 1969, 69, 785.
 7. (a) D. Herlem and F. K-Huu, Tetrahedron, 1979, 35, 633; (b) J. LeMen, L. LeMen-Oliver, J. Levy, M. C. Levy-Appert-Colin, and J. Hannart, German patent 2,410,651; C.A., 1975, 82, 43640.
 8. G. W. Gribble, J. Org. Chem., 1972, 37, 1833.
 9. Dracontomelum mangiferum: S. R. Johns, J. A. Lamberton, and J. L. Occolowitz, Chem. Comm., 1966, 421; Aust. J. Chem., 1966, 19, 1951. The N(12)-methyl derivative of 1 is found in a cranberry species: K. Jankowski, S. Godin, and N. E. Cundasawmy, Can. J. Chem., 1974, 52, 2064.
 10. For experimental conditions and discussions on the use of this reagent, see G. W. Gribble, R. M. Leese, and B. E. Evans, Synthesis, 1977, 172; G. W. Gribble, W. J. Kelly, and S. E. Emery, Synthesis, 1978, 763; G. W. Gribble, Eastman Org. Chem. Bull., 1979, 51, 1.
 11. In connection with an earlier study on the reduction/alkylation of indoles we reported the reduction of 1 but did not determine the product stereochemistry: G. W. Gribble, et al., J. Am. Chem. Soc., 1974, 96, 7812.
 12. 2: ^1H NMR (CDCl_3) δ 1.1-2.2 (m, 11H), 2.6-3.1 (m, 3H), 3.55 (dd, 1H), 6.5-7.2 (m, 4H); ^{13}C NMR (CDCl_3) δ 24.9 (C-2), 25.6 (C-3), 29.6 (C-1), 29.8 (C-7), 40.3 (C-7a), 54.6 (C-6), 56.7 (C-4), 62.8 (C-12b), 63.9 (C-12a), 110.5 (C-11), 118.6 (C-9), 122.9 (C-10), 127.0 (C-8), 134.7 (C-7b), 149.6 (C-11a); mass spectrum, m/e 228.1620 (M^+ , calcd 228.1626), 144, 111, 98 (100%), 83, 69, 55. IR (CHCl_3) 3390, 2940, 2855, 2805, 2765, 1610, 1485, 1465, 1355, 1245, 1125, 1105 cm^{-1} . Anal. calcd for $\text{C}_{15}\text{H}_{20}\text{N}_2$: C, 78.90; H, 8.83; N, 12.27. Found: C, 78.74; H, 8.85; N, 12.21.
 13. There was also isolated 5% of 1 (flash chromatography); two minor reaction components (tlc) have not been identified but are probably diastereomers.
 14. For comparison, the corresponding proton (H-9a) in cis-hexahydrocarbazole resonates at 3.67 ppm: S. A. Monte and R. R. Schmidt, Tetrahedron, 1971, 27, 3331.
 15. G. W. Gribble and R. B. Nelson, J. Org. Chem., 1974, 39, 1845.
 16. Brinkmann Instruments, Inc. Cantiague Road, Westbury, N.Y. 11590, USA.

17. M. Karplus, J. Chem. Phys., 1959, 30, 11.
18. Although vicinal proton coupling depends also on substituent electronegativity, bond angles, and bond lengths, dihedral angle measurements give a good approximation to the relative magnitude ("large", "medium", or "small") of vicinal couplings: M. Karplus, J. Am. Chem. Soc., 1963, 85, 2870.
19. For excellent discussions on the use (and abuse) of the Karplus relationship, see S. Sternhell, Quart. Rev., 1969, 23, 236; L. M. Jackman and S. Sternhell, "Applications of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry," 2nd edition, Pergamon, Oxford, 1969, pp. 280-301.
20. H. Booth and T. Masamune, J. Chem. Soc., Perkin Trans. II, 1972, 354.
21. H. O. House, "Modern Synthetic Reactions," 2nd ed., W. A. Benjamin, Menlo Park, CA, 1972, pp. 574-5.
22. For example, reduction of 4-t-butylcyclohexanone with Li(sec Bu)₃BH gives 96.5% stereoselective equatorial attack: H. C. Brown and S. Krishnamurthy, J. Am. Chem. Soc., 1972, 94, 7159. For highly stereoselective reductions of N-methylcyclohexylimines with this reagent (and others), see D. A. Evans, A. M. Golob, N. S. Mandel, and G. S. Mandel, J. Am. Chem. Soc., 1978, 100, 8170.
23. Maryanoff has recently described a similar case of axial protonation of an indole double bond followed by equatorial attack on the resulting iminium ion with borane/TFA: B. E. Maryanoff, D. F. McComsey, R. J. Taylor, Jr., and J. F. Gardocki, J. Med. Chem., 1980, 24, 79.

Received, 17th August, 1981