

OLEFINIC CYCLIZATION OF α -CARBAMOYL- α -THIOCARBOCATION

Yasumitsu Tamura, Hiroshi Maeda, Shuji Akai, Ko-ichi Ishiyama,
and Kiroyuki Ishibashi

Faculty of Pharmaceutical Sciences, Osaka University,

1-6 Yamada-oka, Suita, Osaka, Japan

The α -carbamoyl- α -thiocarbocation ($RS=CHCONR'_2$), the Pummerer reaction intermediate of α -sulfinylacetamide [$RS(O)CH_2CONR'_2$], was found to behave as a highly reactive initiating center for cationic olefin cyclization. Thus, treatment of N-(2-methyl-2-propenyl)-N-methyl- α -(methylsulfinyl)acetamide (Ia) with $(CF_3CO)_2O$ in CH_2Cl_2 caused the cyclization through a Pummerer reaction intermediate (A, $R^1=Me$, $R^2=R^3=H$) to give the six-membered lactams II and III in 43 and 35% yields. Under the same reaction conditions, N-2-butenyl-N-methyl- α -(methylsulfinyl)acetamide (Ib) afforded the five-membered lactam IV in 92% yield and N-(3-methyl-2-butenyl)-N-methyl- α -(methylsulfinyl)acetamide (Ic) gave a mixture of the five-membered lactams V and VI (ca. 4:1) in 93% yield. Treatment of N-2-propenyl-N-methyl- α -(methylsulfinyl)acetamide (Id) by a similar method to that employed for the cyclization of Ia-c did not afford a cyclized product but a Pummerer rearrangement product VII. However, further treatment of VII with CF_3COOH (in the absence of CH_2Cl_2) caused the cyclization to give the five-membered lactams VIII and IX in 9 and 39% yields. Similar treatment of X derived from N-2-propenyl-N-methyl- α -(phenylsulfinyl)acetamide (Ie) gave the tricyclic compound XI in 33% yield along with the five-membered lactams XII (21%) and XIII (14%).

