


RING-ENLARGEMENT REACTIONS OF BENZENE DERIVATIVES
WITH ALKOXYCARBONYLNITRENE

Tsutomu KUMAGAI, Kyosuke SATAKE, Kazuo KIDOURA and Toshio MUKAI
Department of Chemistry, Faculty of Science, Tohoku University
Sendai 980, Japan.

Thermal reactions of methyl azidoformate with 1,4-di-*tert*-butylbenzene(1) were investigated. We report first the formation of 1H-azepines(2 and 3) and dinitrogenated products(4 and 5) as well as the reaction mechanism for these products. The reaction mixture that was obtained by heating 1 and methyl azidoformate at 125°C for 3 hrs was separated by means of preparative low pressure liquid chromatography to give 2,3,4 and 5 in 8.2,5.8,17.7 and 8.8% yield respectively, and another unknown product. The dinitrogenated product 4 could also be derived from 2 in 25% yield by heating with methyl azidoformate, but not from 3. In addition, heating of 5 at 230°C for 6 hrs led to 4 in quantitative yield.



Dehydrogenation of 4 and decarbonylation of 4 and its dehydrogenated product 6 were studied to gain insight into the properties of pyrrolo[3,2-*b*]pyrrole derivatives. Heating of 4 with DDQ produced 6 in quantitative yield. When 7 or 4 was treated with KOH in ethanol at room temperature or at 50°C, decarbonylation partially occurred to afford mono methoxycarbonyl derivative 7 or 8 in 96 or 55% yields. On the other hand, refluxing 4 with alcoholic-KOH produced pyrrole derivative 9 in 68% yield.

