

THERMAL RING-EXPANSION OF 6-ETHYNYL-1,2,5,6-TETRAHYDRO-
PYRIDINE N-IMIDES INTO DIAZONINE DERIVATIVES

Haruki Sashida and Takashi Tsuchiya*

School of Pharmacy, Hokuriku University,
Kanagawa-machi, Kanazawa, 920-11, Japan

Abstract — The thermolysis of the 6-ethynyl-1,2,5,6-tetrahydro-
pyridine N-imides (5 and 6) resulted in the [2,3]-sigmatropic
rearrangement with the ethynyl group to give the novel dihydro-
diazonines (9) via the allene intermediates (8).

Much attention has been given recently to the utilization of thermal sigma-
tropic rearrangements of various kinds of ylides in organic synthesis.¹ As regards to
aminimides, open-chain allylamine N-imides are known to undergo competing [1,2]-
and [2,3]-sigmatropic rearrangements, but the latter rearrangement usually predomi-
nates over the former, whereas pentadienylamine N-imides undergo [1,2]- and [2,5]-
rearrangements predominantly over the [2,3]-rearrangement.² As for cyclic amines,
saturated piperidine and pyrrolidine N-imides have been shown to undergo the [1,2]-
rearrangement to give the corresponding ring-expansion products.³ These results
and those of the thermolysis of cyclic S-ylides⁴ prompted us to examine the thermal
behavior of unsaturated cyclic amine ylides and we have already reported that the




Chart 1

N-imides (1a-3a) undergo only [2,3]-rearrangement and no [1,2]-rearrangement.⁵ In contrast, the thermolysis of the N-oxides (1b-3b) results in the [1,2]-rearrangement to give the corresponding 1,2-oxazepine derivatives.⁶



In connection with these results and the thermal behavior of prop-2-ynylamine ylides,⁷ we studied the thermolysis of the title N-imides with an ethynyl group and report here the formation of novel nine-membered heterocycles, diazonines, by the [2,3]-rearrangement with the triple bond via allene intermediates.

The 6-ethynyl-1,2,5,6-tetrahydropyridines (4a-c)⁸ were aminated with O-mesitylenesulfonylhydroxylamine (MSH)⁹ and the resulting N-aminopyridinium salts were acetylated with acetic anhydride followed by base treatment to give ca. 4:1 mixtures of N-acylimides (5) and (6) in 50-75% yields, which were separated by alumina chromatography. Structures including stereochemistry of these N-imide diastereomers were elucidated by ¹H-NMR spectral analyses.¹⁰ By analogy to the piperidine alkylation,¹⁰ axial approach of the amination reagent on the diequatorial conformer of the amine (4) may predominate to give the diastereomer (5) as the major product.

Thermolysis of both N-imides (5) and (6) in refluxing xylene for 1-2 h gave the tetrahydropyrazoles (7) and the 5,6-trans-dihydrodiazonines (9) in the yields shown in Chart 2, respectively.¹¹ The pyrazoles (7) may be formed directly from the imides by the [2,3]-rearrangement with the double bond in the ring analogous to that observed for the N-imides (1).⁵ On the other hand, the formation of the ring-expansion products (9) may involve initial formation of the allene intermediates (8) by another [2,3]-rearrangement with the triple bond by analogy with the cases of open-chain prop-2-ynylamine ylides.⁷ The allene intermediates (8) then undergo a 1,3-hydrogen shift to give the products (9).



The results collected in Chart 2 show that the yields of 9 from the imides (6), in which the imide group and the ethynyl group are trans, are 3-4 times higher than those from 5. In addition, the thermolysis of 5 required somewhat longer times than those for 6. This lower or reduced susceptibility of the cis-imides (5) toward the reaction is probably due to the transition state of the concerted rearrangement requiring the unfavored boat form of the tetrahydropyridine ring, in contrast, the trans-diastereomers (6) may be able to form the favored chair form transition state, as indicated in the structures (5') and (6').

Treatment of the trans-dihydrodiazonines (9) with sodium ethoxide in toluene



5a	7%	15%
5b	8%	18%
5c	—	16%
6a	6%	55%
6b	7%	48%
6c	—	70%

Chart 2



resulted in the isomerization to give the cis-isomers (10) in high yields.¹² Studies on the detailed stereochemistry of the present reaction and on synthetic application of the results are further in progress.

References and Notes

- 1) For reviews, see W.J. Mckillip, E.A. Sedor, B.M. Culbertson, and S. Wawzonek, Chem. Rev., 1973, 73, 255; T.L. Gilchrist and C.J. Moody, ibid., 1977, 77, 409; E.C. Taylor and I.J. Turchi, ibid., 1979, 79, 181; H. Hayashi, J. Syn. Org. Chem. Japan, 1976, 34, 396; T. Nakai and K. Mikami, ibid., 1980, 38, 381.
- 2) K. Chantrapromma, W.D. Ollis, and I.O. Sutherland, J. Chem. Soc. Chem. Commun., 1977, 97.
- 3) S. Wawzonek and R.C. Gueldner, J. Org. Chem., 1965, 30, 3031; S. Wawzonek and J.G. Stephanie, ibid., 1971, 36, 2467.
- 4) For examples, E. Vedejs, J.P. Hagen, B.L. Roach, and K.L. Spear, J. Org. Chem., 1978, 43, 1185; E. Vedejs, M.J. Arco, D.W. Powell, J.M. Renga, and S.P. Singer, ibid., 1978, 43, 4831; V. Cere, C. Paolucci, S. Pollicino, E. Sandri, and A. Fava, ibid., 1979, 44, 4128.
- 5) T. Tsuchiya, H. Sashida, and H. Sawanishi, Chem. Pharm. Bull., 1978, 26, 2880; T. Tsuchiya and H. Sashida, ibid., 1981, 29, 1887.
- 6) T. Tsuchiya and H. Sashida, Heterocycles, 1980, 14, 1925.
- 7) J.C. Greig, N.N. Ekwuribe, and L.D. Gruenke, Tetrahedron Lett., 1979, 4025; G. Hallstrom, B. Lindke, A.H. Khuthier, and A.M. Al-Iraqi, ibid., 1980, 667; S. Mageswaran, W.D. Ollis, D.A. Southam, I.O. Sutherland, and Y. Thebtranth, J. Chem. Soc. Perkin I, 1981, 1969.
- 8) These compounds were prepared from 2-bromopyridine by successive ethynylation, methylation, and NaBH₄ reduction.
- 9) Y. Tamura, J. Minamikawa, and M. Ikeda, Synthesis, 1977, 1.
- 10) T.M. Bare, N.D. Hershey, H.O. House, and C.G. Swain, J. Org. Chem., 1972, 37, 997.
- 11) Satisfactory elemental analyses and spectral data were obtained for all new compounds reported.
- 12) The ¹H-NMR spectra of the trans-diazonines (9) showed a temperature dependence consistent with the predictable temperature-dependent inversion of the ring, in contrast, those of the cis-isomers (10) showed no such dependence.

Received, 11th November, 1981