





THE N,N' -DIACYL-4,5-DIHYDROIMIDAZOLIUM ION AS AN ELECTROPHILE

Jan Bergman*, Harris Goonewardena and Birger Sjöberg
 Royal Institute of Technology, Department of Organic Chemistry
 S-100 44 Stockholm 70, Sweden

Abstract - 2-Methyl-4,5-dihydroimidazole/acetic anhydride reacts smoothly with indole at room temperature giving N,N' -diacetyl-2-methyl-2-(3-indolyl)-imidazolidine. This adduct can be hydrolysed to 3-acetylindole. Other aromatics, like 1,3-dimethoxybenzene, undergo similar reactions.

When heated with an imidazole/acetic anhydride reagent, indole afforded^{1,2} the adduct 1. The more reactive reagent imidazole/trifluoroacetic anhydride similarly yielded *e.g.* the adducts 2 and 3 with thiophene and 1,3-dimethoxybenzene, respectively.³ The electrophilic species attacking the aromatic ring was assumed to have the structure 4. We therefore felt that it would be of interest to generate the less stabilized and hence more reactive ion 5 from 4,5-dihydroimidazoles and suitable anhydrides in the presence of aromatic compounds. The results of such a study are reported herein.















Indole reacted smoothly at room temperature⁴ with the 2-methyl-4,5-dihydroimidazole/acetic anhydride reagent yielding 6a. 2-Ethyl-4,5-dihydroimidazole and 2-phenyl-4,5-dihydroimidazole similarly (55°, 5 h) gave 6b and 6c, respectively. In contrast, reactions at reflux temperature (140°C) yielded the ring opened product 7. Compounds 6a and 7 could be readily hydrolysed (5% sodium hydroxide in ethanol/water) to 3-acetylindole in high yield. Compound 6c similarly yielded 3-benzoylindole⁵, whereas treatment with sodium borohydride in hot methanol smoothly yielded 3-benzylindole in 93% yield. The yields of 6a - 6c were in the range 85-95%, whereas the yield of 7 was 62%.

Our interest in precursors⁶ to the indole alkaloid ellipticine and related compounds also induced us to treat various indoles with 2-(3-pyridyl)-4,5-dihydroimidazoles and acetic anhydride. The yields of 8 as well as 9 were good to excellent. The hydrolysis of the adduct 8d was in contrast to 8a-c extremely sluggish, a fact that seems to indicate that deprotonation of 8a-c, followed by indolenine-formation (*i.e.* ring-cleavage of the tetrahydroimidazole ring) are important steps preceding the hydrolysis.

From these results, it appears that the ion 5, as expected, is a more powerful electrophile than the ion 4. However, the ion 5 ($R_1=CH_3$) seems to decompose under more forcing conditions and in several attempts to substitute 1,3-dimethoxybenzene, compound 10a and even 10b were the only products.

Adducts 11a and 11b could however be prepared from 1,3-dimethoxybenzene and the appropriate 4,5-dihydroimidazole with trifluoroacetic anhydride in acetonitrile (yields 80-90%).





In contrast 2-methyl-4,5-dihydroimidazole with the same reactants at the same conditions yielded quite a different product assigned structure 12⁹. This assignment is based on MS and NMR data ¹⁰, as well as degradation studies. Alkaline hydrolysis yielded compound 13, whose structure was confirmed by an independent condensation of the carbinol 14 with 1,3-dimethoxybenzene. Furthermore it was found that 2-methyl-4,5-dihydroimidazole and trifluoroacetic anhydride could be converted to compounds 15a and 15b and 2-phenyl- and 2-ethyl-4,5-dihydroimidazoles similarly yielded 15c and 15d, respectively. Compound 15b could be readily hydrolysed to the strongly hydrogen-bonded compound 16. When 2-methyl-4,5-dihydroimidazole was refluxed with (CF₃CO)₂O in acetonitrile for 2 h, the related C-di-COCF₃-substituted compound 17¹¹ could be immediately isolated in high yield as a stable crystalline solid when the reaction mixture was poured into water.

REFERENCES AND NOTES

1. J. Bergman, *Tetrahedron Letters* 4723 (1972).
2. J. Bergman, B. Sjöberg and L. Renström, *Tetrahedron* 36, 2505 (1980).
3. These adducts can readily be converted² to the corresponding aldehydes.
4. In contrast indole fails to give an adduct with the 2-methylimidazole/acetic anhydride reagent even at reflux temperature.
- 5a. This route to 3-benzoylindole is much more convenient than the route given by Stütz and Stadler^{5b}, involving attack of indole with a 1,3-dithiolium ion in the first step.
- 5b. P. Stütz and P.A. Stadler, *Org. Synth.* 56, 8 (1977).
6. J. Bergman and R. Carlsson, *Tetrahedron Letters*, 4663 (1977).
7. Compound 7; m.p. 130-131°C; MS, *m/e* (% relative intensity): 370 (23), 369 (100), 327 (17), 285 (24), 268 (17), 267 (16), 266 (23), 255 (20) and 253 (14). Only peaks stronger than 10% of the base peak and *m/e* above 250 are given; ¹H NMR (CDCl₃, 200 MHz): δ = 2.18 (s, 3 H, COCH₃), 2.46 (s, 6 H, COCH₃), 2.65 (s, 3 H, COCH₃), 3.63 (dd, 2 H, CH₂), 3.96 (dd, 2 H, CH₂), 5.36 (s, 1 H, C = CH₂), 5.83 (s, 1 H, C = CH₂), 7.35-7.69 (several peaks, 4 H) and 8.51 (d, 1 H).
8. Compound 11a; m.p. 145-146°C; MS, *m/e* (% relative intensity): 477 (25), 476 (100), 408 (10), 407 (42), 399 (11) and 350 (17). Only peaks stronger than 10% of the base peak and above *m/e* 250 are given.
- 9a. Compound 11c was not formed at all.
- 9b. Thiophene yielded an adduct analogous with 12.
- 10a. Compound 12; m.p. 139-140°C; MS, *m/e* (% relative intensity): 631 (28.8), 630 (100), 590 (16.7), 423 (25.4), 411 (14.3), 219 (14.3), 70 (56.0); Calculated for M.w. C₂₆H₂₄F₉N₂O₆; 630.1412. Found: 630.1414. The yield of this compound was 84%.
- 10b. We thank Dr G. Höhne, Technische Universität, Berlin, for this measurement.
11. Compound 17; m.p. 213-214°C; IR_(KBr): 3360 (NH), 1603 (C = O); ¹³C-NMR (CDCl₃): a = 43.4, b = 163.4, c = 87.8, d = 175.5 (J_{FC}_d = 36 Hz), e = 116.8 (J_{FC}_e = 292 Hz). The yield of this compound was 92%.

Received, 29th September, 1981