


SYNTHESIS OF SOME OXAZOLIN-5-ONE-4-SPIRO-1'-CYCLOPROPANES HAVING FUNCTIONAL GROUPS AND THEIR THERMAL REARRANGEMENTS

Otohiko Tsuge*, Michihiko Noguchi, and Hiroyuki Moriyama
 Research Institute of Industrial Science, Kyushu University 86,
 Kasuga, Kasuga-shi, Fukuoka 816, Japan

Abstract — 4-Cycloalkylidene-2-phenyl-2-oxazolin-5-ones reacted with carbonyl-stabilized sulfur methylides to give the corresponding dispiro compounds, which on heating were converted into retro-ene reaction products. The reaction of 4-cinnamylidene-2-phenyl-2-oxazolin-5-one with sulfur allylides afforded the Cope rearrangement products, accompanied by a spiro-cis-divinylcyclopropane in a certain case.

Recently, we have reported the cyclopropanation of 4-methylene-2-phenyl-2-oxazolin-5-ones with dimethylsulfonium phenacylide leading to the formation of two stereoisomeric oxazolin-5-one-4-spiro-1'-cis disubstituted cyclopropanes which are precursors for biologically interesting cyclopropyls of α -amino acids¹. This result is a great contrast to the formation of cis and trans




isomers, or a single cis isomer in the cyclopropanation of 3-arylmethyleneindolin-2-ones², or 4-arylmethylene-2-isoxazolin-5-ones³ with the phenacylide, respectively.

As an extension of the above reaction, we planned to investigate the synthesis of analogous spirocyclopropanes bearing appropriate functional groups favoring a thermal rearrangement of the cyclopropane skeleton, because rearranged products are also possible to be transformed into novel α -amino acid derivatives. In this communication we wish to report the synthesis of some such spirocyclopropanes and their thermal rearrangements.

Oxazolin-5-one-4,1'-spirocyclopropane-2',1"-spirocycloalkanes (dispiro compounds). First, we have investigated the synthesis of the above dispiro compounds by the reaction of 4-cycloalkylidene-

2-phenyl-2-oxazolin-5-ones 1 with carbonyl-stabilized sulfur methylides. 4-Cyclohexylidene- (1a) or 4-cycloheptylidene-2-phenyl-2-oxazolin-5-one (1b)⁴ reacted with dimethylsulfonium phenacylide, generated in situ from dimethylphenacylsulfonium bromide and sodium hydride, in dry THF at 0°C for 5 h to give the corresponding dispiro compound 2a or 2b as the single product in 84 or 35% yield, respectively⁵. Two stereoisomers, 2-1 and 2-2, are possible for the structure of dispiro compound 2. On the basis of the mode of formation as well as the ¹H nmr spectra, however, we assumed that the dispiro compound 2 has the structure 2-1 rather than 2-2. The methine protons of 2a and 2b appeared at δ 3.42 and 3.47, respectively, which are comparable to the value of chemical shift of the methine proton of spirodimethylcyclopropane 3-1 but not that of stereoisomer 3-2¹ (Scheme 1). The previous results also disclosed that in the cyclopropanation of oxazolinones at low temperature spirocyclopropanes such as 3-1 arising from the preferred conformational intermediate were predominantly formed than stereoisomers such as 3-2¹.




Scheme 1

On the other hand, the reaction of 1a with ethyl (dimethylsulfuranylidene)acetate under similar conditions afforded a mixture of two stereoisomeric dispiro compounds, whose ratio was estimated to be ca 3:1 by the nmr spectroscopy, in 97% yield. However, isolation of the minor product was unsuccessful. Although the assignment of stereochemistry of two isomers was very difficult on the basis of their spectral data⁶, it was assumed that the major product is 4 arising from the

preferred conformational intermediate and the minor one is 5.

Since the above dispiro compounds have a cyclopropane moiety in which a carbonyl group (hydrogen acceptor) and ring methylene group (hydrogen donor) are *cis*, they are expected to undergo a retro-ene reaction. On heating in refluxing toluene for 3 h, 2a and 2b were converted into the expected retro-ene reaction products 6a and 6b in 73 and 26% yields, respectively. Structural elucidation of 6a and 6b, 4-(1-cyclohexenyl)- and 4-(1-cycloheptenyl)-4-phenacyl-2-phenyl-2-oxazolin-5-one, was accomplished on the basis of spectral data⁷.

On the other hand, thermolysis of 4 in refluxing xylene for 5 h afforded a 94% yield of a mixture of two isomeric rearrangement products, 7 and 8, which was found to be ca 3:2 by the nmr spectroscopic estimation. Although isolation of each product was unsuccessful, 7 and 8 were assumed to be stereoisomeric 4-(1-cyclohexenyl-ethoxycarbonyl)methyl-2-phenyl-2-oxazolin-5-ones on the basis of spectral data⁸ as well as chemical conversion. When a mixture of 7 and 8 (3:2) was treated with 1M NaOH aqueous solution in refluxing methanol for 7 h, two hydrolyzed products 9, mp 151-152°C, and 10, mp 141-142°C, were obtained in 34 and 21% yields, respectively⁹. However, the stereochemistry of 9 and 10 was not clear.







Scheme 2

As shown in Scheme 2, in the thermolysis of 4 the benzoyl carbonyl group functions as a hydrogen

acceptor to give 6 via A, whereas in 4 the carbonyl group in the oxazolinone ring serves as a hydrogen acceptor to yield isomers 7 and 8 via B.

Oxazolin-5-one-4-spiro-1'-2',3'-cis-divinylcyclopropanes (spiro-cis-divinylcyclopropanes). Next, we have investigated the synthesis of spiro-cis-divinylcyclopropanes which are able to undergo the Cope rearrangement. The reaction of 4-cinnamylidene-2-phenyl-2-oxazolin-5-one (11)¹ with dimethylsulfonium 3-methoxycarbonylallylide¹⁰, generated in situ from 3-methoxycarbonylallyldimethylsulfonium bromide and sodium hydride, in dry THF at room temperature for 3 h afforded a





for 3 h 4%

9%

for 42 h —

32%



Scheme 3

crystalline compound **13** as the sole isolated product in 10% yield. On the basis of spectral data¹¹, the compound **13** was deduced as cyclohepta-1,4-diene-3-spiro-4'-oxazolinone arising from the Cope rearrangement of an initially formed spiro-cis-divinylcyclopropane **12**.

On the other hand, **11** reacted with dimethyloxosulfonium-3-ethoxycarbonyl-2-phenylallylide¹² in dry THF at room temperature for 3 h to give two products **14** and **15**, whereas **15** was only obtained from the reaction for 42 h. On the basis of spectral data¹³, **14** was assigned as the spiro-cis-divinylcyclopropane and **15** as its Cope rearrangement product. In fact, on heating in refluxing benzene for 4 h **14** was transformed into **15** in 50% yield. The assigned structure **14** was strongly supported by comparison with ¹H nmr spectral data of **16-1** and **16-2** obtained from the reaction of **11** with phenacylide¹ (Scheme 3). The configurations of spiro carbons in the Cope rearrangement products **13** and **15** are not clear, although the phenyl and ester groups at 6- and 7-positions are assumed to be trans taking account a concerted Cope rearrangements of **12** and **14**, respectively.

REFERENCES AND NOTES

- O. Tsuge, M. Noguchi, and H. Moriyama, *Heterocycles*, 1981, **16**, 209.
- M. Chiericato, P. D. Croce, and E. Licandro, *J. Chem. Soc. Perkin I*, 1979, 211.
- P. D. Croce, *J. Heterocyclic Chem.*, 1976, **13**, 1109.
- The compound **1a**, mp 141-142°C (lit.¹⁴ mp 137-138°C), was prepared from hippuric acid and cyclohexanone according to the Erlenmeyer method¹⁵. The compound **1b** was prepared from the reaction of the enamine, derived from cycloheptanone and morpholine, with 2-phenyl-2-oxazolin-5-one by the modified Lawson's method¹⁶, since the yield of **1b** according to the Erlenmeyer method was very poor. **1b**: yield 62%; mp 88-89°C; ir (KBr) 1790, 1640 cm⁻¹; MS m/e 255 (M⁺). All new compounds in this paper gave satisfactory elemental analyses.
- 2a**: mp 134-135°C; colorless prisms; ir (KBr) 1800, 1690, 1635 cm⁻¹; ¹H nmr (CDCl₃) δ 3.42 (1H, s, =CH); ¹³C nmr (CDCl₃) δ 46.4 (s), 46.9 (d), 59.3, 161.0, 175.4 (each s); MS m/e 359 (M⁺). **2b**: mp 106-107°C; colorless prisms; ir (KBr) 1790, 1690, 1640 cm⁻¹; ¹H nmr (CDCl₃) δ 3.47 (1H, s, =CH); MS m/e 373 (M⁺).
- 4**: colorless oil; ir (neat) 1805, 1740, 1640, 1635 cm⁻¹; ¹H nmr (CDCl₃) δ 1.27 (3H, t), 2.68 (1H, s, =CH), 4.16 (2H, q). **5**: ¹H nmr (CDCl₃) δ 1.27 (3H, t), 2.86 (1H, s, =CH), 4.20 (2H, q).
- 6a**: mp 166-171°C (dec); colorless needles; ir (KBr) 1820, 1680, 1640 cm⁻¹; ¹H nmr (CDCl₃) δ 4.63 (2H, m, CH₂COPh), 5.67 (1H, m, =CH); ¹³C nmr (CDCl₃) δ 59.2 (t), 64.9 (s), 126.1 (t), 127.9, 162.6, 176.9, 196.3 (each s); MS m/e 359 (M⁺). **6b**: mp 163-165°C; colorless prisms; ir (KBr) 1830, 1675, 1650 cm⁻¹; ¹H nmr (CDCl₃) δ 4.78 (2H, m, CH₂COPh), 5.91 (1H, m, =CH); MS m/e 373 (M⁺).
- A mixture of **7** and **8**: yellow oil; ¹H nmr (CDCl₃) δ 1.16 (t, ca 1.2H), 1.24 (t, ca 1.8H), 3.56

- (1H, d, \approx CH, J=4.0 Hz), 5.15 (1H, dd, \approx CH, J=4.0, 9.0 Hz), 5.49 (1H, m, =CH), 7.60 (1H, d, NH, J=9.0 Hz), 8.29 (1H, broad, OH); MS m/e 345 (M^+). 10: mp 141-142°C; colorless prisms; ir (KBr) 3375, 3200-2300, 1730, 1720, 1620 cm^{-1} ; 1H nmr (DMSO- d_6) δ 1.26 (3H, t), 3.65 (1H, d, \approx CH, J=7.0 Hz), 5.09 (1H, dd, \approx CH, J=7.0, 10.0 Hz), 5.82 (1H, m, =CH), 6.69 (1H, d, NH, J=10.0 Hz), 7.30-7.80 (6H, m, ArH + OH); MS m/e 345 (M^+).
10. J. R. Neff, R. R. Gruetzmacher, and J. E. Nordlauder, *J. Org. Chem.*, 1974, 39, 3814.
11. 13: mp 161-162.5°C; colorless needles; ir (KBr) 1820, 1740, 1645 cm^{-1} ; 1H nmr (CDCl₃) δ 3.71 (3H, s, OCH₃), 4.28 (1H, dddd, \approx CH, J=1.0, 1.3, 4.2, 5.5 Hz), 5.07 (1H, ddd, \approx CH, J=1.9, 4.2, 6.0 Hz), 5.76 (1H, ddd, =CH, J=1.3, 1.3, 11.8 Hz), 5.92 (1H, ddd, =CH, J=1.3, 1.9, 11.0 Hz), 6.02 (1H, dd, =CH, J=5.5, 11.8 Hz), 6.20 (1H, ddd, =CH, J=1.0, 6.0, 11.0 Hz); ^{13}C nmr (CDCl₃) δ 46.4 (s), 51.9 (d), 70.7, 160.5, 171.9, 179.6 (each s); MS m/e 373 (M^+).
12. J. Ide and Y. Kishida, *Tetrahedron Lett.*, 1966, 1787.
13. 14: mp 148-149.5°C; yellow crystals; ir (KBr) 1810, 1710, 1620, 1610 cm^{-1} ; 1H nmr (CDCl₃) δ 0.96 (3H, t), 3.28 (1H, dd, H_b, J=10.0, 10.0 Hz), 3.68 (1H, dd, H_a, J=2.4, 10.0 Hz), 3.86 (2H, q), 5.62 (1H, dd, H_c, J=10.0, 16.2 Hz), 6.42 (1H, d, H_e, J=2.4 Hz), 6.44 (1H, d, H_d, J=16.2 Hz); MS m/e 463 (M^+). 15: mp 155-156°C; pale yellow prisms; ir (KBr) 1815, 1710, 1635, 1620 cm^{-1} ; 1H nmr (CDCl₃) δ 1.06 (3H, t), 2.93 (1H, m, \approx CH), 3.72 (1H, dd, \approx CH, J=2.4, 12.0 Hz), 4.01 (2H, q), 6.32 (1H, d, =CH, J=2.4 Hz), 6.62 (2H, m, =CH); ^{13}C nmr (CDCl₃) δ 40.3, 42.6 (each d), 59.5, 162.1, 165.6, 175.2 (each s); MS m/e 463 (M^+).
14. V. Boekelheide and L. M. Schramm, *J. Org. Chem.*, 1949, 43, 298.
15. H. E. Carter, *Org. React.*, 1946, 3, 198; R. Filler, *Adv. Heterocyclic Chem.*, 1965, 4, 75.
16. A. M. Knowles, A. Lawson, G. D. Boyd, and R. A. Newberry, *J. Chem. Soc. (C)*, 1970, 598.

Received, 14th June, 1982