

OXYGEN-BRIDGED THIA[15]ANNULENES: DEMONSTRATION OF A STRONG PARAMAGNETIC RING CURRENT IN A  $4n\pi$ -THIA-ANNULENE ( $n = 4$ )

Haru Ogawa\* and Taiji Imoto

Faculty of Pharmaceutical Sciences, Kyushu University, Fukuoka 812, Japan

Izumi Miyamoto and Yōichi Taniguchi


Kurume Technical College, Komorino, Kurume 830, Japan

Toshitaka Koga and Yasuyoshi Nogami

Daichi College of Pharmaceutical Sciences, Fukuoka 815, Japan

**Abstract** - The synthesis of 4,7:10,13-diepoxy-2-trans-8,14-di-cis-thia[15]annulene (9) and 4,7:10,13-diepoxy-2,8,14-tri-cis-thia[15]annulene (10) is described. The  $^1\text{H-nmr}$  spectrum of (9) revealed the existence of a paramagnetic ring current, showing that (9) has a planar and diminished bond alternate rigid frame, as expected for an anti-aromatic  $16\pi$  system. Conversely, the  $^1\text{H-nmr}$  spectrum of (10) revealed that (10) is atropic and consequently non-planar.

A hypothetical [15]annulenylium ion system (1) whose geometry is constructed formally with regular pentadecagon seems to offer an angle-strain-free annulene frame, since all the inside and outside angles of (1) are confined to the inside angle of cyclopentadienide anion, viz.  $108^\circ$ . However, replacement of some of the inside hydrogen atoms with divalent atoms is necessary to realize a planar 15-membered molecular framework, because the non-bridged 15-membered ring does not possess a sufficient large cavity as to accommodate all the inner hydrogens without causing serious H-H interactions. This is the reason why that chemistry of pentadecagons made up of  $\text{sp}^2$  carbon skeleton and of its heteroatom incorporated variants has been scarcely studied until oxygen-bridged oxa[15]annulene (2)<sup>1</sup> and [15]annulenones (3 and 4)<sup>2</sup> were prepared as real members. Available findings so far obtained revealed that the oxygen bridges keep the perimeters relatively




flexible as to allow the uni-directional thermal isomerizations both in the [15]annulene (3 → 4) and the corresponding [15]annulenylium ion (5 → 6), respectively,<sup>3</sup> whereas oxa[15]annulene (2) adopts exclusively all *cis* configuration as a thermally rigid frame.

We now report the synthesis of thia[15]annulenes (9 and 10), potential 16 $\pi$  anti-aromatic species, whose preparations were previously unsuccessful by us.<sup>4</sup> In terms of the successful sulphur incorporation, although it be still in lower yields, we became intrigued with following two points:

- (1) Should the thia-annulenes be more paratropic than oxa[15]annulene (2), based on the increasing availability of the sulphur lone pair?
- (2) What arranged geometry should have the thia-annulene, if the accession of the increased paratropicity does hold for the molecule?

The double Wittig reaction of the dialdehyde (7)<sup>5</sup> with the phosphonium salt (8)<sup>6</sup> in DMF, at 60°C, LiOMe as a base under an enforced high-dilution condition<sup>7</sup> followed by a subsequent low temperature work-up, and chromatographic separation on SiO<sub>2</sub> (n-hexane) gave mono-*trans* thia[15]-annulene (9) and all-*cis* counterpart (10) [for physical data, see Table I].



The <sup>1</sup>H-nmr spectrum of (9) indicated that the proton within the perimeter is deshielded, whereas those outside are shielded, i.e., a doublet at  $\delta$  14.59 ppm for the inner proton on the *trans* double bond and a doublet at  $\delta$  5.02 ppm for the outer proton on the *trans* double bond, together with other outer protons at  $\delta$  4.65 - 5.50 ppm (9H) [in CDCl<sub>3</sub> at 25°C, Fig. 1]. Thus it became clear that (9) is strongly paratropic. In contrast, the <sup>1</sup>H-nmr spectrum of (10) consisted only of three signals appearing at the ordinary olefinic region [ $\delta$  6.26, s, 2H (*cis* double bond), 6.54, s, 4H (furanoid H), and 6.57, AB system with  $J/\nu_0\delta > 5^{\text{eq}}$  (*cis* double bonds adjacent to the S-atom)], indicating that (10) is atropic and consequently non-planar.

A comparison of the chemical shifts of the fifteen membered ring protons in the related system is shown in Table II. Crucial role of the sulphur atom for the demonstration of paratropic

character became evident in (9). A confirmative evidence was given, which implies that forcible participation of the sulphur lone pair occurred only in the mono-trans geometry, not in the all-cis geometry. Model examination shows that an extensive twisting of the cis double bonds of (10) hinders the effective overlap of the sulphur lone pair. The difference of chemical shifts of the inner and outer proton resonances on the trans double bond of (9) [ $\Delta\delta = \delta_i - \delta_{o_{av}} = 9.6$  ppm] gave the largest value yet recorded in neutral hetero[15]annulenes so far prepared, and the value is close to the corresponding value obtained from di-oxygen-bridged [16]-annulene dioxide (11)<sup>9</sup> [ $\Delta\delta = 12.8$  ppm], an iso-electronic  $16\pi$  annulene frame possessing same number of oxygen-bridge [see Table II].




Fig. 1.  $^1\text{H-Nmr}$  Spectrum of (9) in  $\text{CDCl}_3$  at  $25^\circ\text{C}$






In agreement with the large paratropic shift of (9), electronic spectrum of (9) in *n*-hexane showed a very close resemblance in shape with that of (11)<sup>9</sup> [Fig. 2]. Indeed, the position of the longest wavelength bands of (9) and (10), which appeared at 503 nm ( $\log \epsilon = 2.57$ ) and at 530 nm ( $\log \epsilon = 2.43$ ), respectively, indicate clearly the participation of lone pair electrons of sulphur atom. The  $^1\text{H-nmr}$  and electronic spectral properties of (9) reveal that the molecule should be planar with delocalized  $16\pi$ -electron system. In contrast, the electronic spectrum of (10) remains markedly different from those of (9) and (11).

It was found that no thermal isomerization occurred in both thia-annulenes, since the  $^1\text{H-nmr}$  spectra did not show significant changes in  $\text{DMF-d}_7$ , up to  $150^\circ\text{C}$ . In contrast, a facile isomerization (9)  $\rightarrow$  (10) occurred in the presence of  $\text{H}^+$  ( $\text{CF}_3\text{COOH}$ ) at room temperature (uv and  $^1\text{H-nmr}$  spectroscopy).

Interestingly, neither the sulphone (12) nor sulphoxide (13) [obtainable by the oxidation of (9) and (10) with *m*-chloroperbenzoic acid in  $\text{CH}_2\text{Cl}_2$  at  $-20^\circ\text{C}$  for 2 h, respectively] showed any

tendency to exhibit diatropicity ( $14\pi$ ), in agreement with the theoretical prediction, which implies that ring current intensity is substantially more higher for delocalized  $4n\pi$  compounds than  $(4n+2)\pi$  compounds with same size, if pronounced bond alternation does not occur in the  $4n\pi$  system.<sup>a,b,ec</sup>

Table II.  $^1\text{H-Nmr}$  Parameters of the Oxygen-bridged  $16\pi$ - and related Compounds in  $\text{CDCl}_3$ , ( $\delta$  values in ppm,  $J$  values in Hz, at  $25^\circ\text{C}$ )

Compound	$\text{H}_a$	$\text{H}_{\text{trans}}$	$\text{H}_{\text{cis}}$	$\text{H}_{\text{furan}}$
 (2)	5.20, s		4.78 d, $J=6.4$ 5.56 d, $J=6.4$	5.62 d, $J=3.4$ 5.73 d, $J=3.4$
 (9)	5.00, s	5.20 d, $J=15.2$ 14.55 d, $J=15.2$	4.65 d, $J=13.5$ 5.14 d, $J=13.5$	5.23 d, $J=3.4$ 5.35 d, $J=3.4$ 5.45 d, $J=3.4$ 5.50 d, $J=3.4$
 (10)	6.26, s		6.57 AB system ( $J/\nu_0\delta > 5$ )	6.54, s
 (11)		4.43 d, $J=15.0$ 17.18 d, $J=15.0$	4.41 d, $J=10.0$ 4.60 d, $J=10.0$	4.86 d, $J=3.5$ 4.93 d, $J=3.5$
 (14)	6.40, s		5.97 d, $J=12.5$ 6.50 d, $J=12.5$	6.54 d, $J=3.5$ 6.65 d, $J=3.5$




Fig. 2. Electronic Spectra of (9) ———, (10) ·····, and (11) - - - - in n-hexane

Table I. Physical Data of Compounds

Thia[15]annulene dioxide (9): mp 98 - 100°C (n-hexane); yield 1.6 %; MS: m/e 242 ( $M^+$ ); electronic spectrum  $\lambda_{\max}$  (n-hexane): 227 (10700), 280.5 (49100), 336 sh (1700), 350 sh (1000), 380 sh (480), 503 (370);  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ ) see Text;  $^{13}\text{C-nmr}$  ( $\text{CDCl}_3$ ) 106.5 (d), 110.9 (d), 114.4 (d), 114.5 (d), 115.8 (d), 116.2 (d), 116.5 (d), 117.2 (d), 120.4 (d), 129.8 (d); signals due to the four singlet carbons on the furan rings were obscured in the  $^{13}\text{C-nmr}$  spectrum; ir (KBr) 1600, 1038, 1020, 933  $\text{cm}^{-1}$ .

Thia[15]annulene dioxide (10): mp 92 - 93°C (n-hexane); yield 5.5 %; MS: m/e 242 ( $M^+$ ); electronic spectrum  $\lambda_{\max}$  (n-hexane): 228 (31400), 265 (23450), 271 (22500), 296 (31700), 425 (2750);  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ ) see Text;  $^{13}\text{C-nmr}$  ( $\text{CDCl}_3$ ) 151.4 (s), 149.9 (s), 120.8 (d), 114.4 (d), 113.5 (d), 111.4 (d); ir (KBr) 1330, 1182, 1030  $\text{cm}^{-1}$ .

The sulphone (12): mp 130°C (decomp) (EtOH); yield 75 %; MS: m/e 274 ( $M^+$ );  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ ) 6.12 (d, J = 15 Hz, 1H), 6.17 (d, J = 13.5 Hz, 1H), 6.66 (s, 2H), 6.76 (d, J = 3.2 Hz, 1H), 6.78 (d, J = 13.5 Hz, 1H), 6.83 (d, J = 3.2 Hz, 1H), 6.93 (d, J = 3.2 Hz, 1H), 7.04 (d, J = 3.2 Hz, 1H), 7.57 (d, J = 15 Hz, 1H).

The sulphoxide (13): mp 188 - 190°C (EtOH); yield 90 %; MS: m/e 258 ( $M^+$ );  $^1\text{H-nmr}$  ( $\text{CDCl}_3$ ) 7.08 (s, 2H), 7.09 (d, J = 11 Hz, 2H), 7.26 (d, J = 3.5 Hz, 2H), 7.33 (d, J = 3.5 Hz, 2H), 7.46 (d, J = 11 Hz, 2H).

## REFERENCES AND NOTES

1. H. Ogawa, M. Kubo, and H. Saikachi, *Tetrahedron Lett.*, 4859 (1971).
2. H. Ogawa, N. Shimojo, H. Kato, and H. Saikachi, *Tetrahedron*, **30**, 1033 (1974).
3. H. Ogawa, N. Kariya, T. Imoto, and Y. Taniguchi, *Croatia Chemica Acta*, **53**, 637 (1980).
4. H. Ogawa and M. Kubo, *Tetrahedron*, **29**, 809 (1973).
5. H. Saikachi, H. Ogawa, and K. Sato, *Chem. Pharm. Bull.*, **19**, 97 (1971).
6. K. Dimroth, H. Follmann, and G. Pohl, *Chem. Ber.*, **99**, 642 (1966).
7. To a solution of (7) (1.62 g, 7.5 mM) and (8) (5.58 g, 7.5 mM) in dry DMF (240 ml) was added a solution of LiOMe (208 mg of lithium in 30 ml of absolute MeOH) dropwisely at 60°C over 7 h under nitrogen.
8. H. Günther, "NMR Spectroscopy", John Wiley and Sons, 1980, (a) p. 152; (b) pp. 83 - 86; (c) J.A. Pople and K.G. Untch, *J. Am. Chem. Soc.*, **88**, 4811 (1966).
9. H. Ogawa, C. Fukuda, T. Imoto, I. Miyamoto, and Y. Taniguchi, *Angew. Chem.*, in submission, 1982.

Received, 8th July, 1982