

SYNTHESIS OF 13-AZA-16-OXA-, 13-AZA-16-THIA- AND 13,16-DIAZASTEROIDAL COMPOUNDS THROUGH CYCLIZATION OF N-ACYL-IMINIUM ION INTERMEDIATES


Shinzo Kano* and Yoko Yuasa

Tokyo College of Pharmacy, 1432-1 Horinouchi, Hachioji,
Tokyo 192-03, Japan

Abstract — 5,5-Dimethyl-2,4-oxazolidinedione, 2,4-thiazolidinedione and 5,5-dimethylhydantoin were coupled with 2-(3,4-dihydro-1-naphthyl)ethyl alcohol by an application of Mitsunobu's method to give the corresponding N-substituted products (5)-(7), respectively. Reduction of 5-7 with diisobutylaluminium hydride, followed by cyclization of the reduction products with formic acid yielded the 13-aza-16-oxa- (11), 13-aza-16-thia- (12) and 13,16-diaza-1,3,5(10),9(11)-estratetraen-17-one (13), respectively.


A great deal of synthetic efforts to synthesize heterocyclic steroidal compounds have been made from chemical and biological points of view¹. Much attention has been devoted to a synthesis of a number of azasteroids^{1a,2}. 13-Azasteroids (1)-(3) were prepared via N-acyliminium-induced C-C bond formation³ (Chart 1).

Chart 1



In a similar way, synthesis of some 13-azasteroids with a modification of the A- or B-ring have been reported⁴. In view of the current interest in an efficient synthesis of heterocyclic steroidal systems with a modification of the D-ring, we examined the synthesis of 13-azasteroids through ring closure of N-acyliminium ion intermediates^{3,4}. The results of our studies will be described in this paper. N-[2-(3,4-Dihydro-1-naphthyl)ethyl]-2,4-oxazolidinedione, 2,4-thiazolidinedione and hydantoin derivatives were used as the starting materials for this purpose because of easy handling and ready conversion to the desired N-acyliminium ions. 2-(3,4-Dihydro-1-naphthyl)ethyl alcohol (4)⁵ was coupled with 5,5-dimethyl-2,4-oxazolidinedione, 2,4-thiazolidinedione, and 5,5-dimethylhydantoin by the modified Mitsunobu's method⁶ using diisopropyl azodicarboxylate instead of diethyl azodicarboxylate to give the corresponding N-substituted products (5), (6) and (7), respectively. Reduction of 5 with diisobutylaluminum hydride, followed by treatment of the resulting 5-hydroxy derivative (8) with formic acid at room temperature for 1 h gave 3-deoxy-18-nor-15,15-dimethyl-13-aza-16-oxa-1,3,5(10),9(11)-estratetraen-17-one (11) in 65 % yield, mp 114-115°C, the structure of which was confirmed by the spectroscopic method together with decoupling technique in its ¹H NMR spectra. In this reaction, neither the $\Delta^{8,9}$ -isomer (14) nor the 9-oxygenated product (15) was obtained (Chart 2). In a similar fashion, 13-aza-16-thia (12) and 13,16-diaza derivatives (13) were easily obtained from 9 and 10,

Chart 2



which were derived from 6 and 7, respectively. Thus, the synthesis of heterocyclic steroidal systems with D-ring modification was achieved by Speckamp's method³.

EXPERIMENTAL

All melting points are not corrected. ¹H NMR spectra were determined on a Varian EM-390 spectrometer operating at 90 MHz by using CDCl₃ as a solvent and tetramethylsilane as an internal standard. Mass spectra were taken with a Hitachi RMU-7L instrument.

N-[2-(3,4-Dihydro-1-naphthyl)ethyl]-5,5-dimethyl-2,4-oxazolidinedione (5) To a stirred mixture of 4 (3.48 g, 20 mmol), triphenylphosphine (6.29 g, 24 mmol), 5,5-dimethyl-2,4-oxazolidinedione (3.10 g, 24 mmol) and THF (30 ml) was added a solution of diisopropyl azodicarboxylate (4.85 g, 24 mmol) in THF (20 ml) under ice-cooling. After the stirring had been continued for 14 h at room temperature, the solvent was evaporated. The remaining residue was chromatographed on silica gel (25 g) by using benzene-hexane (1:2) as an eluent. Evaporation of the solvent (80-100 ml) gave 5 (3.71 g, 65 % yield) as an oil; ¹H NMR (CDCl₃) δ 1.47 (6H, s), 1.73-2.72 (4H, m), 2.80 (2H, t, J=8 Hz), 3.70 (2H, t, J=8 Hz), 5.94 (1H, t, J=5 Hz), 7.03-7.20 (4H, m); Mass spectrum m/e 285.1366 (M⁺, calcd. for C₁₇H₁₉NO₃ m/e 285.1364)

N-[2-(3,4-Dihydro-1-naphthyl)ethyl]-2,4-thiazolidinedione (6) This compound was obtained by the same procedure as above except the use of 2,4-thiazolidinedione (2.81 g, 24 mmol) instead of 5,5-dimethyl-2,4-oxazolidinedione (3.10 g) in 75 % yield (4.1 g) as an oil; ¹H NMR (CDCl₃) δ 1.77-2.33 (4H, m), 2.71 (2H, t, J=8 Hz), 3.77 (2H, s), 3.80 (2H, t, J=8 Hz), 5.60 (1H, t, J=5 Hz), 7.04-7.29 (4H, m); Mass spectrum m/e 273.0819 (M⁺, calcd. for C₁₅H₁₅NO₂S m/e 273.0822).

N-[2-(3,4-Dihydro-1-naphthyl)ethyl]-5,5-dimethylhydantoin (7) This compound was obtained by the same procedure as above except the use of 5,5-dimethylhydantoin (3.07 g, 24 mmol) instead of 5,5-dimethyl-2,4-oxazolidinedione (3.10 g) in 72 % yield (4.09 g) as an oil; ¹H NMR (CDCl₃) δ 1.34 (6H, s), 1.70-2.33 (2H, m), 2.63-2.87 (4H, m), 3.70 (2H, t, J=8 Hz), 5.96 (1H, t, J=5 Hz), 7.03-7.20 (4H, m); Mass spectrum m/e 284.1521 (M⁺, calcd. for C₁₇H₂₀N₂O₂ m/e 284.1523).

Preparation of 11 To a stirred solution of 5 (1.5 g, 5.26 mmol) in toluene (30 ml) was added diisobutylaluminium hydride (1.49 g, 6.87 ml of 25 % toluene solution: 10.52 mmol) at -78°C. After the stirring had been continued for 1 h at the same temperature, the mixture was decomposed with 5 % H₂SO₄ (60 ml) and

extracted with CHCl_3 . The extract was washed with water, dried (Na_2SO_4) and evaporated. The resulting residue was mixed with formic acid (20 ml) under stirring at room temperature. After the stirring had been continued for 1 h, the mixture was made basic with 28 % ammonia and extracted with CHCl_3 . The extract was washed with water, dried (Na_2SO_4) and evaporated to give 11 (0.92 g, 65 % yield), mp 114-115°C (methanol-ether); $^1\text{H NMR}$ (CDCl_3) δ 1.47 (3H, s), 1.64 (3H, s), 3.26 (1H, d, $J=10$ Hz), 3.89 (1H, t, $J=2$ and 18 Hz), 4.28 (1H, t,d, $J=2$ and 18 Hz), 6.33-6.44 (1H, m), 7.16-7.33 (3H, m), 7.61-7.84 (1H, m); Mass spectrum m/e 269 (M^+). Anal. Calcd for $\text{C}_{17}\text{H}_{19}\text{NO}_2$: C, 75.81; H, 7.11; N, 5.20. Found: C, 75.60; H, 7.10; N, 5.16.

Preparation of 12 The compound (6) (1.5 g, 5.49 mmol) was reduced with diisobutylaluminum hydride (1.56 g, 7.17 ml of 25 % toluene solution: 10.98 mmol) by the same procedure as above and resulting reduction product was mixed with formic acid (20 ml) and worked up as above to yield 12 (1.2 g, 85 % yield), mp 181-182°C (methanol-ether); $^1\text{H NMR}$ (CDCl_3) δ 3.82 (1H, t,d, $J=2$ and 18 Hz), 4.45 (1H, t,d, $J=2$ and 18 Hz), 6.26-6.37 (1H, m), 7.12-7.26 (3H, m), 7.53-7.68 (1H, m); Mass spectrum m/e 257 (M^+). Anal. Calcd for $\text{C}_{15}\text{H}_{15}\text{NOS}$: C, 70.01; H, 5.88; N, 5.44. Found: C, 69.88; H, 5.84; N, 5.27.

Preparation of 13 The compound (7) (1.5 g, 5.28 mmol) was reduced with diisobutylaluminum hydride (1.50 g, 6.90 ml of 25 % toluene solution: 10.56 mmol) by the same procedure as above and resulting reduction product was mixed with formic acid (20 ml) and worked up as above to yield 13 (1.25 g, 88 % yield), mp 80-83°C (methanol-ether); $^1\text{H NMR}$ (CDCl_3) δ 1.30 (3H, s), 1.44 (3H, s), 3.77 (1H, t,d, $J=2$ and 18 Hz), 4.26 (1H, t,d, $J=2$ and 18 Hz), 6.37-6.48 (1H, m), 7.20-7.33 (3H, m), 7.63-7.86 (1H, m); Mass spectrum m/e 268 (M^+). Anal. Calcd for $\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}$: C, 76.08; H, 7.51; N, 10.44. Found: C, 75.80; H, 7.29; N, 10.16.

REFERENCES AND NOTES

- (a) H. O. Huisman, Angew. Chem. Int. Ed. Eng., 10, 450 (1971); (b) P. Morand and J. Lyall, Chem. Rev., 68, 85 (1968); (c) S. R. Ramada, P. C. Chenciah, N. S. C. Kumar, M. V. Krishna, P. S. Srinivasan, V. V. K. Sastry, and J. A. Rao, Heterocycles, 19, 861 (1982); (d) R. T. Blickenstaff, A. C. Ghosh, and G. C. Wold, "Total Synthesis of Steroids", Academic Press, New York, 1974.
- For instance; (a) J. C. Hubert, W. N. Speckamp, and H. O. Huisman, Tetrahedron Lett., 1969, 1553; (b) W. N. Speckamp, R. J. P. Barends, A. J. de Gee, and

- H. O. Huisman, Tetrahedron Lett., 1970, 383; (c) K. Matoba, E. Ishigami, H. Takahata, A. Fukushima, and T. Yamazaki, Chem. Pharm. Bull., 29, 651 (1981); (d) K. Matoba, A. Fukushima, H. Takahata, Y. Hirai, and T. Yamazaki, Chem. Pharm. Bull., 30, 1300 (1982).
3. (a) J. C. Hubert, W. N. Speckamp, and H. O. Huisman, Tetrahedron Lett., 1972, 4493; (b) J. Dijkink and W. N. Speckamp, Tetrahedron Lett., 1977, 935; (c) J. Dijkink and W. N. Speckamp, Tetrahedron, 34, 173 (1978); (d) J. A. M. Hamersma and W. N. Speckamp, Tetrahedron, 38, 3255 (1982).
4. (a) R. R. Trehan, E. Bala, D. K. Trikhara, and J. Singh, Indian J. Chem., 20B, 1022 (1981); (b) I. R. Trehan, J. K. Battu, R. L. Kaul, and J. Singh, ibid., 20B, 1026 (1981); (c) I. R. Trehan, K. Bala, and J. Singh, ibid., 18B, 295 (1979); (d) I. R. Trehan, R. L. Kaul, R. K. Sharma, and J. Singh, ibid., 21B, 197 (1982); (e) I. R. Trehan, J. Ghose, and J. Manju, ibid., 19B, 243 (1980).
5. W. R. Schleigh, A. Catala, and F. D. Popp, J. Heterocyclic Chem., 2, 379 (1965).
6. O. Mitsunobu, W. Wada, and T. Sano, J. Am. Chem. Soc., 94, 679 (1972).

Received, 19th January, 1983