

A SHORT STEP SYNTHESIS OF LESPEDAMINE¹

Masanori Somei,^{*} Haruhiko Sato, and Chikara Kaneko
 Faculty of Pharmaceutical Sciences, Kanazawa University
 13-1 Takara-machi, Kanazawa 920, Japan

Abstract— A convenient synthetic method for 1-hydroxy-, 1-methoxy-, and 1-acetoxy-2-oxindole was disclosed starting from methyl 2-nitrophenylacetate. A five-step synthesis of lespedamine was achieved utilizing this method.

In this report, we describe a practical synthetic method for 1-hydroxy-2-oxindole derivatives and a short step synthesis of lespedamine (1),^{2,3} one of eight naturally occurring 1-methoxyindole derivatives.⁴

I. Syntheses of 1-Hydroxy-2-oxindole Derivatives


Various synthetic methods so far reported for 1-hydroxy- (2) and 1-methoxy-2-oxindole (3) are known to give unsatisfactory results.⁵ However, we found that the readily available methyl 2-nitrophenylacetate (4) simply upon treatment with zinc (20 mol eq.) and ammonium chloride (3.8 mol eq.) in methanol for 3h afforded 2⁶ in 48% yield. When an excess amount of reducing agents was used or a longer reaction time was adopted, the yield of 2 was decreased mainly due to its sensitivity toward reductive decomposition, resulting in the formation of 2-oxindole.

We have also found that a significant amount of 2 was lost by the formation of a complex⁷ with zinc iron, which was rather insoluble in organic solvents. The structure of the complex was tentatively assigned to be 5 based mainly on its mass spectrum which showed the ratio of 2 and zinc iron to be 2 to 1.

Direct treatments of the reaction mixture, obtained by the reaction of 4 with zinc and ammonium chloride, with diazomethane and acetic anhydride and pyridine were found to give 1-methoxy- (3)⁸ and 1-acetoxy-2-oxindole (6)⁹ in 77% and 70% overall yields, respectively. Furthermore, hydrolysis of 6 with sodium carbonate afforded 2 in 94% yield.

II. Synthesis of Lespedamine

The reaction of 3 with ethylene dibromide in the presence of sodium hydride afforded spiro compound (7)¹⁰ in 90% yield. Subsequent treatment of 7 with aq. dimethylamine (50 mol eq.) and its hydrochloride (9 mol eq.) in *N,N*-dimethylformamide produced the desired 3-(2-*N,N*-dimethylaminoethyl)-1-methoxy-2-oxindole (8)¹¹ in 54% yield, together with 10% yield of a phenylhydroxylamine derivative (9).¹² The reduction of 8 with lithium aluminum hydride (LiAlH_4) in ether was found to produce 3-(2-*N,N*-dimethylaminoethyl)-2-hydroxy-1-methoxy-2,3-dihydroindole (10)¹³ in 62% yield as a mixture of diastereoisomers. The compound (10) was unstable and instantaneously changed by treatment with aq. hydrochloric acid to lespedamine¹⁴ (1) in 95% yield. On the basis of the above results, the final step was improved as follows. Thus, after reduction of 8 with LiAlH_4 , the reaction mixture was treated briefly with aq. hydrochloric acid. By this modification, lespedamine (1) was prepared in 64% yield directly from 8. Thus, the total synthesis³ of 1 was achieved in five steps with 24% overall yield from 4. The spectral data of synthetic material and melting point of its picrate were identical with those of lespedamine.² In conclusion, building blocks such as 2, 3, and 6 for 1-hydroxyindole derivatives have now become readily available from 4 in excellent yields. Investigation of their reactions and preparation of other naturally occurring 1-methoxyindole derivatives are currently in progress.



REFERENCES AND NOTES

1. Presented partly at the 103rd Annual Meeting of the Pharmaceutical Society of Japan, Tokyo, April, 1983. This report is part XX of a series entitled "The Chemistry of Indoles." Part XIX: M. Somei, S. Tokutake, and C. Kaneko, Chem.

- Pharm. Bull., in press.
- H. Morimoto and H. Oshio, Liebigs Ann. Chem., **682**, 212 (1965).
 - Total synthesis of lespedamine was reported with 2.6% overall yield. R.M. Acheson, P.G. Hunt, D.M. Littlewood, B.A. Murrer, and H.E. Rosenberg, J. Chem. Soc., perkin I, 1978, 1117.
 - E. Wenkert, J.C. Orr, S. Garratt, J.H. Hansen, B. Wickberg, and C.L. Leicht, J. Org. Chem., **27**, 4123 (1963); S.R. Johns, J.A. Lamberton, and J.L. Occolowitz, Aust. J. Chem., **20**, 1737 (1967); M. Nomoto and S. Tamura, Agr. Biol. Chem., **34**, 1590 (1970); D.W. Nagel, K.G.R. Pachler, P.S. Steyn, R. Vleggaar, and P.L. Wessels, Tetrahedron, **32**, 2625 (1976); H. Wagner and T. Nestler, Tetrahedron Letters, 1978, 2777; Y. Konda, M. Onda, A. Hirano, and S. Omura, Chem. Pharm. Bull., **28**, 2987 (1980).
 - A. Reissert, Ber., **41**, 3921 (1908); W.B. Wright, Jr. and K.H. Collins, J. Am. Chem. Soc., **78**, 221 (1956); T. Andre, M. Georges, and R. Gilbert, C. R. Acad. Sci., Ser. C, **273**, 1378 (1971) [C.A., **76**, 59359w (1972)]; C.W. Muth and R.N. Beers, Proc. W. Va. Acad. Sci., **41**, 235 (1969) [C.A., **74**, 42104v (1971)].
 - mp 200.5-202.0°C (lit.⁵ mp 199-200°C). IR (KBr): 1675, 1617 cm⁻¹. ¹H-NMR (10% CD₃OD in CDCl₃) δ: 3.35 (1H, br s), 3.43 (2H, s), 6.65-7.41 (4H, m).
 - mp >300°C. IR (KBr): 1630, 1605 cm⁻¹. ¹H-NMR (pyridine-d₅) δ: 3.28 (4H, s), 6.62-7.38 (8H, m). High MS m/z: Calcd for C₁₆H₁₂N₂O₄Zn: 360.0087 and 362.0057. Found: 360.0109 and 361.9963.
 - mp 84.5-86.0°C (lit.⁵ mp 84-86°C). IR (KBr): 1712, 1617 cm⁻¹. ¹H-NMR (CDCl₃) δ: 3.42 (2H, s), 3.95 (3H, s), 6.65-7.42 (4H, m). MS m/z: 163 (M⁺), 132.
 - mp 97-99°C. IR (KBr): 1807, 1727 cm⁻¹. ¹H-NMR (CDCl₃) δ: 2.33 (3H, s), 3.55 (2H, s), 6.50-7.35 (4H, m). Anal. Calcd for C₁₀H₉NO₃: C, 62.82; H, 4.75; N, 7.33. Found: C, 63.00; H, 4.72; N, 7.04.
 - Oil. IR (film): 1723, 1619 cm⁻¹. ¹H-NMR (CCl₄) δ: 1.17-1.54 (2H, m), 1.54-1.87 (2H, m), 3.92 (3H, s), 6.41-7.21 (4H, m). High MS m/z: Calcd for C₁₁H₁₁NO₂: 189.0789. Found: 189.0795
 - Oil. IR (film): 1727, 1616 cm⁻¹. ¹H-NMR (CCl₄) δ: 1.69-2.50 (4H, A₂B₂, m), 2.06 (6H, s), 3.32 (1H, t, J=5.6 Hz), 3.86 (3H, s), 6.57-7.29 (4H, m). High MS m/z: Calcd for C₁₃H₁₈N₂O₂: 234.1367. Found: 234.1375.
 - Oil. IR (film): 3480, 1647 cm⁻¹. ¹H-NMR (CDCl₃) δ: 1.67-2.57 (4H, m), 2.32 (6H, s), 2.73 (3H, s), 2.86 (3H, s), 3.66 (3H, s), 3.90 (1H, dd, J=8.8 and 5.2 Hz), 6.50-7.30 (4H, m), 6.93 (1H, br s). High MS m/z: Calcd for C₁₅H₂₅N₃O₂: 279.1944. Found: 279.1937.
 - Oil. IR (film): 3340, 1612, 1596, 1475, 1463 cm⁻¹. ¹H-NMR (CCl₄) δ: 1.84-2.75 (5H, m), 2.25 (6H, s), 3.82 (3H, s), 4.60 and 4.92 (total 1H, each d, J=8 Hz), 5.83 (1H, br s), 6.44-7.15 (4H, m). High MS m/z: Calcd for C₁₃H₂₀N₂O₂: 236.1523. Found: 236.1539.
 - Spectra of IR and ¹H-NMR were identical with those of lespedamine. Charts of IR and ¹H-NMR spectra of lespedamine are reported in the ref. 2. Oil. IR (CHCl₃): 1459 cm⁻¹. ¹H-NMR (CCl₄) δ: 2.19 (6H, s), 2.32-2.96 (4H, m), 3.92 (3H, s), 6.62-7.45 (5H, m). MS m/z: 218 (M⁺), 187 (M⁺-OMe). Picrate: mp 161-163°C (lit.² mp 160-162°C).

Received, 9th May, 1983