

THE ACIDITIES AND THE TAUTOMERIC STRUCTURE OF 5-ARYL-2-MERCAPTO-1,3,4-OXADIAZOLES

Ahmad S. Shawali,*† Mahmoud S. Rizk, Abdou O. Abdelhamid,* and Magda A. Abdalla
 Department of Chemistry, Faculty of Science, University of Cairo,
 Giza, Egypt

Cyril Párkányi* and Magdalena E. Wojciechowska§
 Department of Chemistry, The University of Texas at El Paso,
 El Paso, Texas 79968, U.S.A.

Abstract - The acidity constants of a series of substituted 5-phenyl-2-mercapto-1,3,4-oxadiazoles were determined by potentiometric and spectrophotometric methods in 80% (vol.) ethanol-water at 25°C. The data obtained by the two methods are in good agreement. The pK_a values correlate with the σ^* constants of the substituted phenyl group XC_6H_4 ($\rho = -0.985$, $r = 0.936$), however, a better correlation of the pK_a data with the Hammett substituent constants σ_X ($\rho = -0.983$, $r = 0.959$) was obtained. These linear correlations exclude the possibility of the presence of the thiol tautomer $\underline{1}$ in equilibrium with the thioamide tautomer $\underline{2}$ and indicate that the ionization of the compounds takes place in the form of the thioamide tautomer $\underline{2}$. According to the results of the HMO calculations, the thioamide form $\underline{2}$ is more stable than the thiol tautomer $\underline{1}$. A satisfactory correlation between the observed pK_a values and the difference between the π -electronic energies (ΔE_π) of the thioamide tautomer and of the common resonance-stabilized anion was obtained, thus supporting the assigned tautomeric structure $\underline{2}$ for the series under study.

†Present address: Department of Chemistry, Faculty of Science, King Abdul Aziz University, P. O. Box 9028, Jeddah, Saudi Arabia.


*Present address: Department of Chemistry, The University of Texas at El Paso, El Paso, Texas 79968, U.S.A.

§Present address: Physical Science Department, Troy State University, Troy, Alabama 36081, U.S.A.


INTRODUCTION

Although several studies concerning the transmission of substituent effects by heterocyclic rings have been reported,¹⁻³ the 1,3,4-oxadiazole ring, to our knowledge, has not been investigated from this point of view so far.

In the present contribution, the acid dissociation constants of a series of 5-aryl-2-mercapto-1,3,4-oxadiazoles $\overset{1a-1j}{\sim\sim}$ ($R = XC_6H_4$) have been determined in 80% (vol.) ethanol-water at 25°C and the results have been correlated by using the Hammett equation. The pK_a values for only three compounds, $\overset{1a}{\sim}$ ($R = Ph$), $\overset{1k}{\sim}$ ($R = H$), and $\overset{1l}{\sim}$ ($R = Me$) have been reported in the literature.^{4,5}



$\overset{1}{\sim}$



$\overset{2}{\sim}$


The test of the applicability of the Hammett equation to the effective ionization constants of $\overset{1}{\sim}$ is also of interest because it will help to obtain further insight into the possible tautomerism between $\overset{1}{\sim}$ and $\overset{2}{\sim}$. On the basis of the spectral data, these compounds were reported to exist in their thioamide form $\overset{2}{\sim}$ in solution.^{4,5}

Furthermore, it seemed worthwhile to study the acidity of a systematic series of compounds of type $\overset{1}{\sim}$ from a quantum-chemical viewpoint. In the present work, the HMO π -electronic energies of the two tautomers $\overset{1}{\sim}$ and $\overset{2}{\sim}$ and of their common anion $\overset{3}{\sim}$, $E_{\pi}(\overset{1}{\sim})$, $E_{\pi}(\overset{2}{\sim})$, and $E_{\pi}(\overset{3}{\sim})$, respectively, have been calculated⁶ and correlations between the pK_a data and each of the energy differences [$E_{\pi}(\overset{3}{\sim}) - E_{\pi}(\overset{1}{\sim})$ and $E_{\pi}(\overset{3}{\sim}) - E_{\pi}(\overset{2}{\sim})$] were examined.

RESULTS AND DISCUSSION

The reaction sequence followed in the synthesis of the compounds $\overset{1a-1j}{\sim\sim}$ is shown in Scheme 1.⁷ Five of the 1,3,4-oxadiazoles prepared are new compounds and their structures have been confirmed on the basis of their elemental and spectral analyses (see Experimental).

The compounds $\overset{1a-1j}{\sim\sim}$ behave as weak acids in 80% (vol.) ethanol-water at 25°C. If the prototropic tautomerism is taken into account, the ionization of $\overset{1a-1j}{\sim\sim}$ can be represented by Scheme 2. The effective ionization constants, pK_a , of these compounds were determined both by spectroscopic and by potentiometric titration methods. The uv spectra of $\overset{1}{\sim}$, in each case, vary with increasing pH of the medium (Fig. 1). For all the compounds under study, the dependence of the absorbance on the pH value gives sigmoid curves. Similarly, the potentiometric titration curves were sigmoid as well. Typical plots are shown in Fig. 2.



a, X = H

b, X = p-MeO

c, X = p-Me

d, X = m-Me

e, X = p-Cl


f, X = m-Cl

g, X = p-Br


h, X = m-Br

i, X = m-NO₂

j, X = p-NO₂



Scheme 1



Scheme 2




Fig.1. Ultraviolet absorption spectra of 5-(p-nitrophenyl)-2-mercapto-1,3,4-oxadiazole at different pH values; $c = 5.0 \times 10^{-5} M$, in 80% (vol.) ethanol-water at 25°C and $\mu = 0.10$.




Fig. 2. Typical spectrophotometric titration curve of $1j$ (A) at $\lambda = 345$ nm and potentiometric titration curves of $1b$ (B) and $1a$ (C).

The pK_a values were calculated from the absorbance-pH data by eq. [1].

$$pK_a = pH_i + \log \frac{A_b - A_j}{A_j - A_a} \quad [1]$$

where A_j is the absorbance of the test solution at pH_i , and A_b and A_a are the absorbance values of the strongly alkaline and strongly acid solutions, respectively. In the potentiometric method, the pK_a values were calculated by using eq. [2].

$$K_a = (a I_1 + [H^+])[H^+]/I_1 (1 - a) - [H^+] \quad [2]$$

where a is the molar ratio of the base added to the compound, and I_1 is the total acid concentration. It should be noted that the pK_a values from potentiometric titrations agree well with the data obtained spectrophotometrically. The pK_a values of the compounds studied are summarized in Table 1.

Table 1. Experimental and Calculated Quantities for Substituted
5-Aryl-2-mercapto-1,3,4-oxadiazoles, 1a-1j

| Compound | Substituent | pK_a (a) | σ_X | $\sigma_{XC_6H_4}^*$ | E_π (β) | | |
|-----------|---------------------------|------------|------------|----------------------|---------------------|--------|--------|
| | | | | | Form | | |
| | | | | | 1 | 2 | 3 |
| <u>1a</u> | None | 5.25 | 0.00 | 0.58 | 20.813 | 21.672 | 23.976 |
| <u>1b</u> | <u>p</u> -MeO | 5.29 | -0.27 | 0.36 | 23.995 | 25.875 | 28.127 |
| <u>1c</u> | <u>p</u> -Me | 5.49 | -0.17 | 0.47 | 23.516 | 25.393 | 27.673 |
| <u>1d</u> | <u>m</u> -Me | 5.40 | -0.07 | 0.53 | 23.740 | 25.591 | 27.885 |
| <u>1e</u> | <u>p</u> -Cl | 5.07 | 0.23 | 0.75 | 23.847 | 25.721 | 28.096 |
| <u>1f</u> | <u>m</u> -Cl | 4.98 | 0.37 | 0.85 | 23.856 | 25.720 | 28.120 |
| <u>1g</u> | <u>p</u> -Br | 4.92 | 0.23 | 0.74 | - | - | - |
| <u>1h</u> | <u>m</u> -Br | 4.93 | 0.39 | 0.86 | - | - | - |
| <u>1i</u> | <u>m</u> -NO ₂ | 4.60 | 0.71 | 1.09 | 29.077 | 30.739 | 33.312 |
| <u>1j</u> | <u>p</u> -NO ₂ | 4.56 | 0.78 | 1.50 | 28.874 | 30.754 | 33.385 |

(a) Standard deviation $s = \pm 0.02-0.04$.

The results show that, to some extent, the acidity of 1 is influenced by the substituent in the 5-phenyl ring. Two correlations of the pK_a data with the simple Hammett equation⁸ were attempted. In the first case, the pK_a values were plotted against the Hammett substituent constants σ_X (Fig. 3), whereas in the second plot the phenyl group is included with the substituent and the pK_a values are correlated with the polar substituent constants $\sigma_{XC_6H_4}^*$ for the substituted phenyl groups.⁹




Fig. 3. Correlation of pK_a of 5-aryl-2-mercapto-1,3,4-oxadiazoles with the Hammett substituent constants.

The corresponding regression lines expressing such correlations are given by eqs. [3] and [4], respectively.

$$pK_a = 5.29 - 0.983\sigma_X \quad r = 0.959, \underline{s} = \pm 0.055 \quad [3]$$

$$pK_a = 5.83 - 0.985\sigma_{XC_6H_4}^* \quad r = 0.936, \underline{s} = \pm 0.123 \quad [4]$$

From the values of the correlation coefficient, r , and the standard deviation, \underline{s} , it is obvious that the pK_a values of $\underline{1}$ are correlated better by the Hammett substituent constants σ_X than by

the inductive constants $\sigma_{\text{XC}_6\text{H}_4}^*$ of the substituted phenyl group. This linear correlation between $\text{p}K_a$ and σ_X indicates further that the compounds in question exist predominantly in one tautomeric form regardless of the nature of the substituent on the 5-phenyl group. If the two forms 1 and 2 coexisted in equilibrium, a good correlation would not be observed.^{10,11} This is because for such an equilibrium (Scheme 2), the overall acid dissociation constant K_a is related to the dissociation constants K_1 and K_2 of the thiol and of the thioamide forms, respectively, and the tautomeric equilibrium constant K_T by eqs. [5] and [6]. According to these equations,

$$K_a = K_1 / (1 + K_T) \quad [5]$$

$$K_a = K_2 K_T / (1 + K_T) \quad [6]$$

a linear correlation between $\text{p}K_a$ and σ_X should be observed only if $K_1 = K_2$. In general, because of the ionization of the SH and NH groups, respectively, the values of K_1 and K_2 are different. The data from the correlation analysis make it possible to decide which of the possible tautomeric forms of mercaptooxadiazoles should be given preference. The reaction constant ($\rho = 0.983$) agrees better with the existence of the thioamide form 2. The absolute value of the reaction constant for the thiol tautomer 1 should be clearly much lower, because the length of the bridge between the substituent and the reaction site in 2 is shorter than in the tautomer 1. The assignment of the tautomeric structure 2 is compatible with the spectral data of the compounds studied. Thus, the infrared spectra of the compounds reveal no band in the 2550-2600 cm^{-1} region that could be assigned to the thiol tautomer 1.⁴ The spectra possess, however, in all cases, an absorption band close to 3100 cm^{-1} which can be assigned to an NH stretch, and a stretching frequency at 1152-1160 cm^{-1} attributable to the C=S bond. The NMR spectra do not possess a signal which would be due to a proton of an SH group; they contain signals in the 7.0-8.2 ppm range (aromatic protons) and, after exchange with D_2O , a signal appears at 4.65 ppm due to HDO and resulting from the NH + ND exchange. Furthermore, the electronic absorption pattern of 2a is identical to that of 5-phenyl-3-methyl-1,3,4-oxadiazole-2-thione 4 (Scheme 1).⁴ To assess further the relative stabilities of the two tautomeric forms 1 and 2 and of their common resonance-stabilized anion 3, their π -electronic energies, E_π , were calculated using the HMO method.⁶ The parameters used in the calculations are listed in Tables 2 and 3 and the results of the calculations are summarized in Table 1. In all cases it has been found that the thioamide form 2 has a higher π -electronic energy than the thiol tautomer 1. The value of 0.859β for the term $E_\pi(2) - E_\pi(1)$ found for compound 1a is similar to that (0.908β) reported⁴ for the same compound.

Table 2. HMO Heteroatom Parameters Used in the Calculations

| $\alpha_X = \alpha + h_X\beta$ (a) | | $\beta_{XY} = k_{XY}\beta$ (a) | |
|------------------------------------|-----------------|--------------------------------|--|
| SH form (1) | NH form (2) | Anion (3) | |
| $h_N(3) = 0.5$ | $h_N(3) = 1.5$ | $h_N(3) = 1.75$ | |
| $h_N(4) = 0.5$ | $h_N(4) = 0.5$ | $h_N(4) = 1.75$ | |
| $h_S = 1.0$ | $h_S = 0.6$ | $h_S = 0.8$ | |
| $h_O = 2.0$ | $h_O = 2.0$ | $h_O = 2.0$ | |
| $k_{CN} = 1.0$ | $k_{C=N} = 1.0$ | $k_{CN} = 0.9$ | |
| $k_{CO} = 0.8$ | $k_{C-O} = 0.8$ | $k_{CO} = 0.9$ | |
| $k_{NN} = 0.8$ | $k_{C-O} = 0.8$ | $k_{NN} = 0.9$ | |
| $k_{CS} = 0.7$ | $k_{NN} = 0.8$ | $k_{CS} = 0.8$ | |
| | $k_{CS} = 0.9$ | | |

(a) α_X is the Coulomb integral of the p_z atomic orbital of atom X; α is the same quantity for $2p_z$ atomic orbital of carbon; β_{XY} is the resonance integral of the XY bond and β is the same quantity for the CC bond in planar conjugated hydrocarbons.


Table 3. HMO Parameters for Substituents in the Phenyl Group

| Substituent, X | Coulomb integral, α_X | Resonance integral, β_{CX} |
|-------------------|--------------------------------------|-------------------------------------|
| Cl | $\alpha_{Cl} = \alpha + 2.0\beta$ | $\beta_{CCl} = 0.4\beta$ |
| NO ₂ | $\alpha_N = \alpha + \beta$ | $\beta_{CN} = \beta$ |
| | $\alpha_{C(N)} = \alpha + 0.2\beta$ | $\beta_{NO} = \beta$ |
| | $\alpha_O = \alpha + 1.5\beta$ | |
| Me (a) | $\alpha_{Me} = \alpha + 1.83\beta$ | $\beta_{CMe} = 0.6\beta$ |
| | $\alpha_{C(Me)} = \alpha - 0.2\beta$ | |
| MeO | $\alpha_{MeO} = \alpha + 2\beta$ | $\beta_{C-OMe} = 0.8\beta$ |

(a) Heteroatom model, see ref.¹²

These findings seem to substantiate the above conclusion that the compounds studied exist predominantly in the thioamide form $\tilde{2}$. Furthermore, the differences $[\Delta E_{\pi}(i) = E_{\pi}(\tilde{3}) - E_{\pi}(i)]$ between the π -electronic energy of the anion $\tilde{3}$ and that of each of the two tautomers indicate that the acidity of $\tilde{1}$ is higher than that of $\tilde{2}$. Because in acid-base equilibria of various tautomers the tautomer with higher acidity is considered to be less stable,¹³ it is not unreasonable to conclude that the NH tautomer $\tilde{2}$ is more stable than the SH form $\tilde{1}$. In all cases which we have studied, the structure of the anion $\tilde{3}$ is intermediate between the thiol $\tilde{1}$ and the thioamide $\tilde{2}$ forms as it can be seen from the molecular diagrams (Table 4). Also, if it is assumed that the changes in free energy due to entropy, ΔS , and energy of solvation of the anion $\tilde{3}$ and its conjugate

Table 4. Molecular Diagram for 5-(*p*-Tolyl)-2-mercapto-1,3,4-oxadiazole (a)



| Position, <i>i</i> | Electron densities, q_i | | | Bond, <i>ij</i> | Bond orders, p_{ij} | | |
|-----------------------|---------------------------|-------------|-------------|--------------------|-----------------------|-------------|-------------|
| | $\tilde{1}$ | $\tilde{2}$ | $\tilde{3}$ | | $\tilde{1}$ | $\tilde{2}$ | $\tilde{3}$ |
| 1 | 0.967 | 0.983 | 1.053 | 1-2 | 0.801 | 0.682 | 0.707 |
| 2 | 1.066 | 1.024 | 1.027 | 1-6 | 0.440 | 0.608 | 0.523 |
| 3 | 0.973 | 0.960 | 1.051 | 2-3 | 0.468 | 0.644 | 0.619 |
| 4 | 1.008 | 1.024 | 1.027 | 3-4 | 0.468 | 0.644 | 0.619 |
| 5 | 0.993 | 0.983 | 1.054 | 3-13 | 0.196 | 0.203 | 0.182 |
| 6 | 1.010 | 1.026 | 1.083 | 4-5 | 0.801 | 0.682 | 0.707 |
| 7 | 0.897 | 0.884 | 0.982 | 5-6 | 0.440 | 0.608 | 0.523 |
| 8 | 1.277 | 1.224 | 1.828 | 6-7 | 0.398 | 0.400 | 0.566 |
| 9 | 1.360 | 1.757 | 1.790 | 7-8 | 0.766 | 0.806 | 0.371 |
| 10 | 0.815 | 0.807 | 0.670 | 7-11 | 0.349 | 0.311 | 0.325 |
| 11 | 1.805 | 1.820 | 1.774 | 8-9 | 0.419 | 0.291 | 0.114 |
| 12 | 1.807 | 1.589 | 1.723 | 9-10 | 0.750 | 0.457 | 0.500 |
| 13 | 1.961 | 1.958 | 1.963 | 10-11 | 0.380 | 0.376 | 0.436 |
| | | | | 10-12 | 0.394 | 0.720 | 0.613 |

(a) Heteroatom model of the methyl group; see ref.¹²

acid are negligible in comparison with changes in the π -electronic energy due to dissociation, $\Delta E_{\pi}(i)$ can be approximated as

$$pK_a = a \Delta E_{\pi}(i) + b \quad [7]$$

where a and b are numerical constants. The correlation between the experimentally determined pK_a values and $\Delta E_{\pi}(3,2)$ [i.e., $E_{\pi}(3) - E_{\pi}(2)$] appears (Fig. 4) to be quite acceptable, particularly if the heterogeneity of the substituents as a group is taken into account. The equation of the regression line shown in Fig. 4 is

$$pK_a = 11.52 - 2.68 \Delta E_{\pi}(3,2); r = 0.971, s = \pm 0.086 \quad [8]$$

However, the values of $\Delta E_{\pi}(3,1)$ (i.e., $E_{\pi}(3) - E_{\pi}(1)$) for the thiol tautomers $\underline{1}$ do not correlate with the pK_a data. This result provides an additional support for the conclusion that the compounds examined in this study exist predominantly in the thioamide form $\underline{2}$ regardless of the nature of the substituent in the 5-phenyl group.

EXPERIMENTAL

The melting points were determined on a Gallenkamp electrothermal melting point apparatus and are uncorrected. Elemental analyses were carried out by the Microanalytical Laboratory of the University of Cairo, Giza, Egypt. Infrared spectra were recorded with a Beckman 200 infrared spectrophotometer, and the NMR spectra were taken on a Varian T60-A instrument. The δ values were obtained in $CDCl_3$ at room temperature, with TMS as the internal reference. The ultraviolet spectra of $\underline{1a-1j}$ were measured in ethanol on Pye-Unicam AR55 and Cary 118 spectrophotometers.

5-Aryl-1,3,4-oxadiazole-2-thiones, $\underline{1a-1j}$

They were prepared by a previously described method.^{7,14} The physical constants of the known compounds ($\underline{1a}$, $\underline{1b}$, $\underline{1d}$, $\underline{1e}$) were in agreement with the data published in the literature. The remaining compounds were new. They were purified by crystallization from ethanol until a constant melting point was obtained.

The characteristics of the synthesized compounds were as follows [compound no., substituent in the 5-phenyl ring, m.p. [m.p. lit. (ref.)], uv spectrum, λ^{EtOH} (log ϵ), molecular formula, anal. %S, found (calc.): $\underline{1a}$, H, 218°C [219-220°C (15)], 250 nm (4.05), 295 nm (4.32), $C_8H_6N_2OS$, - (-); $\underline{1b}$, p-MeO, 203°C [204-206°C (7)], 245 nm (3.90), 296 nm (3.98), $C_9H_8N_2O_2S$, - (-); $\underline{1c}$, p-Me, 223°C [-],




Fig. 4. Correlation of pK_a 's of 5-aryl-2-mercapto-1,3,4-oxadiazoles with ΔE_{π} .

240 nm (4.26), 300 (4.27), $C_9H_8N_2OS$, 16.59 (16.66); $\underline{1d}$, \underline{m} -Me, 160°C [158-160°C (16)], 247 nm (4.17), 295 nm (4.46), $C_9H_8N_2OS$, - (-); $\underline{1e}$, \underline{p} -Cl; 174°C [176-178°C (7)], 248 nm (4.50), 340 nm (4.16), $C_8H_5ClN_2OS$, - (-); $\underline{1f}$, \underline{m} -Cl, 179°C [-], 257 nm (4.11), 310 nm (4.17), $C_8H_5ClN_2OS$, 14.95 (15.06); $\underline{1g}$, \underline{p} -Br, 189°C [-], 240 nm (4.13), 305 nm (4.18), $C_8H_5BrN_2OS$, 12.35 (12.45); $\underline{1h}$, \underline{m} -Br, 212°C [-], 255 nm (4.12), 305 nm (4.17), $C_8H_5BrN_2OS$, 12.37 (12.45); $\underline{1j}$, \underline{m} -NO₂, 150°C [-], 224 nm (4.40), 330 nm (4.10), $C_8H_5N_3O_3S$, 14.25 (14.35); $\underline{1j}$, \underline{p} -NO₂, 200°C [-], 257 nm (4.35), 340 nm (4.16), $C_8H_5N_3O_3S$, 14.29 (14.35).

All compounds exhibit $\tilde{\nu}_{NH}$ (3031-3175 cm^{-1}) and $\tilde{\nu}_{CS}$ (1152-1160 cm^{-1}) bands in their ir spectra, and there is no $\tilde{\nu}_{SH}$ absorption at 2550-2600 cm^{-1} . The NMR spectra do not possess a signal due to an SH proton; they contain signals at 7.0-8.2 ppm (aromatic protons). After exchange with D₂O, a signal appears at 4.65 ppm (HDO; NH + ND exchange).

Determination of pK_a

The acid dissociation constants of $1a-1j$ were determined by both the potentiometric and the spectrophotometric methods in 80% (vol.) ethanol-water at 25°C. An Extech digital pH-meter type 671 fitted with an Orion combination glass electrode type 91-06 was used to measure the pH values. The instrument was calibrated by two standard Beckman buffer solutions with pH 4.01 and 10.00, respectively. The pH-meter readings B were converted to hydrogen ion concentration, $[H^+]$, by means of eq. [9].¹⁶

$$-\log[H^+] = \underline{B} + \log U_H \quad [9]$$

where $\log U_H = -0.4$ is the correction factor for the solvent composition and ionic strength for which B is read off at 25°C.

The procedures followed in the determination of the pK_a constants and their calculation from the spectrophotometric and potentiometric data have already been described.^{17,18} The pK_a values obtained by the two methods were in good agreement and were reproducible to within ± 0.03 pK_a unit. The results are summarized in Table 1.

HMO Calculations

The HMO calculations were carried out in the usual way on an IBM 360/65 computer.⁶ The values of the semiempirical parameters adopted in this work are given in Tables 2 and 3 along with those used for the corresponding anions.

ACKNOWLEDGEMENT

The financial support of the Robert A. Welch Foundation, Houston, Texas, through Grant No. AH-461 is greatly appreciated.

REFERENCES

1. A. Perjéssy, R. Frimm, and P. Hrnčiar, Collect. Czech. Chem. Commun., 1972, 37, 3302.
2. A. Perjéssy, P. Hrnčiar, and A. Krutošková, Tetrahedron, 1972, 28, 1025.
3. F. Fringuelli, G. Marino, and G. Savelli, Tetrahedron, 1969, 25, 5815.
4. J. Sandström and I. Wennerbeck, Acta Chem. Scand., 1966, 20, 57.
5. D. E. Horning and J. M. Muchowski, Can. J. Chem., 1972, 50, 3079.
6. A. Streitwieser, Jr., "Molecular Orbital Theory for Organic Chemists," J. Wiley, New York, N. Y. 1961.
7. E. Hoggarth, J. Chem. Soc., 1952, 4811.
8. O. Exner, in: "Advances in Linear Free Energy Relationships," N. B. Chapman, Ed., Plenum Press, New York, N.Y., 1972, Chapter 1.

9. Y. Nagai, H. Matsumoto, T. Nakano, and H. Watanabe, Bull. Chem. Soc. Jpn., 1972, 45, 2560.
10. M. I. Kabachnik, T. A. Mastyukova, A. E. Shipov, and T. A. Melent'eva, Tetrahedron, 1960, 9, 10.
11. W. L. Garbrecht and R. M. Herbst, J. Org. Chem., 1953, 18, 1003, 1269; R. M. Herbst and W. L. Garbrecht, J. Org. Chem., 1953, 18, 1283.
12. C. Párkányi, W. C. Herndon, and A. S. Shawali, J. Org. Chem., 1980, 45, 3529.
13. J. Arriau, J. M. Campillo, J. Elguero, and J. M. Pereillo, Tetrahedron, 1974, 30, 1345.
14. R. W. Young and K. H. Wood, J. Am. Chem. Soc., 1955, 77, 400.
15. M. Baron and C. V. Wilson, J. Org. Chem., 1958, 23, 1021.
16. L. G. Van Uitert and W. C. Fernelius, J. Am. Chem. Soc., 1954, 76, 5887.
17. A. Albert and E. P. Serjeant, "Ionization Constants of Acids and Bases," J. Wiley, New York, N. Y., 1962, p. 20.
18. A. S. Shawali and B. M. Altahou, Tetrahedron, 1977, 33, 1625.

Received, 6th May, 1983