

REACTIONS OF URACILS; 31

PYRAZOLO[3,4-d]- AND PYRIDO[2,3-d]PYRIMIDINES FROM 5-FORMYL-1,3-DIMETHYLURACILS

Péter Mátyus^{a2}, Pál Sohár^b, and Heinrich Wamhoff^{*a}^a Institut für Organische Chemie und Biochemie der Universität Bonn, Gerhard-Domagk-Str.1, D-5300 Bonn 1, Bundesrepublik Deutschland^b EGYT Pharmacochemical Works, H-1475 Budapest, P.O.Box 100, Hungary


Abstract — Starting from the known 5-formyl-1,3-dimethyluracils (1a,b) some novel derivatives having a great biological interest have been synthesized. The constitutions of the products obtained were established based on spectral data.

The biological activities of pyrimidines containing acyl and/or olefinic functions at position-5 have stimulated considerable research in this field³⁻⁵. Continuing our work on the synthesis of uracils of potential biological activity¹, we report here some reactions of 5-formyl-1,3-dimethyluracils to new derivatives and the synthesis of several, otherwise hardly accessible new 2,4,7-trioxypyrido[2,3-d]pyrimidines. It has been reported that the reaction of 1a with phenyl- or methylhydrazine yielded the Schiff's bases which upon heating gave N-substituted pyrazolo[3,4-d]-pyrimidines⁶. We have extended this reaction to (6-chloro-3-pyridazinyl)-hydrazine 2 in order to investigate the influences of the heteroaromatic ring and the adjacent ring N-atoms on the stability of the intermediate as well as on the formation of pyrazolo[3,4-d]pyrimidine derivative. Thus, 1a condensed with 2 to afford the arylidene derivative 3, which reacted smoothly with aqueous methylamine to give 4. In contrast to the behaviour of the phenyl analogue, the intramolecular cyclization tendency of 3 was considerably low due to the -I effect of the 3-pyridazinyl ring: only the thermal reaction at 180°C of 3 afforded 5 in moderate yield, while the cyclization was unsuccessful in ethanol in the presence of equimolar amounts of triethylamine. On the basis of spectral data the formation of the theoretically also possible tricycle 6 could be ruled out. The E-3-(5-uracilyl)acrylic acid derivatives 7a,b were synthesized by Wittig or Knoevenagel-Doebner reaction, respectively.


An investigation was undertaken to explore the potential utility of the reactions

of some nucleophiles with 7a as a route for the synthesis of fused ring systems. It has been found that under kinetic control primary aliphatic amines or ammonia react with 7a to yield 8a-c. Upon heating of 8a,b in triethylamine in the presence of catalytic amount of 1,5-diazabicyclo[4.3.0]non-5-ene (DBN), the novel 2,4,7-trioxopyrido[2,3-d]pyrimidines 9a,b were obtained in high yield.

Since the alkylation of 1,3-dimethyl-2,4,7-trioxopyrido[2,3-d]pyrimidine 10⁸ with butyl bromide (or iodide) leads to the formation of the 7-butoxy derivative 11, our method provides a simple and efficient approach to the synthesis of the potentially biological active 8-substituted 2,4,7-trioxopyrido[2,3-d]pyrimidines.



In contrast to the behaviour of 8a toward the triethylamine/DBN system, it reacted in Dowtherm[®]A at 200°C to the desalkylated derivative 12, presumably *via* two consecutive sigmatropic 1,5-H shifts:



The constitution of 12 was established based on spectral data. Thus, in the ¹H NMR spectrum the absence of an AB quartet being characteristic for the olefinic protons and the lack of the butyl protons, as well as the presence of -(CH₂)₂- and NH₂ protons pointed unambiguously to the constitution of 12. Moreover, also IR, MS, and ¹³C NMR⁹ data were of further evidence for this structural proposal. Now, the behaviour of several other mono- and bifunctional nucleophiles towards 7a as well as the mechanism of formation of 12 is under investigation. Work in progress together with the full preparative details will be published elsewhere¹⁰.

Table 1. List of compounds 3-5, 7a,b, 8a-c, 9a,b, 11, and 12[†]

Comp.	Solvent of Cryst.	M p [°C]	Yield [%]	Molecular Formula
<u>3</u>	ethanol	224-225	90	C ₁₁ H ₁₀ Cl ₂ N ₆ O ₂
<u>4</u>	ethanol	313-314	84	C ₁₂ H ₁₄ ClN ₇ O ₂
<u>5</u>	ethanol	269-270	41	C ₁₁ H ₉ ClN ₆ O ₂
<u>7a</u>	ethanol	174-175	72	C ₁₁ H ₁₃ ClN ₂ O ₄
<u>7b</u>	ethanol	281-282	78	C ₉ H ₁₀ N ₂ O ₄
<u>8a</u>	i-propanol	118-119	54	C ₁₅ H ₂₃ N ₃ O ₄
<u>8b</u>	i-propanol	170-171	54	C ₁₂ H ₁₇ N ₃ O ₄
<u>8c</u>	ethanol	237-238	61	C ₁₁ H ₁₅ N ₃ O ₄
<u>9a</u>	ether	99-100	80	C ₁₃ H ₁₇ N ₃ O ₃
<u>9b</u>	ethanol	209-210	56	C ₁₀ H ₁₁ N ₃ O ₃
<u>11</u>	i-propanol	91-92	68	C ₁₃ H ₁₇ N ₃ O ₃
<u>12</u>	i-propanol	163-164	48	C ₁₁ H ₁₇ N ₃ O ₄

[†]Satisfactory elemental analyses (C,H,N) and MS data were obtained for all the newly synthesized compounds.

Table 2. IR and ¹H NMR data of compounds 3-5, 7a,b, 8a-c, 9a,b, 11, and 12

Comp.	IR (KBr, [cm ⁻¹])	¹ H NMR ([D ₆]DMSO, δ [ppm])
<u>3</u>	1715 (CO), 1670 (CO), 1615 (C=C);	3.22 (s, 3H, 1-N-CH ₃), 3.53 (s, 3H, 3-N-CH ₃), 7.47 (d, (J=10 Hz), 1H, 4'-CH), 7.63 (d (10), 1H, 5'-CH), 8.16 (s, 1H, CH=N), 11.7 (s, 1H, NH);
<u>4</u>	3205 (NH), 1695 (CO), 1660 (CO), 1615 (C=C);	3.12 (d (6), 3H, NHCH ₃) ⁺ , 3.16 (s, 3H, 1-N-CH ₃) ⁺ , 3.33 (s, 3H, 3-N-CH ₃) ⁺ , 7.24 (d (10), 1H, 4'-CH), 7.56 (d (10), 1H, 5'-CH), 8.42 (s, 1H, CH=N), 9.3 (broad, 1H, NH-CH ₃), 11.35 (s, 1H, N-NH);
<u>5</u>	1720 (CO), 1665 (CO), 1615 (C=C);	3.24 (s, 3H, 1-N-CH ₃), 3.47 (s, 3H, 3-N-CH ₃), 8.18 (d, (9), 1H, 4'-CH), 8.29 (d (9), 1H, 5'-CH), 9.33 (s, 1H, CH=N);
<u>7a</u>	1715 (CO) ⁺ , 1710 (ester CO) ⁺ , 1670 (CO), 1295, 1175 (C-O);	1.22 (t (7.5), 3H, 11-CH), 3.22 (s, 3H, 1-N-CH ₃), 3.57 (s, 3H, 3-N-CH ₃), 4.17 (q, 2H, 10-CH), 7.10 (d (16), 1H, 8-CH), 7.58 (d (17), 1H, 7-CH);

Table 2. — Continued

Comp.	IR(KBr, [cm ⁻¹])	¹ H NMR ([D ₆]DMSO, δ [ppm])
<u>7b</u>	1730 (CO), 1695 (CO), 1645 (C=C);	3.00 (s, 3H, 1-N-CH ₃), 3.16 (s, 3H, 3-N-CH ₃); 6.56 (d, (16), 1H, 8-CH), 7.11 (d (16), 1H, 7-CH), 8.13 (s, 1H, 6-CH);
<u>8a</u>	3345 (NH), 1705 (CO), 1675 (CO);	0.87 (t (6.5), 3H, CH ₃ (Bu)), 1.22 (t (7), 3H, OCH ₂ CH ₃ (Bu)) ⁺ , 1.4 (m, 2H, CH ₂ CH ₃ (Bu)) ⁺ , 1.7 (m, 2H, NHCH ₂ CH ₂ -), 3.16 (s, 3H, 1-N-CH ₃), 3.3 (m, 2H, NHCH ₂) ^o , 3.33 (s, 3H, 3-N-CH ₃) ^o , 4.11 (q, 2H, OCH ₂), 6.73 (d (15), 1H, 8-CH), 7.50 (d (15), 1H, 7-CH);
<u>8b</u>	3360 (NH), 1705 (CO);	1.21 (t (7), 3H, CH ₂ CH ₃), 3.00 (d, 3H, NHCH ₃), 3.16 (s, 3H, 1-N-CH ₃), 3.36 (s, 3H, 3-N-CH ₃), 4.12 (q, 2H, CH ₂ CH ₃), 6.69 (d (15), 1H, 8-CH), 7.00 (broad, 1H, NH), 7.62 (d (15), 1H, 7-CH);
<u>8c</u>	3380, 3250 (NH), 1690 (CO), 1670 (CO);	1.22 (t (7), 3H, CH ₂ CH ₃), 3.14 (s, 3H, 1-N-CH ₃), 3.36 (s, 3H, 3-N-CH ₃), 4.13 (q, 2H, CH ₂ CH ₃), 6.96 (d (15), 1H, 8-CH), 7.67 (d (15), 1H, 7-CH);
<u>9a</u>	1715 (CO), 1665 (CO);	0.84 (t (7), 3H, CH ₂ CH ₃), 1.32 (sx, 2H, CH ₂ CH ₃), 1.68 (qui, 2H, NCH ₂ CH ₂ -), 3.22 (s, 3H, 1-N-CH ₃), 3.50 (s, 3H, 3-N-CH ₃), 4.16 (t (7.5), 2H, N-CH ₂ -), 6.29 (d (9), 1H, 6-CH), 7.85 (d (9), 1H, 5-CH);
<u>9b</u> ⁺⁺	1715 (CO), 1660 (CO);	3.38 (s, 3H, 3-N-CH ₃), 3.57 (s, 3H, 1-N-CH ₃), 3.59 (s, 3H, 8-N-CH ₃), 6.37 (d (9), 1H, 6-CH), 7.95 (d (9), 1H, 5-CH);
<u>11</u> ⁺⁺	1710 (CO), 1665 (CO);	0.98 (t (7), 3H, CH ₂ CH ₃), 1.44 (m, 2H, CH ₂ CH ₃), 1.80 (m, 2H, O-CH ₂ CH ₂ -), 3.46 (s, 3H, 1-N-CH ₃), 3.67 (s, 3H, 3-N-CH ₃), 4.44 (t, 2H, -O-CH ₂ -), 6.58 (d (9), 1H, 6-CH), 8.29 (d (9), 1H, 5-CH);
<u>12</u> ⁺⁺	3340, 3360 (NH), 1705 (CO), 1680 (CO);	1.24 (t (7), 3H, 11-CH), 2.63 (s, 4H, 7,8-CH), 3.32 (s, 3H, 1-N-CH ₃), 3.44 (s, 3H, 3-N-CH ₃), 4.10 (q, 2H, 10-CH), 5.32 (s, 2H, NH ₂);

^{+,o} superimposed signals

⁺⁺ ¹H NMR spectrum in CDCl₃ solution.

ACKNOWLEDGEMENT This work was supported by the *Fonds der Chemischen Industrie*, the *Minister für Wissenschaft und Forschung des Landes Nordrhein-Westfalen*, and the *BAYER AG.* - P.M. thanks the *Alexander von Humboldt-Stiftung* for granting a fellowship.

REFERENCES AND NOTES

1. Part 2: G. Szilágyi and H. Wamhoff, Angew.Chem., 1983, 95, 156; Angew.Chem. Suppl., 1983, 120; Angew.Chem.Int.Ed.Engl., 1983, 22, 161; cf. also H. Wamhoff, "Some Model Reactions for Studying Electronic, Steric, and Other Effects on the Photocycloaddition to Nucleic Acids" in G. Náray-Szabó (ed), *Steric Effects in Biomolecules*, Akadémiai Kiadó, Budapest 1982, p. 195.
2. Alexander von Humboldt-Fellow 1982/83.
3. J.P. Jonak, S.F. Zakrzewski, and L.H. Mead, J.Med.Chem., 1971, 14, 408.
4. J.D. Fissekis and F. Sweet, J.Org.Chem., 1973, 28, 264.
5. V. Warin, M. Foulon, F. Baert, J.L. Bernier, and J.P. Henichart, Acta Cryst. 1980, B36, 1721.
6. S. Senda, K. Hirota, and G.N. Yang, Chem.Pharm.Bull. (Japan), 1972, 20, 399.
7. 1b was obtained in 84% yield from the Vilsmeier reaction of 1,3-dimethyluracil.
8. A.D. Broom, J.L. Shim, and G.L. Anderson, J.Org.Chem., 1976, 41, 1095.
9. ¹³C NMR data of 12 (CDCl₃, δ, [ppm]): 15.7 (C-11), 20.9 (C-8), 29.0 (C-12), 31.4 (C-13), 34.0 (C-7), 61.2 (C-10), 85.8 (C-5), 152.6 (C-2), 153.2 (C-4), 163.1 (C-6), 174.5 (C-9).
10. P. Mátyus, P. Sohár, and H. Wamhoff, Synthesis, in preparation.

Received, 24th August, 1983