

1-ACYLOXYPYRIDINIUM ION: THE REACTIVE INTERMEDIATE IN A MODIFIED REISSERT-HENZE REACTION

Wilmer K. Fife^{*1}

Indiana University-Purdue University at Indianapolis

Indianapolis, IN 46223

and


Brett D. Boyer²

Indiana Central University

Indianapolis, IN 46227


Abstract - Cyanation of 3-X-1-dimethylaminocarbonyloxypyridinium ions with trimethylsilane-carbonitrile or cyanide ion gives 3-methyl-2-pyridinecarbonitrile (~90%) contaminated with 5-methyl-2-pyridinecarbonitrile (~10%) when X = -CH₃, and approximately equal amounts of the 3- and 5-X derivatives when X = -COOCH₃. These product mixtures are identical to those obtained with the corresponding pyridine 1-oxides and dimethylcarbamoyl chloride in a modified Reissert-Henze reaction.


The recently reported quantitative conversion of pyridine 1-oxides to 2-pyridinecarbonitriles³ has now been shown to involve the intermediacy of 1-acyloxypyridinium ion, eq. 1. 1-Acyloxypyridinium ions have been



presumed to be the reactive intermediate in the few documented examples of Reissert-Henze cyanation of pyridine 1-oxides in the presence of benzoyl chloride.⁴ Treatment of the 1-dimethylaminocarbonyloxypyridinium ion,⁵ **1**, formed by prior reaction between a pyridine 1-oxide and dimethylcarbamoyl chloride, with either trimethylsilanecarbonitrile in dichloromethane or potassium cyanide in acetonitrile (or water) furnishes the related 2-pyridinecarbonitrile, **2**, in essentially quantitative yield (Table 1).

Table 1. Cyanation of Pyridine Derivatives^a






(Yields %)^b

X	Y	Cyanating Agent	Solvent	3-X 2-CN	5-X 2-CN	3-X 4-CN	Overall Isolated Yield (%)
H ³	O ⁻	Me ₃ SiCN/Me ₂ NCOCI	CH ₂ Cl ₂	100	—	0	94
H	OOCNMe ₂	Me ₃ SiCN	CH ₂ Cl ₂	100	—	0	95
		KCN	CH ₃ CN	100	—	0	90
		KCN	H ₂ O	100	—	0	88
CH ₃ ³	O ⁻	Me ₃ SiCN/MeNCOCI	CH ₂ Cl ₂	90	10	0	95
CH ₃	OOCNMe ₂	Me ₃ SiCN	CH ₂ Cl ₂	94	6	0	91
		KCN	CH ₃ CN	86	14	0	90
		KCN	H ₂ O	89	11	0	87
COOCH ₃ ⁶	O ⁻	Me ₃ SiCN/Me ₂ NCOCI	CH ₂ Cl ₂	40	60	0	70
COOCH ₃	OOCNMe ₂	Me ₃ SiCN	4:1 DMSO-d ₆ - CDCl ₃ ^c	40	60	trace	—
		KCN	CH ₃ CN	48	42	10	90
		KCN	D ₂ O	55	43	2	—
CH ₃	OCH ₃	Me ₃ SiCN	1:1 DMSO-d ₆ - CDCl ₃	0	0	0	0
		KCN	CH ₃ CN	22	5	73	50
		KCN	H ₂ O	24	7	69	65
COOCH ₃	OCH ₃	Me ₃ SiCN	1:1 DMSO-d ₆ - CDCl ₃	0	0	0	0
		KCN	CH ₃ CN	16	26	58	60
		KCN	H ₂ O ^d	2	11	87	75

^aAnalytical experiments utilized reaction mixtures at room temperature (~27°C) that contained 2.5x10⁻⁴ mole pyridinium salt, 2.8x10⁻⁴ mole trimethylsilanecarbonitrile or 7.5x10⁻⁴ mole potassium cyanide in 0.50 mL solvent. Preparative experiments were conducted with the same proportions of reactants and 0.005-0.010 mole pyridinium salt. See reference 3 for experimental details. ^bProduct compositions were determined by VPC analysis (OV-1 5% on Chromosorb W) at 160°C. All products were compared to samples that gave melting points and/or IR and NMR spectra that were in excellent agreement with literature data, or had appropriate elemental compositions. ^c3-Carbomethoxy-1-dimethylaminocarbonyloxypyridinium chloride reverts to a mixture of the N-oxide and dimethylcarbamoyl chloride in chloroform-d. ^dThe reported result was obtained by addition of an aqueous solution containing 10-fold excess potassium cyanide to an aqueous solution containing the pyridinium salt. A reversed order of addition produced the cyanation product mixture: 11% 2-CN, 18% 6-CN, 71% 4-CN (overall isolated yield of cyano product-70%).

Product mixtures obtained by cyanation of 1 ($X = 3\text{-CH}_3$ and 3-COOCH_3) are identical to those obtained by treatment of related pyridine 1-oxides with trimethylsilanecarbonitrile and dimethylcarbonyl chloride,³ and they differ markedly from product mixtures derived from the corresponding 1-methoxypyridinium ions, 3, eq. 2. 1-Methoxypyridinium ions are not cyanated with trimethylsilanecarbonitrile in dichloromethane,



but they react with cyanide ion to give high percentages of 4 in agreement with the literature.⁷


In contrast, products derived from 1, are almost exclusively the result of cyanation at a ring position adjacent to nitrogen. The only exception is the cyanation of the 3-COOCH_3 derivative with potassium cyanide in acetonitrile which gave 4 in 10% yield along with nearly equal amounts (48% and 42%, respectively) of 3- and 5-carbomethoxy-2-pyridinecarbonitrile.

Table 2. Relative Reactivity of 3-X-Pyridine 1-Oxide and 3-X-1-Dimethylaminocarbonyloxypyridinium Ion to Cyanation^a


Pyridine Derivative	Cyanating Conditions	$t_{1/2}$	Rel. Rate
X = H	$\text{Me}_3\text{SiCN/Me}_2\text{NCOCl/CDCl}_3$	15 min	20
X = 3-CH_3		20 min	15
X = 3-COOCH_3		5 hours	1
	X = 3-CH_3 $\text{Me}_3\text{SiCN/4:1 DMSO-d}_6\text{-CDCl}_3$	1 min	300
X = 3-COOCH_3		1 min	300

^aReactions were carried out at ambient probe temperatures ($\sim 32^\circ\text{C}$) in an NMR tube with equimolar mixtures (2.5×10^{-4} mole) of reactants in 0.50 mL solvent. The 3-carbomethoxy-1-dimethylaminocarbonyloxypyridinium ion was studied in 4:1 $\text{DMSO-d}_6\text{-CDCl}_3$ to avoid equilibration with N-oxide and dimethylcarbonyl chloride. Half-lives ($t_{1/2}$) were estimated by integration of NMR signals of reaction mixtures.

The relative reactivity of 1 with trimethylsilanecarbonitrile is 15–300X that of related N-oxide-dimethylcarbonyl chloride mixtures (Table 2). Furthermore, the reactivity of the 1-acyloxypyridinium ions is relatively insensitive to substituent, whereas the order of reactivity of the N-oxides toward cyanation parallels their ability to be acylated ($3\text{-CH}_3 > \text{H} > \text{-COOCH}_3$).⁸ The inertness of 3 to trimethylsilanecarbonitrile and the appreciable amounts of product produced by attack of cyanide ion at the 4-position of 3 strongly supports the intervention of an intermediate such as 5 or 6 in reactions of 1.⁹



5



6

Further work is currently underway to more fully characterize this reaction. Related studies in progress show that 1-acyloxy pyridinium ions are extremely versatile intermediates for the synthesis of a wide-variety of pyridine derivatives. The results of these studies will be reported at a later date.

ACKNOWLEDGEMENT This research was supported by the gift of starting materials from Reilly Tar and Chemical Corporation.

REFERENCES AND NOTES

1. Author to whom correspondence should be addressed.
2. An undergraduate research participant under the Consortium for Urban Education, Indianapolis. Portions of this report were taken from the B.Sc. Thesis of B. D. Boyer.
3. W. K. Fife, *J. Org. Chem.*, 1983, 48, 1375.
4. a) E. Ochiai and I. Nakayama, *Yakugaku Zasshi*, 1945 65, 582. b) Y. Kobayashi and I. Kumadaki, *Chem. Pharm. Bull.*, 1969 17, 510-514.
5. The preparation of 1-dimethylaminocarbonyloxy pyridinium ions has been reported previously by: a) P. Bergthaller, *Ger. Offen.* 2, 408, 813 (Cl. C07D), Sept. 4, 1975, 25 pp., *Chem. Abstr.*, 84, P43859p (1976); c) P. Bergthaller, W. Himmelmann, W. Sauerteig, and L. Rosenhahn, *Ger. Offen.* 2, 408, 814 (Cl. G03C), Sept. 4, 1975, 37 pp., *Chem. Abstr.*, 84, P67803s (1975).
6. W. K. Fife, *Heterocycles*, in press.
7. E. Ochiai, "Aromatic Amine Oxides," Elsevier Publishing Company, Amsterdam, 1974, p. 269. A. R. Katritzky and J. M. Lagowski, "Chemistry of the Heterocyclic N-Oxides," Academic Press, London, 1971, p. 300. R. A. Abramovitch and E. M. Smith, "The Chemistry of Heterocyclic Compounds," A. Weissberger and E. C. Taylor, eds., John Wiley and Sons, New York, Vol. 14, Suppl. Part 2, 1974, pp. 114-117.
8. Studies currently in progress confirm that a wide variety of substituted 1-dimethylaminocarbonyloxy pyridinium ions react rapidly with either trimethylsilanecarbonitrile or cyanide ion ($t_{1/2}$ 5 min. or less). Similar reactions with related pyridine-1-oxides and dimethylcarbamoyl chloride have half-lives that vary from 0.2 to 1.5×10^3 hours.
9. Complexes similar to 6 have been invoked in recent studies of silicon compounds. See R.J.P. Corriu and C. Guerin, *Adv. Organomet. Chem.*, 1982 20, 265.

Received, 12th December, 1983