

SYNTHESIS OF 3-HYDROXYETHYL-4-OXOAZETIDIN-2-YLPHOSPHONATE DERIVATIVES:
POTENTIAL PRECURSORS TO CARBAPENEM AND α -AMINOPHOSPHONIC ACID DERIVATIVES

Masao Shiozaki and Hidekazu Masuko

Chemical Research Laboratories, Sankyo Co., Ltd.

Hiromachi 1-2-58, Shinagawa-ku, Tokyo 140, Japan


Abstract - Optically active 3-hydroxyethyl-4-oxoazetidin-2-ylphosphonate derivatives were synthesized.

4-Oxoazetidin-2-ylphosphonate derivatives, which should be potential precursors to the carbapenems, obtained from treatment of the corresponding 4-acetoxiazetidin-2-one derivatives with trialkyl phosphites have already been reported by Campbell¹ as the starting material of alanyl dipeptide containing organophosphorous moiety via β -phosphono- β -alanine derivatives. Such organophosphorous-containing molecules as the α -aminophosphonic acid dipeptide (for example: alaphosphine - an inhibitor of alanine racemase) are becoming increasingly important for pharmaceutical use. In this paper we wish to describe a new synthetic procedure for 4-oxoazetidin-2-ylphosphonate derivative by the ring closure of 2-bromo-3-hydroxybutyramides via the corresponding 2,3-epoxybutyramides which possess an activated methylene adjoining the amide nitrogen.

The starting phosphorous-containing butyramides were synthesized as follows: Treatment of anisidine or *p*-methoxybenzylamine with paraformaldehyde according to the procedure reported by Ratcliffe et al.² gave hexahydrotriazines, 1a (72%; mp 128-130° C) or 1b (95%; mp 113-115°C). Reaction of the hexahydrotriazines (1a and 1b) with diethyl phosphite at 100°C for 4-6 h gave secondary amines (2a, 61% and 2b, 65%). Amide formation reaction of the amines (2a and 2b) with 1 eq of (2S,3R)-2-bromo-3-hydroxybutyric acid (3)³ in THF in the presence of 1 eq of DCC gave 4a (21%) and 4b (77%).

Next, these phosphorous-containing butyramides were cyclized to β -lactams in a stereospecific manner.⁴ Treatment of 4a and 4b with 1 eq of lithium hexamethyldisilazide in THF at ice-cooling temperature for 5 min, and then another 1 eq of $\text{LiN}(\text{SiMe}_3)_2$ at 24°C gave chiral diethyl 3-hydroxyethyl-4-oxoazetidin-2-ylphosphonate derivatives, 5a (61%) and 5b (37%), and *cis*-isomers were not isolated. Protection of the hydroxy group of 5a with *p*-nitrobenzylchloroformate by use of 4-dimethylamino-

pyridine as a base in CH_2Cl_2 gave 6a (79%) as a foam. Deprotection of the methoxyphenyl group of 6a with 3 eq of cerium (IV) ammonium nitrite (CAN) in acetone- H_2O (3:2) gave 7A (81%) as a crystalline solid; mp $74-76^\circ\text{C}$. Hydrogenolysis of 7A with 5% Pd-C/ H_2 at room temperature gave 7C (92%); mp $70-71^\circ\text{C}$; $[\alpha]_{\text{D}}^{24} -4.9^\circ$ ($c=0.59$, CHCl_3). Similarly, protection of the hydroxy group of 5b with *t*-butyldimethylsilyl chloride by use of 4-dimethylaminopyridine as a base in DMF gave 6b (67%). Deprotection of the methoxybenzyl group of 6b with $\text{K}_2\text{S}_2\text{O}_8$ - K_2HPO_4 in MeCN- H_2O (1:1) at $70-75^\circ\text{C}$ under argon for 30 min gave N-free β -lactam 7B (57%) as a gum. These compounds (5a,b, 6a,b and 7A,B,C) should be potential precursors to new carbapenems and α -aminophosphonic acid analogues of aspartic acid.



a: R=p-Methoxyphenyl



1a

2a

b: R=p-Methoxybenzyl

1b

2b



7A, $\text{R}^1=\text{p-Nitrobenzyloxycarbonyl}$

7B, $\text{R}^1=\text{t-Butyldimethylsilyl}$

7C, $\text{R}^1=\text{H}$

REFERENCE

1. M. M. Campbell and N. Carruthers, *J. Chem. Soc., Chem. Commun.*, 1980, 730.
2. R. W. Ratcliffe and B. G. Christensen, *Tetrahedron Letters*, 1973, 4645.
3. Y. Shimohigashi, M. Waki and N. Izumiya, *Bull. Chem. Soc. Japan*, 1979, 52, 949.
4. M. Shiozaki, N. Ishida, T. Hiraoka and H. Maruyama, *Tetrahedron Letters*, 1981, 5205.

Received, 1st May, 1984