

SYNTHESIS AND CRYSTAL STRUCTURE OF DECAHYDRO-DISPIRO[OXIRANE-2,3'-(2'H)BENZOFURAN-2',2"-[2H]PYRAN]

Chuzo Iwata*, Katsuya Nakamura, Hiroshi Atarashi, and Shuji Uchida
Faculty of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565, Japan

Masaru Kido

Laboratory of Natural Products Chemistry, Ohtsuka Pharmaceutical Co. Ltd., Kawauchi-cho, Tokushima 770, Japan



Abstract — (2'S*,3'R*,3'aS*,7'aR*)-Decahydro-dispiro[oxirane-2,3'-(2'H)benzofuran-2',2"-[2H]pyran] (E-6) and its isomer E-6' with epimeric stereochemistry of the epoxide ring were prepared stereoselectively and an X-ray analysis of E-6' is reported.

In the preceding paper,¹ we described an efficient route of E- and Z-1,6-dioxaspiro-[4.5]decane ring systems, important moieties of many natural products including insect pheromones and polyether antibiotics.

In this paper, we report the synthesis of dispiro compounds, E-6 and its isomer E-6', the essential framework of phyllanthocin 1², and an X-ray analysis of E-6' which confirms the correctness of our previous proposal about the stereochemistry of a series of compounds 2, 3, 4 and 5.

Though various attempts³ to introduce the epoxy function into the E-olefin 4 were unsuccessful, resulting in recovery of the starting material or decomposition, epoxidation in conventional way (MCPBA, CH₂Cl₂, 0°C, 48 h) afforded the two diastereomeric isomers, E-6 [the minor isomer (5.6%); bp 100°C/0.01 mmHg; δ_H(CCl₄): 2.68 and 2.76 (each 1H, d, J=5 Hz), 3.30-3.58 (1H, m), 3.62-3.89 (1H, m), 4.15 (1H, m); δ_C(CDCl₃): 38.8(d), 61.8(t), 70.2(s), 73.5(d), 103.2(s)] and E-6' [the major isomer (84%); mp 57.5-58°C; δ_H(CCl₄): 2.58 and 2.62 (each 1H, d, J=5 Hz), 3.40-3.98 (2H, m), 4.23 (1H, m); δ_C(CDCl₃): 40.5(d), 61.9(t), 70.5(s), 73.0(d), 100.7(s)].

On the other hand, this minor isomer E-6 was derived from E-ketone 5 with high selectivity in a two-step sequence (46% overall yield): the formation of β-hydroxy-sulfide E-7 [PhSCH₂Li⁴ (5 equiv.), THF, r.t., 15 h] and methylation at the sulfur with



trimethyloxonium tetrafluoroborate (CH_2Cl_2 , r.t., 1 h) followed by treatment of aq. base (5% NaOH, r.t., 15 h).⁵

An attempt to determine the stereochemistry of these isomers by the NMR spectra was uninformative. Thus, in order to establish these structures and ensure our previous stereochemical assignments of a series of the synthetic intermediates 2, 3, 4 and 5 based on the ^{13}C NMR spectral data, the nicely crystalline compound E-6' was subjected to an X-ray analysis. The crystal data and intensity data were derived from the measurements on a Syntex R3 four-circle diffractometer with graphite-monochromated $\text{MoK}\alpha$ radiation. Crystal data: $\text{C}_{13}\text{H}_{20}\text{O}_3$, monoclinic, space group $\text{P2}_1/\text{c}$, $a=11.770(4)$, $b=9.573(4)$, $c=10.769(7)\text{\AA}$, $\beta=92.01(4)^\circ$, $D_x=1.23\text{ g.cm}^{-3}$, $z=4$ and $\mu(\text{MoK}\alpha)=0.9\text{ cm}^{-1}$. Intensity data for 1293 reflections ($I>1.96\sigma(I)$) were collected using an ω -scan mode within 2θ less than 45° . Lorentz and polarization corrections were applied, but no absorption corrections were made. The structure was solved by the direct method using MULTAN on a Syntex XTL program. All the hydrogen atoms were found on a difference Fourier map. The refinement of atomic parameters was carried

out by block-diagonal least-squares calculations. The final R-value was 0.041 assuming anisotropic thermal motions for non-hydrogen atoms and isotropic ones for hydrogen atoms. The molecular structure is illustrated in Figure. Once the structure of E-6' was established, the structures of the synthetic intermediates 2, 3, 4 and 5 then logically followed.

In conclusion, we have achieved a stereoselective synthesis of dispiro compound E-6 which possesses the requisite stereochemistry among the tetracyclic rings contained in phyllanthocin 1. Further synthetic approach toward phyllanthocin and related compounds is in progress.




Figure. The molecular structure of E-6'

REFERENCES AND NOTES

1. C.Iwata, H.Atarashi, K.Nakamura, and S.Uchida, submitted for publication in Heterocycles.
2. S.M.Kupchan, E.J.LaVoie, A.R.Branfman, B.Y.Fei, W.M.Bright, and R.F.Bryan, J. Am. Chem. Soc., 1977, 99, 3199; P.R.McGuirk and D.B.Collum, ibid., 1982, 104, 4496, J. Org. Chem., 1984, 49, 843.
3. For example: G.B.Payne, P.H.Deming, and P.H.Williams, J. Org. Chem., 1961, 26, 659; W.Cocker and D.H.Grayson, Tetrahedron Lett., 1969, 4451; R.C.Carlson and R.Ardon, J. Org. Chem., 1971, 36, 216; K.B.Sharpless and R.C.Michaelson, J. Am. Chem. Soc., 1973, 95, 6136; M.Parrilli, G.Barone, M.Adinolfi, and L.Mangoni, Tetrahedron Lett., 1976, 207; T.Yamamoto and M.Kimura, Chem. Comm., 1977, 948;

- D.R.Williams, B.A.Barner, K.Nishitani, and J.G.Phillips, J. Am. Chem. Soc., 1982, 104, 4708.
4. E.J.Corey and D.Seebach, J. Org. Chem., 1966, 31, 4097.
5. J.R.Shanklin, C.R.Johnson, J.Ollinger, and R.M.Coates, J. Am. Chem. Soc., 1973, 95, 3429.

Received, 28th May, 1984